

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):


- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.

(51) International Patent Classification <sup>6</sup> : C12N 15/34, 15/86, 15/85, 5/10, C07K 14/075, A61K 48/00		A2	(11) International Publication Number: <b>WO 98/13499</b>
		(43) International Publication Date: 2 April 1998 (02.04.98)	
(21) International Application Number: PCT/EP97/05251		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 24 September 1997 (24.09.97)		<p><b>Published</b> Without international search report and to be republished upon receipt of that report.</p>	
(30) Priority Data: 08/719,806 25 September 1996 (25.09.96) US			
(71) Applicants (for all designated States except US): NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH). THE SCRIPPS RESEARCH INSTITUTE [US/US]; 10666 North Torrey Pines Road, La Jolla, CA 92037 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): MEMEROW, Glen, R. [US/US]; 462 Cerro Street, Encinitas, CA 92024 (US). VON SEGGERN, Daniel, J. [US/US]; 7528 Charmant Drive #428, San Diego, CA 92122 (US).			
(74) Agent: ROTH, Bernhard, M.; Novartis AG, Patent- und Markenabteilung, Lichtstrasse 35, CH-4002 Basel (CH).			

(54) Title: PACKAGING CELL LINES FOR USE IN FACILITATING THE DEVELOPMENT OF HIGH-CAPACITY ADENOVIRAL VECTORS



(57) Abstract

The present invention relates to gene therapy, especially to adenovirus-based gene therapy, and related cell lines and compositions. In particular, novel packaging cell lines are disclosed, for use in facilitating the development of high-capacity vectors. The invention also discloses a variety of high-capacity adenovirus vectors and related compositions and kits including the disclosed cell lines and vectors. Finally, the invention discloses methods of preparing and using the disclosed vectors, cell lines and kits.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## PACKAGING CELL LINES FOR USE IN FACILITATING THE DEVELOPMENT OF HIGH-CAPACITY ADENOVIRAL VECTORS

This invention was made with U.S government support under NIH Grant No. HL 54352. The government has certain rights in the invention.

The present invention relates to gene therapy, especially to adenovirus-based gene therapy. In particular, novel packaging cell lines are disclosed, for use in facilitating the development of high-capacity vectors. High-capacity adenovirus vectors are also disclosed herein, as are related compositions, kits, and methods of preparation and use of the disclosed vectors, cell lines and kits.

Enhanced transfer of DNA conjugates into cells has been achieved with adenovirus, a human DNA virus which readily infects epithelial cells (Horwitz, "Adenoviridae and their replication", in Virology, Fields and Knipe, eds., Raven Press, NY (1990) pp. 1679-1740).

Although adenovirus-mediated gene therapy represents an improved method of DNA transfer into cells, a potential limitation of this approach is that adenovirus replication results in disruption of the host cell. In addition, adenovirus also possesses oncogenic properties including the ability of one of its proteins to bind to tumor suppressor gene products. The use of so-called replication defective strains of adenovirus (which typically possess E1A and/or E1B deletions that render the virus unable to replicate in host cells) is in principle more suitable for *in vivo* therapy; however, the potential of co-infection of epithelial cells with wild-type strains of virus resulting in transactivation of the recombinant virus may represent a significant safety concern for *in vivo* applications. Furthermore, it is not yet known which recombinant adenoviruses are capable of integrating their genome into host cell DNA allowing for long-term stable expression of any foreign genes they may be transporting.

Another undesirable aspect of using intact or replication-competent adenovirus as a gene transfer means is that it is an oncogenic virus whose gene products are known to interfere with the function of host cell tumor suppressor proteins as well as immune recognition molecules, such as the major histocompatibility complex (MHC). In addition, pre-existing circulating antibodies to adenovirus may significantly reduce the efficiency of *in vivo* gene delivery. Lastly, only a foreign gene of 6 kilobases (kb) or less can be incorporated into the intact adenovirus genome for gene transfer experiments, whereas DNA segments of greater than 15 kb can be transferred using the methods of this invention.



In order to make Ad vectors more replication-incompetent, some investigators have attempted to construct recombinant Ad-derived vectors which have nearly all of their genome deleted, except for portions known to be required for packaging of virus particles. For example, helper-dependent vectors lacking all viral ORFs but including essential *cis* elements (the inverted terminal repeats -- ITRs -- and the contiguous packaging sequence) have been constructed, but the virions package less efficiently than the helper and package as multimers part of the time, which suggests that the virus may "want" to package a fuller DNA complement (see, e.g., Fisher, et al., Virology 217: 11-22 (1996)). Mitani et al. (PNAS USA 92: 3854-3858 (1995)) also describe a helper-dependent Ad vector that was apparently not completely replication-defective.

Amalfitano, et al. (PNAS USA 93: 3352-3356 (1996)) describe the construction of an Ad packaging cell lines that support the growth of E1- and polymerase-deleted Ad vectors, in an effort to block the replication of Ad vectors *in vivo*. Similarly, Armentano, et al. (Hum. Gene Ther. 6: 1343-53 (1995)) describes Ad vectors with most -- but not all -- of the E4 sequence deleted therefrom. However, since such a small amount of genetic material is deleted from the vectors, their ability to transport therapeutic sequences is rather limited.

In view of the aforementioned problems, the design and construction of the within-disclosed packaging cell lines and systems provides a novel and elegant solution, as described further herein. The use of the recombinant sequences and vectors of this invention to mediate the transfer of foreign genes into recipient cells both *in vitro* and *in vivo* overcomes the limitations of the above-described gene transfer systems. This invention utilizes recombinant constructs which duplicate the cell receptor binding and DNA delivery properties of intact adenovirus virions and thus represents an improved method for gene therapy as well as for antisense-based antiviral therapy.

In contrast to the disadvantages of using intact adenovirus, modified adenovirus vectors requiring a helper plasmid or virus, or so-called replication-deficient adenovirus, the use of recombinant adenovirus-derived vectors according to the present invention provides certain advantages for gene delivery. First, the Ad-derived vectors of the present invention possess all of the functional properties required for gene therapy including binding to epithelial cell receptors and penetration of endocytic vesicles. Therapeutic viral vectors of the present

invention may also be engineered to target the receptors of and achieve penetration of non-epithelial cells; means of engineering viral vectors to accomplish these ends are described in detail hereinbelow.

Second, the vectors of the present invention have deletions of substantial portions of the Ad genome, which not only limits the ability of the Ad-derived vectors to "spread" to other host cells or tissues, but allows significant amounts of "foreign" (or non-native) nucleic acids to be incorporated into the viral genome without interfering with the reproduction and packaging of the viral genome. Therefore, the vectors of the present invention are ideal for use in a wide variety of therapeutic applications.

Third, while the vectors disclosed herein are safe for use as therapeutic agents in the treatment of a variety of human afflictions, they do not require the presence of any "helpers" for propagation and packaging, largely because of the novel cell lines in which they are reproduced. Such cell lines -- referred to herein as packaging cell lines -- comprise yet another aspect of the invention.

To reduce the frequency of contamination with wild-type adenovirus, it is desirable to improve either the viral vector or the cell line to reduce the probability of recombination. For example, an adenovirus from a group with less homology to the group C viruses may be used to engineer recombinant viruses with little propensity for recombination with the Ad5 sequence in 293 cells. Similarly, an epithelial cell line -- 293 or another -- may be prepared according to within-disclosed methods which stably expresses adenovirus proteins or polypeptides from Ad3 and/or proteins or polypeptides from another non-group-C or group C serotype; such a cell line would be useful for supporting adenovirus-derived viral vectors bearing deletions of regulatory and/or-structural genes, irrespective of the serotype from which such a vector was derived.

It is also contemplated that the constructs and methods of the present invention will support the design and engineering of chimeric viral vectors which express amino acid residue sequences derived from two or more Ad serotypes. Thus, unlike methods and constructs available prior to the advent of the present disclosure, this invention allows the greatest possible flexibility in the design and preparation of useful viral vectors and cell lines which support their construction and propagation -- all with a decreased risk of recombining with wild-type Ad to produce potentially-harmful recombinants.

In part, the present invention discloses a simpler, alternative means of reducing the recombination between viral and cellular sequences than those discussed in the art. One such means is to increase the size of the deletion in the recombinant virus and thereby reduce the extent of shared sequences between that virus and any Ad genes present in a packaging cell line e.g., the Ad5 genes in 293 cells, or the various Ad genes in the novel cell lines of the present invention.

Deletions of all or portions of structural genes of the adenovirus have been considered undesirable because of the anticipated deleterious effects such deletions would have on viral reproduction and packaging. Indeed, the use of "helper" viruses or plasmids has often been recommended when using Ad-derived vectors containing large deletions in structural protein sequences precisely for this reason.

Contrary to what has been suggested in the art, however, this invention discloses and claims the preparation, propagation and use of recombinant Ad-derived vectors having deletions of all or part of various gene sequences encoding Ad structural proteins, both as a means of reducing the risk of wild-type adenovirus contamination in virus preparations and as a means of allowing foreign DNA to be packaged in such vectors for a variety of diagnostic and therapeutic applications.

Thus, in one embodiment of the present invention, a packaging cell line wherein DNA sequences encoding one or more adenovirus regulatory polypeptides and DNA sequences encoding one or more adenovirus structural polypeptides have been stably integrated into the cellular genome is disclosed.

Thus, in a further embodiment of the present invention, a packaging cell line expressing one or more adenovirus structural proteins, polypeptides, or fragments thereof, wherein said structural protein is selected from the group consisting of:

- a. penton base;
- b. hexon;
- c. fiber;
- d. polypeptide IIIa;
- e. polypeptide V;
- f. polypeptide VI;

- g. polypeptide VII;
- h. polypeptide VIII; and
- i. biologically active fragments thereof is disclosed.

In one variation, the sequences are constitutively expressed; in another, one or more sequences is under the control of a regulatable promoter. In a preferred embodiment expression is constitutive. In various preferred embodiments, the polypeptides expressed by the DNA sequences are biologically active.

In a further and preferred embodiment the packaging cell line of the present invention supports the production of a viral vector. In a preferred embodiment the viral vector is a therapeutic vector.

In one aspect of the present invention, each DNA sequence is introduced into the genome of the within-disclosed cell lines via a separate complementing plasmid. In other embodiments, two or more DNA sequences were introduced into the genome via a single complementing plasmid. In one variation, the complementing plasmid comprises a DNA sequence encoding adenovirus fiber protein, polypeptide or fragment thereof. An example of a useful complementing plasmid according to the present invention is a plasmid having the characteristics of pCLF (for deposit details, see Example 3)

In another aspect of the present invention, the complementing plasmid used to transform a cell line of the present invention further comprises a DNA sequence encoding an adenovirus regulatory protein, polypeptide or fragment thereof. In one variation, the regulatory protein is selected from the group consisting of E1A, E1B, E2A, E2B, E3, E4 and L4 (also referred to as "the 100K protein"); an exemplary complementing plasmid has the characteristics of is pE4/Hygro?? (for deposit details, see Example 3). In another aspect, the complementing plasmid used to transform a cell line of the present invention further comprises a DNA sequence encoding two or more of the above mentioned adenovirus regulatory proteins, polypeptides or fragments thereof.

In one variation, the two or more regulatory proteins, polypeptides or fragments thereof are selected from the group consisting of E1A, E1B, E2A, E2B, E3, E4 and L4 (also referred to as "the 100K protein"). In another variation, the structural protein is selected from the group consisting of penton base; hexon; fiber; polypeptide IIIa; polypeptide V; polypeptide VI; polypeptide VII; polypeptide VIII; and biologically active fragments thereof.

In one variation of the present invention, a packaging cell line expresses fiber protein. In one embodiment, the fiber protein has been modified to include a non-native amino acid residue sequence which targets a specific receptor, but which does not disrupt trimer formation or transport of fiber into the nucleus. In another variation, the non-native amino acid residue sequence alters the binding specificity of the fiber for a targeted cell type. In still another embodiment, the structural protein is fiber comprising amino acid residue sequences from more than one adenovirus serotype. As disclosed herein, the nucleotide sequences encoding fiber protein or polypeptide need not be modified solely at one or both termini; fiber protein -- and indeed, any of the adenovirus structural proteins, as taught herein -- may be modified "internally" as well as at the termini.

The present invention also discloses a packaging cell line wherein the viral vector produced in that cell line comprises a nucleic acid sequence having a deletion or mutation of a DNA sequence encoding an adenovirus structural protein, polypeptide, or fragment thereof. In one variation, the viral vector further comprises a nucleic acid sequence having a deletion or mutation of the DNA sequences encoding regulatory polypeptides E1A and E1B. In another variation, the viral vector further comprises a nucleic acid sequence having a deletion or mutation of a DNA sequence encoding one or more of the following regulatory proteins or polypeptides: E2A, E2B, E3, E4, L4, or fragments thereof.

Yet another variation discloses that a foreign DNA sequence encoding one or more foreign proteins, polypeptides or fragments thereof has been inserted in place of any of the deletions in the therapeutic viral vector. In one embodiment, the foreign DNA encodes a tumor-suppressor protein or a biologically active fragment thereof. In another embodiment, the foreign DNA encodes a suicide protein or a biologically active fragment thereof. As before, cell lines as described herein may be procaryotic or eucaryotic in origin, with mammalian cell lines often being preferred. Epithelial and non-epithelial cell lines are useful in the aforementioned variations; some particularly useful cell lines include 293, A549, W162, HeLa, Vero, 211, and 211A cell lines.

The invention further contemplates that the aforementioned cell lines support the production of viral vectors including foreign DNA sequences encoding one or more foreign proteins, polypeptides or fragments thereof has been inserted in place of any structural and/or

regulatory proteins (or portions thereof) that have been deleted. Thus, in one embodiment, the foreign DNA encodes a tumor-suppressor protein; a suicide protein; a cystic fibrosis transmembrane conductance regulator (CFTR) protein; or a biologically active fragment of any of them.

Any of the within-disclosed cell lines may have a DNA sequence encoding all or part of a fiber protein -- including modified or chimeric proteins -- stably integrated into the genome. thus, in one variation, the fiber protein has been modified to include a non-native amino acid residue sequence which targets a specific receptor, but which does not disrupt trimer formation or transport of fiber into the nucleus. In one variation, the non-native amino acid residue sequence is coupled to the carboxyl terminus of the fiber. In yet another, the non-native amino acid residue sequence further includes a linker sequence. Alternatively, the fiber protein further comprises a ligand coupled to the linker. A suitable ligand may be selected from the group consisting of ligands that specifically bind to a cell surface receptor and ligands that can be used to couple other proteins or nucleic acid molecules. In one variation, the ligand is selected from the group consisting of ligands that specifically bind to a cell surface receptor and ligands that can be used to couple other proteins or nucleic acid molecules.

In yet another embodiment, the non-native amino acid residue sequence is incorporated into the fiber amino acid residue sequence at a location other than one of the fiber termini. Alternatively, the non-native amino acid residue sequence alters the binding specificity of the fiber for a targeted cell type. In other embodiments, the linker sequence alters the binding specificity of the fiber for a targeted cell type. The expressed fiber may, in various embodiments, bind to a specific targeted cell type not usually targeted by adenovirus and/or may comprise amino acid residue sequences from more than one adenovirus serotype.

In various aspects of the present invention, a packaging cell line of the present invention is derived from a procaryotic cell line; in another, it is derived from a eucaryotic cell line. While various embodiments suggest the use of mammalian cells, and more particularly, epithelial cell lines, a variety of other, non-epithelial cell lines are used in various embodiments. Thus, while various embodiments disclose the use of a cell line selected from the group consisting of 293, A549, W162, HeLa, Vero, 211, and 211A cell lines, it is understood that various other cell lines are likewise contemplated for use as disclosed herein.

The invention further discloses a wide variety of nucleic acid sequences and viral vectors. Thus, in one embodiment, the invention discloses a nucleic acid sequence encoding any one of the aforementioned adenovirus fiber proteins, polypeptides or fragments thereof -- including, without limitation, those that include deletions or other mutations; those that are chimeric; and those that have linkers, foreign amino acid residues, or other molecules attached for various purposes as disclosed herein. Nucleic acid sequences encoding various other adenovirus structural and/or regulatory proteins or polypeptides are also within the scope of the present invention.

A wide variety of therapeutic viral vectors are also embodiments of the present invention. In one embodiment, a therapeutic viral vector is disclosed which lacks a DNA sequence encoding fiber protein, or a portion thereof. In another variation, a therapeutic viral vector may further or alternatively comprise deletion of a DNA sequence encoding one or more regulatory proteins, polypeptides, or fragments thereof. In various embodiments, foreign DNA sequences are inserted in place of the DNA sequence encoding fiber protein in the viral vectors of the present invention. In other embodiments, the therapeutic viral vectors further comprise foreign DNA sequences inserted in place of the DNA sequences encoding one or more regulatory proteins, polypeptides, or fragments thereof, and/or one or more structural proteins, polypeptides, or fragments thereof.

The present invention further discloses a number of viral vectors. In one variation, a viral vector comprises a deletion or mutation of a DNA sequence encoding an adenovirus structural protein, polypeptide, or fragment thereof. A vector may further comprise deletion or mutation of the DNA sequences encoding regulatory polypeptides E1A and E1B; and it may still further comprise deletion or mutation of the DNA sequence encoding one or more of the following regulatory proteins or polypeptides: E2A, E2B, E3, E4, L4, or fragments thereof. In another variation, in a viral vector of the present invention, the structural protein comprises fiber. Any combination of the foregoing is also contemplated by the present invention. The viral vectors of the present invention are suitable for the preparation of pharmaceutical compositions comprising any of the therapeutic viral vectors disclosed herein -- including combinations thereof -- are also disclosed herein. A further use of the viral vectors of the present invention is for targeting specific cells in a cell population comprising different cell types.

The invention further discloses complementing plasmids and methods of making same. In one embodiment, a complementing plasmid comprises a promoter nucleotide sequence operatively linked to a nucleotide sequence encoding an adenovirus structural polypeptide. In one variation, the complementing plasmid comprises pCLF. In another variation, a complementing plasmid further comprises a nucleotide sequence encoding a first adenovirus regulatory polypeptide, a nucleotide sequence encoding a second regulatory polypeptide, a nucleotide sequence encoding a third regulatory polypeptide; or any combination of the foregoing. In still another embodiment, the adenovirus structural polypeptide is selected from the group consisting of penton base; hexon; fiber; polypeptide IIIa; polypeptide V; polypeptide VI; polypeptide VII; polypeptide VIII; and biologically active fragments thereof.

The present invention also discloses a complementing plasmid comprising a promoter nucleotide sequence operatively linked to a nucleotide sequence encoding an adenovirus structural protein, polypeptide or fragment thereof and a nucleotide sequence encoding an adenovirus regulatory protein, polypeptide or fragment thereof. In one variation, the early region polypeptide is E4; in another, the plasmid comprises pE4/Hygro. In still another variation, the early region polypeptides are E1 and E4. Complementing plasmids further comprising a nucleotide sequence encoding an adenovirus structural protein, polypeptide or fragment thereof are also contemplated, as are plasmids wherein the promoter nucleotide sequence is selected from the group consisting of MMTV, CMV and E4 promoter nucleotide sequences.

Viral vectors are also disclosed which comprise nucleotide sequences encoding a packaging signal and a foreign protein or polypeptide, wherein the nucleotide sequence encoding an adenovirus structural protein has been deleted. In one variation, the nucleotide sequence encoding the foreign protein or polypeptide is a DNA molecule up to about 3 kb in length; in another, the nucleotide sequence encoding the foreign protein or polypeptide is a DNA molecule up to about 9.5 kb in length; in still another, the nucleotide sequence encoding the foreign protein or polypeptide is a DNA molecule up to about 12.5 kb in length. Nucleotide sequences of intermediate lengths are also contemplated by the present invention, as are sequences in excess of 12.5 kb.

The invention also discloses viral vectors wherein the sequence encoding a foreign protein or polypeptide is a sequence encoding an anti-tumor agent, a tumor suppressor protein,



a suicide protein, or a fragment or functional equivalent thereof. In one variation, nucleotide sequences encoding one or more regulatory proteins have also been deleted from the vector. In another, the regulatory proteins are selected from the group consisting of E1A, E1B, E2A, E2B, E3, E4, and L4 (100K protein).

In various embodiments, the adenovirus is a Group C adenovirus selected from serotypes 1, 2, 5, or 6; in other embodiments, adenovirus selected from other serotypes are useful as disclosed herein. The invention also discloses useful vaccines comprising a viral vector according to any of the foregoing specifications, and a pharmaceutically acceptable carrier or excipient.

Various useful compositions are also disclosed herein. One embodiment discloses a composition useful in the preparation of recombinant adenovirus viral vectors comprising a cell containing a delivery plasmid comprising an adenovirus genome lacking a nucleotide sequence encoding fiber. In one variation, the cell further comprises a complementing plasmid containing a nucleotide sequence encoding fiber, the plasmid being stably integrated into the cellular genome of the cell. In another variation, the delivery plasmid further comprises a nucleotide sequence encoding a foreign polypeptide. In one variation, the delivery plasmid is pDV44, p E1B gal, or p E1sp1B.

In another embodiment, the polypeptide is a therapeutic molecule. In yet another, the polypeptide is a therapeutic molecule. Another variation provides that the delivery plasmid further comprises a nucleotide sequence encoding a foreign polypeptide.

Compositions useful in the preparation of recombinant adenovirus viral vectors are also disclosed. In one embodiment, a composition comprises a cell containing a first delivery plasmid comprising an adenovirus genome lacking a nucleotide sequence encoding fiber and incapable of directing the packaging of new viral particles in the absence of a second delivery plasmid; and a second delivery plasmid comprising an adenoviral genome capable of directing the packaging of new viral particles in the presence of the first delivery plasmid.

In another variation, the first and second delivery plasmids interact within the cell to produce a therapeutic viral vector. In yet another, the cell further comprises a complementing plasmid containing a nucleotide sequence encoding fiber, the plasmid being stably integrated into the cellular genome of the cell. In still another, the first or second delivery plasmid

further comprises a nucleotide sequence encoding a foreign polypeptide. In various embodiments, the polypeptide is a therapeutic molecule.

Another embodiment discloses a composition as before, wherein the first delivery plasmid lacks adenovirus packaging signal sequences. In another aspect, the second delivery plasmid contains a LacZ reporter construct. Another variation discloses that the second delivery plasmid further lacks a nucleotide sequence encoding an adenovirus regulatory protein. In one variation, the regulatory protein is E1. In one embodiment of the above-noted compositions, the complementing plasmid has the characteristics of pCLF.

In another embodiment, a composition is disclosed wherein the first delivery plasmid lacks a nucleotide sequence encoding an adenovirus structural protein and the second delivery plasmid lacks a nucleotide sequence encoding adenovirus E1 protein. In another, the first delivery plasmid lacks a nucleotide sequence encoding adenovirus E4 protein and the second delivery plasmid lacks a nucleotide sequence encoding adenovirus E1 protein. In still another, the cell contains at least one complementing plasmid encoding an adenoviral regulatory protein and a structural protein.

In alternative embodiments, the regulatory protein is E4 and the structural protein is fiber; or the regulatory protein is E1 and the structural protein is fiber. In still another embodiment, the adenoviral regulatory protein and the structural protein are encoded by separate complementing plasmids.

Another variation discloses a composition wherein the cell is selected from the group consisting of 293, A549, W162, HeLa, Vero, 211, and 211A. In another embodiment, the delivery plasmid is DV1, or p E1B gal, or p E1sp1B.

Various methods of making and using the vectors, plasmids, cell lines and other compositions and constructs of the present invention are also disclosed herein. The following methods are considered exemplary and not limiting.

Thus, in one variation, the invention discloses a method of constructing therapeutic viral vectors, comprising introducing a delivery plasmid into an Ad fiber-expressing complementing cell line, wherein the DNA sequence encoding Ad fiber protein has been deleted from the delivery plasmid. In one variation, the delivery plasmid further includes a DNA sequence encoding a foreign protein, polypeptide, or fragment thereof. In other embodiments, the delivery plasmid is DV1, p E1B gal, p E1sp1B, or similar constructs.

The invention also discloses methods of transforming a pathologic hyperproliferative mammalian cell comprising contacting the cell with any of the vectors described herein. In another embodiment, methods of infecting a mammalian target cell with a viral vector containing a preselected foreign nucleotide sequence are disclosed. One such variation comprises the following steps: (a) infecting the target cell with a viral vector of the present invention, the viral vector carrying a preselected foreign nucleotide sequence; and (b) expressing the foreign nucleotide sequence in the targeted cell.

The invention also encompasses mammalian target cells infected with a preselected foreign nucleotide sequence produced by the methods disclosed herein. In one variation, the target cells are selected from the group consisting of replicating, slow-replicating and non-replicating human cells.

Methods of treating an acquired or hereditary disease are also disclosed. One method comprises (a) administering a pharmaceutically acceptable dose of a viral vector to a target cell, wherein the vector comprises a preselected therapeutic nucleotide sequence; and (b) expressing the therapeutic sequence in the target cell for a time period sufficient to ameliorate the acquired or hereditary disease in the cell. Method of gene therapy comprising administering to a subject an effective amount of a therapeutic viral vector produced by a packaging cell line of the present invention are also disclosed.

Also contemplated by the present invention are various methods of inhibiting the proliferation of a tumor in a subject comprising administering an effective amount of a therapeutic viral vector of the present invention under suitable conditions to the subject. IN one variation, the gene encodes an anti-tumor agent. In another variation, the agent is a tumor-suppressor gene. In still another embodiment, the agent is a suicide gene or a functional equivalent thereof. In another variation, the vector is administered via intra-tumoral injection.

The invention also discloses systems or kits for use in any of the aforementioned methods. The systems or kits may contain any appropriate combination of the within-described vectors, plasmids, cell lines, and additional therapeutic agents as disclosed. Preferably, each such kit or system includes a quantity of the appropriate therapeutic substance or sequence sufficient for at least one administration, and instructions for administration and use. Thus, one system further comprises an effective amount of a therapeutic agent which enhances the therapeutic effect of the therapeutic viral vector-containing composition.

Another variation discloses that the composition and the therapeutic agent are each included in a separate receptacle or container.

It will also be appreciated that any combination of the preceding elements may also be efficacious as described herein, and that all related methods are also within the scope of the present invention.

### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a schematic diagram of the entire adenoviral E4 transcriptional unit with the open reading frames (ORF) indicated by blocked segments along with the promoter and terminator sequences. The location of primers for amplifying specific portions of E4 are also indicated as further described in Example 1A.

Figure 2 is a schematic map of plasmid pE4/Hygro as further described in Example 1B.

Figure 3 is a schematic map of plasmid pCDNA3/Fiber as further described in Example 1B.

Figure 4 is a schematic map of plasmid pCLF as further described in Example 1B.

Figure 5 is a photograph of a Southern blot showing the presence of intact adenovirus E4 3.1 kilobase (kb) insert in the 211 cell line as further described in Example 1C.

Figure 6 is a photograph of a Western blot showing labeled fiber protein detected under native and denaturing electrophoresis conditions as described in Example 1C. The 293 cells lack fiber while the sublines 211A, 211B and 211R contain fiber protein detectable in functional trimerized form and denatured monomeric form.

Figure 7 is a schematic map of plasmid pDEX/E1 as further described in Example 1D.

Figure 8 is a schematic map of plasmid pE1/Fiber as further described in Example 1F1.

Figure 9 is a schematic map of plasmid pE4/Fiber as further described in Example 1F2).

Figure 10 is a schematic illustration of linearized ~~p3~~E1B gal delivery plasmids for use in cotransfection and recombination to form a recombinant adenoviral vector having multiple adenoviral gene deletions. The plasmids and recombination event are more fully described in Example 2A.

Figure 11 is a schematic of plasmid p11.3 as further described in Example 2A used in the construction of pDV44 delivery plasmid with plasmid p8.2.

Figure 12 is a schematic of plasmid p8.2 as further described in Example 2A used in the construction of pDV44 delivery plasmid with plasmid p11.3.

Figure 13. **Trimeric structure of the recombinant fiber:** 293, 211A, 211B, or 211R cells as indicated were metabolically labeled with [<sup>35</sup>S]methionine, soluble protein extracts prepared, and fiber was immunoprecipitated. A portion of the precipitated protein was electrophoresed on an 8% SDS-PAGE gel under either semi-native or denaturing conditions. The positions of trimeric (T) and monomeric (M) fiber are indicated. As a control for electrophoretic conditions, recombinant Ad2 fiber produced in baculovirus-infected cells was run under identical conditions and stained with Coomassie blue.

**Fig. 14. Complementation of a fiber mutant adenovirus by fiber-producing cells:** The cell lines indicated ( $2 \times 10^6$  cells per sample) were infected with the temperature-sensitive fiber mutant adenovirus H5ts142 at 10 PFU/cell and incubated at either the permissive (32.5 °C, stippled bars) or the restrictive (39.5 °C, solid bars) temperature. 48 hours post-infection, virus was isolated by freeze-thaw lysis and yields determined by fluorescent focus assay on SW480 cells. Each value represents the mean of duplicate samples, and the data shown is representative of multiple experiments.

**Fig. 15. Incorporation of the recombinant Ad5 fiber into Ad3 particles:** A) Alignment of the N-terminal (penton base-binding) domains of fiber proteins from several different adenovirus serotypes. B) Type 3 adenovirus was propagated in 293, 211B, or 211R cells as indicated and purified by two sequential CsCl centrifugations. 10 mg of the purified viral particles was then electrophoresed under denaturing conditions and transferred to a PVDF membrane. Ad5 fiber was detected with a polyclonal rabbit antibody raised against recombinant Ad2 fiber. As a positive control for detection, 400 ng of wild-type Ad2 was run in the lane marked 'Ad2'. Under these conditions, the mobilities of the Ad2 and Ad5 fibers are indistinguishable and the antibody reacts with both proteins.

**Fig. 16. Nuclear localization of the recombinant fiber protein in three packaging cell lines:** Cells were grown on 8-well chamber slides, stained with a rabbit anti-fiber polyclonal antibody and visualized

with a FITC-conjugated goat anti-rabbit antibody. A) line 211A. B) Line 211B C) Line 211R D) 293 cells (negative control). E) 293 cells infected with Ad.RSVbgal at 1 pfu/cell and stained 24 hour post-infection (positive control). F) Infected cells prepared as in (E) but stained without the primary antibody.

DETAILED DESCRIPTION To reduce the frequency of contamination with wild-type adenovirus, it is considered desirable to improve either the viral vector or the cell line to reduce the probability of recombination. For example, an adenovirus from a group with less homology to the group C viruses may be used to engineer recombinant viruses with little propensity for recombination with the Ad5 sequence in 293 cells. Similarly, an epithelial cell line -- e.g. the cell line known as 293 -- may be used or further modified according to within-disclosed methods which stably expresses adenovirus proteins or polypeptides from Ad3 and/or proteins or polypeptides from another non-group-C or group C serotype; such a cell line would be useful to support adenovirus-derived viral vectors bearing deletions of regulatory and/or structural genes, irrespective of the serotype from which such a vector was derived.

It is also contemplated that the constructs and methods of the present invention will support the design and engineering of chimeric viral vectors which express amino acid residue sequences derived from two or more Ad serotypes. Thus, unlike methods and constructs available prior to the advent of the present disclosure, this invention allows the greatest possible flexibility in the design and preparation of useful viral vectors and cell lines which support their construction and propagation -- all with a decreased risk of recombining with wild-type Ad to produce potentially-harmful recombinants.

In part, the present invention discloses a simpler, alternative means of reducing the recombination between viral and cellular sequences than those discussed in the art. One such means is to increase the size of the deletion in the recombinant virus and thereby reduce the

extent of shared sequences between that virus and any Ad genes present in a packaging cell line

— e.g., the Ad5 genes in 293 cells, or the various Ad genes in the novel cell lines of the present invention.

By the term "substantially homologous" is meant having at least 80%, preferably at least 90%, most preferably at least 95% homology therewith.

The amino acid residues described herein are preferably in the "L" isomeric form. However, residues in the "D" isomeric form can be substituted for any L-amino acid residue, as long as the desired functional property is retained by the polypeptide.  $\text{NH}_2$  refers to the free amino group present at the amino terminus of a polypeptide.

DNA Homolog: A nucleic acid having a preselected conserved nucleotide sequence and a sequence encoding a preferred polypeptide according to the present invention.

Foreign Gene: This term is used to identify a DNA molecule not present in the exact orientation and position as the counterpart DNA molecule found in wild-type adenovirus. It may also refer to a DNA molecule from another Ad serotype or from an entirely different species -- e.g. a human DNA sequence.

Penton: The terms "penton" or "penton complex" are preferentially used herein to designate a complex of penton base and fiber. The term "penton" may also be used to indicate penton base, as well as penton complex. The meaning of the term "penton" alone should be clear from the context within which it is used.

Polypeptide and Peptide: These terms are used interchangeably herein to designate a series of no more than about 50 amino acid residues connected one to the other by peptide bonds between the alpha-amino and carboxy groups of adjacent residues.

Receptor: Receptor is a term used herein to indicate a biologically active molecule that specifically binds to (or with) other molecules. The term "receptor protein" may be used to more specifically indicate the proteinaceous nature of a specific receptor.

Transgene or Therapeutic Nucleotide Sequence: As described and claimed herein, such a sequence includes DNA and RNA sequences encoding an RNA or polypeptide. Such sequences may be "native" or naturally-derived sequences; they may also be "non-native" or "foreign" sequences which are naturally- or recombinantly-derived. The term "transgene," which may be used interchangeably herein with the term "therapeutic nucleotide sequence," is often used to describe a heterologous or foreign (exogenous) gene that is carried by a viral vector and transduced into a host cell.

Therefore, therapeutic nucleotide sequences include antisense sequences or nucleotide sequences which may be transcribed into antisense sequences. Therapeutic nucleotide sequences (or transgenes) further comprise sequences which function to produce a desired effect in the cell or cell nucleus into which said therapeutic sequences are delivered. For example, a therapeutic nucleotide sequence may encode a functional protein intended for delivery into a cell which is unable to produce that functional protein.

Expression or Delivery Vector: Any plasmid or virus into which a foreign DNA may be inserted for expression in a suitable host cell -- i.e., the protein or polypeptide encoded by the DNA is synthesized in the host cell's system. Vectors capable of directing the expression of DNA segments (genes) encoding one or more proteins are referred to herein as "expression vectors". Also included are vectors which allow cloning of cDNA (complementary DNA) from mRNAs produced using reverse transcriptase.

Adenoviral Vector or Ad-Derived Vector: Any adenovirus-derived plasmid or virus into which a foreign DNA may be inserted or expressed. This term may also be used interchangeably with "viral vector" This "type" of vector may be utilized to carry nucleotide



sequences encoding therapeutic proteins or polypeptides to specific cells or cell types in a subject in need of treatment, as described further hereinbelow.

Complementing Plasmid: This term is generally used herein to describe plasmid vectors used to deliver particular nucleotide sequences into a packaging cell line, with the intent of having said sequences stably integrate into the cellular genome.

Delivery Plasmid: This term is generally used herein to describe a plasmid vector that carries or delivers nucleotide sequences in or into a cell line (e.g., a packaging cell line) for the purpose of propagating therapeutic viral vectors of the present invention.

The adenovirus (Ad) particle is relatively complex and may be resolved into various substructures. The outer shell is strikingly icosahedral in shape and, at first glance, appears to have a triangulation number of 25. The structures at the fivefold positions ("pentons") are different from the rest ("hexons"), however, and the hexons are chemically trimers rather than hexamers. Thus, the structure really does not correspond to a simple sub-triangulated icosahedral design. (See, e.g., Fields, et al., Virology, Vol. I, Raven Press, NY, pp. 54-56 (1990).)

Fiber plays a crucial role in adenovirus infection by attaching the virus to a specific receptor on the cell surface. The fiber consists of three domains: an N-terminal tail that interacts with penton base; a shaft composed of 22 repeats of a 15-amino-acid segment that forms  $\alpha$ -sheet and  $\alpha$ -bends; and a knob at the C-terminus that contains the type-specific antigen and is responsible for binding to the cell surface receptor. The fiber protein is also responsible for transport of viral nucleic acids into the nucleus. The gene encoding the fiber protein from Ad2 has been expressed in human cells and has been shown to be correctly assembled into trimers, glycosylated and transported to the nucleus. (See, e.g., Hong and Engler, Virology 185: 758-761 (1991).) Thus, alteration of gene delivery mediated by recombinant adenovirus vectors to specific cell types has great utility for a variety of gene therapy applications and is thus one of the objects of the present invention.

Hexon and penton capsomeres are the major components on the surface of the virion. Their constituent polypeptides, nos. II, III and IV, contain tyrosine residues that are exposed on the surface of the virion and can be labeled -- e.g., by iodination of intact particles.

The fiber is an elongated protein which exists as a trimer of three identical polypeptides (polypeptide IV) of 582 amino acids in length. The N-terminus of the fiber mediates binding to the penton base to form what is generally called the penton capsomere. The C-terminus of the fiber is involved in initial binding of the virus to cellular receptors.

The 35,000+ base pair (bp) genome of adenovirus type 2 has been sequenced and the predicted amino acid sequences of the major coat proteins (hexon, fiber and penton base) have been described. (See, e.g., Neumann et al., Gene 69: 153-157 (1988); Herisse et al., Nuc. Acids Res. 9: 4023-4041 (1981); Roberts et al., J. Biol. Chem. 259: 13968-13975 (1984); Kinloch et al., J. Biol. Chem. 259: 6431-6436 (1984); and Chroboczek et al., Virology 161: 549-554 (1987).)

The sequence of Ad5 DNA was completed more recently; its sequence includes a total of 35,935 bp. Portions of many other adenovirus genomes have also been sequenced. It is presently understood that the upper packaging limit for adenovirus virions is about 105% of the wild-type genome length. (See, e.g., Bett, et al., J. Virol. 67(10): 5911-21 (1993).) Thus, for Ad2 and Ad5, this would be an upper packaging limit of about 38kb of DNA.

Adenovirus DNA also includes inverted terminal repeat sequences (ITRs) ranging in size from about 100 to 150 bp, depending on the serotype. The inverted repeats enable single strands of viral DNA to circularize by base-pairing of their terminal sequences, and the resulting base-paired "panhandle" structures are thought to be important for replication of the viral DNA.

For efficient packaging, the ITRs and the packaging signal (a few hundred bp in length) appear to comprise the "minimum requirement." Helper-dependent vectors lacking all viral ORFs but including these essential *cis* elements (the ITRs and contiguous packaging sequence) have been constructed, but the virions package less efficiently than the helper and package as multimers part of the time, which suggests that the virus may "want" to package a fuller DNA complement (see, e.g., Fisher, et al., Virology 217: 11-22 (1996)).

While some prefer to use replication-defective Ad viral vectors for fear that replication-competent vectors raise safety issues, the viral vectors of the present invention may retain their ability to express the genome packaged within -- i.e., they may retain their "infectivity" -- they do not act as infectious agents to the extent that they cause disease in the subjects to whom they are administered for therapeutic purposes.

It is to be appreciated that the viral vectors of the present invention have several distinct advantages over adenoviral and Ad-derived vectors described in the art. For example, recombination of such vectors is rare; there are no known associations of human malignancies with adenoviral infections despite common human infection with adenoviruses; the genome may be manipulated to accommodate foreign genes of a fairly substantial size; and host proliferation is not required for expression of adenoviral proteins.

An extension of this invention is that the Ad-derived viral vectors disclosed herein may be used to target and deliver genes into specific cells by incorporating the attachment sequence for other receptors (such as CD4) onto the fiber protein by recombinant DNA techniques, thus producing a chimeric molecule. This should result in the ability to target and deliver genes into a wide range of cell types with the advantage of evading recognition by the host's immune system. The within-disclosed delivery systems thus provide for increased flexibility in gene design to enable stable integration into proliferating and nonproliferating cell types.

For example, published International App. No. WO95/26412 and Krasnykh, et al. (J. Virol. 70: 6839-46 (1996)), the disclosures of which are incorporated by reference herein, describe modifications that may be made to the adenovirus fiber protein. Such modifications are useful in altering the targeting mechanism and specificity of adenovirus and could readily be utilized in conjunction with the constructs of the present invention to target the novel viral vectors disclosed herein to different receptors and different cells. Moreover, modifications to fiber protein which alter its tropism may permit greater control over the localization of viral vectors in therapeutic applications.

Similarly, incorporation of various structural proteins into cell lines of the present invention, whether or not those proteins are modified, is also contemplated by the present invention. Thus, for example, modified penton base polypeptides such as those described in Wickham, et al. (J. Virol. 70: 6831-8 (1996)) may have therapeutic utility when used according to the within-disclosed methods.

### C. Packaging Cell Lines

The first generation of recombinant adenoviral vectors currently available tend to have a deletion in the first viral early gene region which is generally referred to as E1, which comprises the E1a and E1b regions. (These regions typically span genetic map units 1.30 to 9.24.) Figure 3 in chapter 67 of Fields Virology, 3d Ed. (Fields et al. (eds.), Lippincott-Raven Publ., Philadelphia, (1996), p. 2116) illustrates a transcription and translation map of adenovirus type 2 (Ad2) that is a helpful example.

According to various published reports, deletion of the viral E1 region renders the recombinant adenovirus defective for replication and incapable of producing infectious viral particles in the subsequently-infected target cells. Thus, the ability to generate E1-deleted adenovirus is often based on the availability of the human embryonic kidney packaging cell line called 293. This cell line contains the E1 region of adenovirus, which provides E1 gene region products to "support" the growth of E1-deleted virus in the cell line (see, e.g., Graham et al., J. Gen. Virol. 36: 59-71 (1977)).

Nevertheless, the inherent problems with current first-generation recombinant adenoviruses have raised increasing concerns about their use in patients. For example, several recent studies have shown that E1-deleted adenoviruses are not completely replication-incompetent (see Rich, Hum. Gene. Ther. 4: 461-476 (1993); Engelhardt, et al., Nature Genet. 4: 27-34 (1993)).

Three general limitations are associated with the adenoviral vector technology. First, infection both *in vivo* and *in vitro* with the adenoviral vector at high multiplicity of infection ("MOI") has resulted in cytotoxicity to the target cells, due to the accumulation of penton protein, which is itself toxic to mammalian cells (Kay, Cell Biochem. 17E: 207 (1993)). Second, host immune responses against adenoviral late gene products, including penton protein, cause the inflammatory response and destruction of the infected tissue which received the vectors (Yang, et al., PNAS USA 92: 4407-4411 (1994)). Lastly, host immune responses and cytotoxic effects together prevent the long-term expression of transgenes and cause decreased levels of gene expression following subsequent administration of adenoviral vectors (Mittal, et al., Virus Res. 28: 67-90 (1993)).

The packaging cell lines disclosed herein support viral vectors with deletions of major portions of the viral genome, without the need for helper viruses.

D. Therapeutic Viral Vectors and Related Systems

1. Nucleic Acid Segments

A therapeutic viral vector or composition of the present invention comprises a nucleotide sequence encoding a protein or polypeptide molecule -- or a biologically active fragment thereof -- which may be used for therapeutic applications, as described herein. A therapeutic viral vector or composition may further comprise an enhancer element or a promoter located 5' to and controlling the expression of such a therapeutic nucleotide sequence or gene.

In general, promoters are DNA segments that contain a DNA sequence that controls the expression of a gene located 3' or downstream of the promoter. The promoter is the DNA sequence to which RNA polymerase specifically binds and initiates RNA synthesis (transcription) of that gene, typically located 3' of the promoter. If more than one nucleic acid sequence encoding a particular polypeptide or protein is included in a therapeutic viral vector or nucleotide sequence, more than one promoter or enhancer element may be included, particularly if that would enhance efficiency of expression. For purposes of the present invention, regulatable (inducible) as well as constitutive promoters may be used, either on separate vectors or on the same vector.

A subject therapeutic nucleotide composition or vector consists of a nucleic acid molecule that comprises at least 2 different operatively linked DNA segments. The DNA can be manipulated and amplified by PCR as described herein and by using standard techniques, such as those described in Molecular Cloning: A Laboratory Manual, 2nd Ed., Sambrook et al., eds., Cold Spring Harbor, New York (1989). Typically, to produce a therapeutic viral vector of the present invention, the sequence encoding the selected therapeutic composition and the promoter or enhancer are operatively linked to a DNA molecule capable of autonomous replication in a cell either *in vivo* or *in vitro*. By operatively linking the enhancer element or promoter and the nucleotide sequence encoding the therapeutic nucleotide composition to the vector, the attached segments are replicated along with the vector sequences.

Thus, a recombinant DNA molecule (rDNA) of the present invention is a hybrid DNA molecule comprising at least 2 nucleotide sequences not normally found together in nature. In various preferred embodiments, one of the sequences is a sequence encoding an Ad-derived polypeptide, protein, or fragment thereof. Stated another way, a therapeutic nucleotide sequence of the present invention is one that encodes an expressible protein, polypeptide or fragment thereof, and it may further include an active constitutive or regulatable (e.g. inducible) promoter sequence.

A therapeutic viral vector or composition of the present invention is optimally from about 20 base pairs to about 40,000 base pairs in length. Preferably the nucleic acid molecule is from about 50 bp to about 38,000 bp in length. In various embodiments, the nucleic acid molecule is of sufficient length to encode one or more adenovirus proteins or functional polypeptide portions thereof. Since individual Ad polypeptides vary in length from about 19 amino acid residues to about 967 amino acid residues, corresponding nucleotide sequences will range from about 50 bp up to about 3000 bp, depending on the size and of individual polypeptide-encoding sequences that are "replaced" in the viral vectors by therapeutic nucleotide sequences of the present invention.

Various Ad proteins are comprised of more than one polypeptide sequence. Thus, deletion of the corresponding genes from an Ad vector as taught herein will thus allow the vector to accommodate even larger "foreign" DNA segments. Thus, if the sequences encoding one or more adenovirus polypeptides or proteins are supplanted by a recombinant nucleotide sequence of the present invention, the length of the recombinant sequence can conceivably extend nearly to the packaging limit of the relevant adenovirus-derived vector.

In view of the fact that preferred embodiments disclosed herein are helper-independent Ad-derived vectors, the entire wild-type Ad genome cannot be completely supplanted by recombinant nucleic acid molecules without transforming such a vector into a vector requiring "help" of some kind. However, the Ad-derived vectors of the present invention do not depend on a helper virus; instead, the vectors of the present invention are propagated in cell lines stably expressing proteins or polypeptides that have been removed from said vectors to allow the addition of "foreign" DNA into the vectors. In various disclosed embodiments, specific early region and structural polypeptides are deleted from the vectors of the present invention, thereby enabling the vectors to accommodate recombinant nucleic acid sequences (or

cassettes) of various lengths. For example, Ad-derived vectors of the present invention may easily include 12 kb or more of foreign (or "therapeutic") DNA sequences.

The therapeutic (or foreign) nucleotide sequence can be a gene or gene fragment that encodes a protein or polypeptide -- or a biologically active fragment thereof -- that provides a desired therapeutic effect such as replacement of alpha 1-antitrypsin or cystic fibrosis transmembrane conductance regulator protein (CFTR) and the like. (See, e.g., Crystal, et al., Nature Genetics 8: 42-51 (1994); Zabner, et al., Cell 75: 207-216 (1993); Knowles, et al., NEJM 333(13): 823-831 (1995); and Rosenfeld, et al., Cell 68: 143-155 (1992), the disclosures of which are incorporated by reference herein.)

An Ad-derived vector of the present invention may also comprise a nucleotide sequence encoding a protein, polypeptide or fragment thereof that is effective in regulating the cell cycle -- such as p53, Rb, or mitotin -- or which is effective in inducing cell death, such as thymidine kinase. (See, e.g., published International App. No. WO 95/11984, the disclosures of which are incorporated by reference herein.) It is further contemplated that a therapeutic protein or polypeptide expressed by a therapeutic viral vector of the present invention may be used in conjunction with another therapeutic agent when appropriate -- e.g., a thymidine kinase metabolite may be used in conjunction with the gene encoding thymidine kinase and its gene product -- in order to be even more effective.

Alternatively, a therapeutic viral vector can include a DNA or RNA oligonucleotide sequence that exhibits enzymatic therapeutic activity without needing to be translated into a polypeptide product before exerting a therapeutic effect. Examples of the latter include antisense oligonucleotides that will inhibit the transcription of deleterious genes or ribozymes that act as site-specific ribonucleases for cleaving selected mutated gene sequences. In another variation, a therapeutic nucleotide sequence of the present invention may comprise a DNA construct capable of generating therapeutic nucleotide molecules, including ribozymes and antisense DNA, in high copy numbers in target cells, as described in published PCT application No. WO 92/06693 (the disclosure of which is incorporated herein by reference). Other preferred therapeutic nucleotide sequences according to the present invention are capable of delivering HIV antisense nucleotides to latently-infected T cells via CD4. Similarly, delivery of Epstein-Barr Virus (EBV) EBNA-1 antisense nucleotides to B cells via CR2 is capable of effecting therapeutic results.

As noted elsewhere herein, an Ad-derived vector of the present invention may also include a promoter sequence. Both constitutive and regulatable (often called "inducible") promoters are useful in constructs and methods of the present invention. For example, some useful regulatable promoters are those of the CREB-regulated gene family and include - and - inhibin, - gonadotropin, cytochrome c, glucagon, and the like. (See, e.g., published International App. No. WO96/14061, the disclosures of which are incorporated by reference herein.)

A regulatable or inducible promoter may be described as a promoter wherein the rate of RNA polymerase binding and initiation is modulated by external stimuli. Such stimuli include various compounds or compositions, light, heat, stress, chemical energy sources, and the like. Inducible, suppressible and repressible promoters are considered regulatable promoters.

Regulatable promoters may also include tissue-specific promoters. Tissue-specific promoters direct the expression of the gene to which they are operably linked to a specific cell type. Tissue-specific promoters cause the gene located 3' of it to be expressed predominantly, if not exclusively, in the specific cells where the promoter expressed its endogenous gene. Typically, it appears that if a tissue-specific promoter expresses the gene located 3' of it at all, then it is expressed appropriately in the correct cell types (see, e.g., Palmiter et al., Ann. Rev. Genet. 20: 465-499 (1986)).

When a tissue-specific promoter controls the expression of a gene, that gene will be expressed in a small number of tissues or cell types rather than in substantially all tissues and cell types. Examples of tissue-specific promoters include the immunoglobulin promoter described by Brinster et al., Nature 306: 332-336 (1983) and Storb et al., Nature 310: 238-231 (1984); the elastase-I promoter described by Swift et al., Cell 38: 639-646 (1984); the globin promoter described by Townes et al., Mol. Cell. Biol. 5: 1977-1983 (1985), and Magram et al., Mol. Cell. Biol. 9: 4581-4584 (1989), the insulin promoter described by Bucchini et al., PNAS USA, 83: 2511-2515 (1986) and Edwards et al., Cell 58: 161 (1989); the immunoglobulin promoter described by Ruscon et al., Nature 314: 330-334 (1985) and Grosscheld et al., Cell 38: 647-658 (1984); the alpha actin promoter described by Shani, Mol. Cell. Biol. 6: 2624-2631 (1986); the alpha crystalline promoter described by Overbeek et al., PNAS USA 82: 7815-7819 (1985); the prolactin promoter described by Crenshaw et al., Genes and Development 3: 959-972 (1989); the proopiomelanocortin promoter described by



Tremblay et al., PNAS USA 85: 8890-8894 (1988); the beta-thyroid stimulating hormone (BTSH) promoter described by Tatsumi et al., Nippon Rinsho 47: 2213-2220 (1989); the mouse mammary tumor virus (MMTV) promoter described by Muller et al., Cell 54: 105 (1988); the albumin promoter described by Palmiter et al., Ann. Rev. Genet. 20: 465-499 (1986); the keratin promoter described by Vassar et al., PNAS USA 86: 8565-8569 (1989); the osteonectin promoter described by McVey et al., J. Biol. Chem. 263: 11,111-11,116 (1988); the prostate-specific promoter described by Allison et al., Mol. Cell. Biol. 9: 2254-2257 (1989); the opsin promoter described by Nathans et al., PNAS USA 81: 4851-4855 (1984); the olfactory marker protein promoter described by Danciger et al., PNAS USA 86: 8565-8569 (1989); the neuron-specific enolase (NSE) promoter described by Forss-Pelter et al., J. Neurosci. Res. 16: 141-151 (1986); the L-7 promoter described by Sutcliffe, Trends in Genetics 3: 73-76 (1987) and the protamine 1 promoter described Peschon et al., Ann. New York Acad. Sci. 564: 186-197 (1989) and Braun et al., Genes and Development 3: 793-802 (1989). (The disclosures of all references cited are incorporated by reference herein.)

## 2. Compositions

In various alternative embodiments of the present invention, therapeutic sequences and compositions useful for practicing the therapeutic methods described herein are contemplated. Therapeutic compositions of the present invention may contain a physiologically tolerable carrier together with one or more therapeutic nucleotide sequences of this invention, dissolved or dispersed therein as an active ingredient. In a preferred embodiment, the composition is not immunogenic or otherwise able to cause undesirable side effects when administered to a subject for therapeutic purposes.

As used herein, the terms "pharmaceutically acceptable", "physiologically tolerable" and grammatical variations thereof, as they refer to compositions, carriers, diluents and reagents, are used interchangeably and represent that the materials are capable of administration to or upon a subject -- e.g., a mammal -- without the production of undesirable physiological effects such as nausea, dizziness, gastric upset and the like.

For example, the present invention comprises therapeutic compositions useful in the specific targeting of epithelial or non-epithelial cells as well as in delivering a therapeutic nucleotide sequence to those cells. Therapeutic compositions designed to preferentially target to epithelial cells may comprise an adenovirus-derived vector including a therapeutic

nucleotide sequence. As described herein, a number of adenovirus-derived moieties are useful in the presently-disclosed therapeutic compositions and methods.

While some of the Examples appearing below specifically recite fiber proteins, polypeptides, and fragments thereof, it is expressly provided herein that other structural and non-structural Ad proteins and polypeptides (e.g., regulatory proteins and polypeptides) may be used as components of the various disclosed vectors and cell lines. Moreover, chimeric molecules comprised of proteins, polypeptides, and/or fragments thereof which are derived from different Ad serotypes may be used in any of the within-disclosed methods, constructs and compositions. Similarly, recombinant DNA sequences of the present invention may be prepared using nucleic acid sequences derived from different Ad serotypes, in order to design useful constructs with broad applicability, as disclosed and claimed herein.

It should also be appreciated that, while the members of Group C adenovirus -- i.e., Ad serotypes 1, 2, 5, and 6 -- are specifically recited in various examples herein, the present invention is in no way limited to those serotypes alone. In view of the fact that the adenovirus serotypes are all closely-related in structure and functionality, therapeutic viral vectors, packaging cell lines, and plasmids of the present invention may be constructed from components of any and all Ad serotypes -- and the within-disclosed methods of making and using the various constructs and cell lines of the present invention apply to all of said serotypes.

The preparation of a pharmacological composition that contains active ingredients dissolved or dispersed therein is well understood in the art. Typically such compositions are prepared as injectables -- either as liquid solutions or suspensions -- however, solid forms suitable for solution or suspension in liquid prior to use can also be prepared. A preparation can also be emulsified, or formulated into suppositories, ointments, creams, dermal patches, or the like, depending on the desired route of administration.

The active ingredient can be mixed with excipients which are pharmaceutically acceptable and compatible with the active ingredient and in amounts suitable for use in the therapeutic methods described herein. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol or the like and combinations thereof, including vegetable oils, propylene glycol, polyethylene glycol and benzyl alcohol (for injection or liquid preparations); and petrolatum (e.g., VASELINE), vegetable oil, animal fat and polyethylene glycol (for

externally applicable preparations). In addition, if desired, the composition can contain wetting or emulsifying agents, isotonic agents, dissolution promoting agents, stabilizers, colorants, antiseptic agents, soothing agents and the like additives (as usual auxiliary additives to pharmaceutical preparations), pH buffering agents and the like which enhance the effectiveness of the active ingredient.

The therapeutic compositions of the present invention can include pharmaceutically acceptable salts of the components therein. Pharmaceutically acceptable salts include the acid addition salts (formed with the free amino groups of the polypeptide) that are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, tartaric, mandelic and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine and the like.

Physiologically tolerable carriers are well known in the art. Exemplary of liquid carriers are sterile aqueous solutions that contain no materials in addition to the active ingredients and water, or contain a buffer such as sodium phosphate at physiological pH value, physiological saline or both, such as phosphate-buffered saline. Still further, aqueous carriers can contain more than one buffer salt, as well as salts such as sodium and potassium chlorides, dextrose, polyethylene glycol and other solutes.

Liquid compositions can also contain liquid phases in addition to and to the exclusion of water. Exemplary of such additional liquid phases are glycerin, vegetable oils such as cottonseed oil, and water-oil emulsions.

A therapeutic composition typically contains an amount of a therapeutic nucleotide sequence of the present invention sufficient to deliver a therapeutically effective amount to the target tissue, typically an amount of at least 0.1 weight percent to about 90 weight percent of therapeutic nucleotide sequence per weight of total therapeutic composition. A weight percent is a ratio by weight of therapeutic nucleotide sequence to total composition. Thus, for example, 0.1 weight percent is 0.1 grams of DNA segment per 100 grams of total composition.

The therapeutic nucleotide compositions comprising synthetic oligonucleotide sequences of the present invention can be prepared using any suitable method, such as, the phosphotriester or phosphodiester methods. See Narang et al., Meth. Enzymol., 68:90,

(1979); U.S. Patent No. 4,356,270; and Brown et al., Meth. Enzymol., 68:109, (1979), the disclosures of which are incorporated by reference herein.

For therapeutic oligonucleotides sequence compositions in which a family of variants is preferred, the synthesis of the family members can be conducted simultaneously in a single reaction vessel, or can be synthesized independently and later admixed in preselected molar ratios. For simultaneous synthesis, the nucleotide residues that are conserved at preselected positions of the sequence of the family member can be introduced in a chemical synthesis protocol simultaneously to the variants by the addition of a single preselected nucleotide precursor to the solid phase oligonucleotide reaction admixture when that position number of the oligonucleotide is being chemically added to the growing oligonucleotide polymer. The addition of nucleotide residues to those positions in the sequence that vary can be introduced simultaneously by the addition of amounts, preferably equimolar amounts, of multiple preselected nucleotide precursors to the solid phase oligonucleotide reaction admixture during chemical synthesis. For example, where all four possible natural nucleotides (A,T,G and C) are to be added at a preselected position, their precursors are added to the oligonucleotide synthesis reaction at that step to simultaneously form four variants.

This manner of simultaneous synthesis of a family of related oligonucleotides has been previously described for the preparation of "Degenerate Oligonucleotides" by Ausubel et al. (Current Protocols in Molecular Biology, Suppl. 8, p.2.11.7, John Wiley & Sons, Inc., New York (1991)), and can readily be applied to the preparation of the therapeutic oligonucleotide compositions described herein.

Nucleotide bases other than the common four nucleotides (A,T,G or C), or the RNA equivalent nucleotide uracil (U), can also be used in the present invention. For example, it is well known that inosine (I) is capable of hybridizing with A, T and G, but not C. Examples of other useful nucleotide analogs are known in the art; many may be found listed in 37 C.F.R. §1.822.

Thus, where all four common nucleotides are to occupy a single position of a family of oligonucleotides, that is, where the preselected therapeutic nucleotide composition is designed to contain oligonucleotides that can hybridize to four sequences that vary at one position, several different oligonucleotide structures are contemplated. The composition can contain four members, where a preselected position contains A,T,G or C. Alternatively, the

composition can contain two members, where a preselected position contains I or C, and has the capacity to hybridize at that position to all four possible common nucleotides. Finally, other nucleotides may be included at the preselected position that have the capacity to hybridize in a non-destabilizing manner with more than one of the common nucleotides in a manner similar to inosine.

### 3. Expression Vector Systems

The introduction of exogenous DNA into eucaryotic cells has become one of the most powerful tools of the molecular biologist. The term "exogenous" encompasses any therapeutic composition of this invention which is administered by the therapeutic methods of this invention. Thus, "exogenous" may also be referred to herein as "foreign," "non-native," and the like. The methods of this invention preferably require efficient delivery of the DNA into the nucleus of the recipient cell and subsequent identification of cells that are expressing the foreign DNA.

A widely-used plasmid is pBR322, a vector whose nucleotide sequence and endonuclease cleavage sites are well known. Various other useful plasmid vectors are described in the Examples that follow.

A vector of the present invention comprises a nucleic acid (preferably DNA) molecule capable of autonomous replication in a cell and to which a DNA segment, e.g., a gene or polynucleotide, can be operatively linked so as to bring about replication of the attached segment. In the present invention, one of the nucleotide segments to be operatively linked to vector sequences encodes at least a portion of a therapeutic nucleic acid molecule -- in effect, a nucleic acid sequence that encodes one or more therapeutic proteins or polypeptides, or fragments thereof.

In various embodiments, the entire peptide-coding sequence of the therapeutic gene is inserted into the vector and expressed; however, it is also feasible to construct a vector which also includes some non-coding sequences as well. Preferably, however, non-coding sequences are excluded. Alternatively, a nucleotide sequence for a soluble form of a polypeptide may be utilized. Another preferred therapeutic viral vector includes a nucleotide sequence encoding at least a portion of a therapeutic nucleotide sequence operatively linked to the vector for expression. As used herein with regard to DNA sequences or segments, the phrase

"operatively linked" generally means the sequences or segments have been covalently joined into one piece of DNA, whether in single or double stranded form.

The choice of viral vector into which a therapeutic nucleotide sequence of this invention is operatively linked depends directly, as is well known in the art, on the functional properties desired, e.g., vector replication and protein expression, and the host cell to be transformed -- these being limitations inherent in the art of constructing recombinant DNA molecules. Although certain adenovirus serotypes are recited herein in the form of specific examples, it should be understood that the present invention contemplates the use of *any* adenovirus serotype, including hybrids and derivatives thereof. As one will observe, it is not unusual or outside the scope of the present invention to utilize nucleotide and/or amino acid residue sequences of two or more serotypes in constructs, compositions and methods of the invention.

As one of skill in the art will note, in various embodiments of the present invention, different "types" of vectors are disclosed. For example, one "type" of vector is used to deliver particular nucleotide sequences into a packaging cell line, with the intent of having said sequences stably integrate into the cellular genome; these "types" of vectors are generally identified herein as complementing plasmids. A further "type" of vector described herein carries or delivers nucleotide sequences in or into a cell line (e.g., a packaging cell line) for the purpose of propagating therapeutic viral vectors of the present invention; hence, these vectors are generally referred to herein as delivery plasmids. A third "type" of vector described herein is utilized to carry nucleotide sequences encoding therapeutic proteins or polypeptides to specific cells or cell types in a subject in need of treatment; these vectors are generally identified herein as therapeutic viral vectors or Ad-derived vectors.

In one embodiment, the directional ligation means is provided by nucleotides present in the upstream nucleotide sequence, downstream nucleotide sequence, or both. In another embodiment, the sequence of nucleotides adapted for directional ligation comprises a sequence of nucleotides that defines multiple directional cloning means. Where the sequence of nucleotides adapted for directional ligation defines numerous restriction sites, it is referred to as a multiple cloning site.

A translatable nucleotide sequence is a linear series of nucleotides that provide an uninterrupted series of at least 8 codons that encode a polypeptide in one reading frame.

Preferably, the nucleotide sequence is a DNA sequence. The vector itself may be of any suitable type, such as a viral vector (RNA or DNA), naked straight-chain or circular DNA, or a vesicle or envelope containing the nucleic acid material and any polypeptides that are to be inserted into the cell.

A preferred viral vector in which therapeutic nucleotide compositions of this invention are present is derived from adenovirus (Ad). It is also desirable that the vector contain a promoter sequence. As taught herein, viral vectors of this invention may be designed and constructed in such a way that they specifically target a preselected recipient cell type, depending on the nature of therapy one seeks to administer. Methods of making and using therapeutic viral vectors that target specific cells are further described in the Examples that follow.

Novel vectors and compositions may also be designed and prepared to preferentially target cells that might not otherwise be targeted by wild-type adenovirus virions. For example, in order to target non-epithelial cells, one following the teachings of the present specification may be able to prepare a therapeutic vector including a nucleotide sequence encoding a foreign protein, polypeptide or other ligand directed to a non-epithelial cell or to a different receptor than that generally targeted by a particular adenovirus. Examples of useful ligands directed to specific receptors (identified in parentheses) include the V3 loop of HIV gp120 (CD4); transferrin (transferrin receptor); LDL (LDL receptors); and deglycosylated proteins (asialoglycoprotein receptor). Various useful ligands which may be added to adenovirus fiber - and methods for preparing and attaching same -- are set forth in published International App. No. WO95/26412, the disclosures of which are incorporated by reference herein.

Useful ligands which may be encoded by a foreign nucleotide sequence contained within a viral vector of the present invention, or which may be linked to proteins or polypeptides expressed thereby after said vectors are administered to a subject, also include antibodies and attachment sequences, as well as receptors themselves. For example, antibodies to cell receptor molecules such as integrins and the like, MHC Class I and Class II, asialoglycoprotein receptor, transferrin receptors, LDL receptors, CD4, and CR2 are but a few which are useful according to the present invention. It is also understood that the ligands typically bound by receptors, as well as analogs to those ligands, may be used as cellular targeting agents, as disclosed herein.

#### E. Therapeutic Methods

The vectors of the present invention are particularly suited for gene therapy. Thus, various therapeutic methods are contemplated by the present invention.

For example, it has now been discovered that Ad-derived viral vectors are capable of delivering a therapeutic nucleotide sequence to a specific cell or tissue, thereby expanding and enhancing treatment options available in numerous conditions in which more conventional therapies are of limited efficacy. Accordingly, methods of gene therapy utilizing these vectors are within the scope of the invention. Vectors are typically purified and then an effective amount is administered *in vivo* or *ex vivo* into the subject.

For example, the compositions may be used prophylactically or therapeutically *in vivo* to disrupt HIV infection and mechanisms of action by inhibiting gene expression or activation, via delivery of antisense HIV sequences or ribozymes to T cells or monocytes. Using methods of the present invention, one may target therapeutic viral vectors as disclosed herein to specific cells and tissues, including hematopoietic cells, as infection of such cells appears to be mediated by distinct integrins to which viral vectors of the present invention may readily be targeted. (See, e.g., Huang, et al., J. Virol. 70: 4502-8 (1996).)

Other useful therapeutic nucleotide sequences include antisense nucleotide sequences complementary to EBV EBNA-1 gene. Use of such therapeutic sequences may remediate or prevent latent infection of B cells with EBV. As discussed herein and in the Examples below, targeting and delivery may be accomplished via the use of various ligands, receptors, and other appropriate targeting agents.

Thus, in one embodiment, a therapeutic method of the present invention comprises contacting the cells of a subject infected with EBV or HIV with a therapeutically effective amount of a pharmaceutically acceptable composition comprising a therapeutic nucleotide sequence of this invention. In a related embodiment, the contacting involves introducing the therapeutic nucleotide sequence composition into cells having an EBV or HIV-mediated infection.

Methods of gene therapy are well known in the art (see, e.g., Larrick and Burck, Gene Therapy: Application of Molecular Biology, Elsevier Science Publ. Co., Inc., New York, NY (1991); Kriegler, Gene Transfer and Expression: A Laboratory Manual, W. H. Freeman and Company, New York (1990)). The term "subject" should be understood to include any animal



-- particularly mammalian -- patient, such as any murine, rat, bovine, porcine, canine, feline, equine, ursine, or human patient.

When the foreign gene carried in the vector encodes a tumor suppressor gene or another anti-tumor protein, the vector is useful to treat or reduce hyperproliferative cells in a subject, to inhibit tumor proliferation in a subject or to ameliorate a particular, related pathology. Pathologic hyperproliferative cells are characteristic of various disease states, such as thyroid hyperplasia, psoriasis, eczema, benign prostatic hypertrophy, Li-Fraumeni syndrome including breast cancer, sarcomas and other neoplasms, bladder cancer, colon cancer, lung cancer, various leukemias, and lymphomas.

Non-pathologic hyperproliferative cells are found, for example, in cells associated with wound repair. Pathologic hyperproliferative cells, however, characteristically exhibit loss of contact inhibition and a decline in their ability to selectively adhere which implies a change in the surface properties of the cell and a further breakdown in intercellular communication. These changes include stimulation to divide and the ability to secrete proteolytic enzymes.

The present invention also contemplates methods of depleting suitable samples of pathologic mammalian hyperproliferative cells contaminating hematopoietic precursors during bone marrow reconstitution via the introduction of a wild-type tumor suppressor gene into the cell preparation using a vector of this invention. As used herein, a suitable sample is defined as a heterogeneous cell preparation obtained from a patient, e.g., a mixed population of cells containing both phenotypically normal and pathogenic cells.

Administration includes -- but is not limited to -- the introduction of therapeutic agents of the present invention into a cell or subject via various means, including direct injection, intravenously, intraperitoneally, via intra-tumor injection, via aerosols, or topically. Therapeutic agents as disclosed herein may also be combined for administration of an effective amount of the agents with a pharmaceutically-acceptable carrier, as described herein.

As used herein, "effective amount" generally means the amount of vector (or proteins produced/released thereby) which achieves a positive outcome in the subject to whom the vector is administered. The total volume administered will necessarily vary depending on the mode of administration, as those of skill in the relevant art will appreciate, and dosages may vary as well.

The dose of a biologic vector is somewhat complex and may be described in terms of the concentration (in plaque-forming units per milliliter (pfu/ml)), the total dose (in pfus), and the estimated number of vectors administered per cell (the estimated multiplicity of infection or MOI). Thus, if a vector is administered via infusion -- say, across nasal epithelium -- at a constant total volume, the respective concentration, etc. may be described as follows:

Concentration (pfu/ml)	Volume (ml)	Dose (pfu)	Estimated MOI
$10^7$	2	$2 \times 10^7$	1
$10^8$	2	$2 \times 10^8$	10
$10^9$	2	$2 \times 10^9$	100
$10^{10}$	2	$2 \times 10^{10}$	1000

In general, when adenoviral vectors are administered via infusion across the nasal epithelium, administered amounts producing an estimated MOI of about 10 or greater are much more effective than lower dosages. (See, e.g., Knowles, et al., New Eng. J. Med. 333: 823-831 (1995).) Similarly, when direct injection is the preferred treatment modality -- e.g., direct injection of a viral vector into a tumor -- doses of  $1 \times 10^9$  pfu or greater are generally preferred. (See, e.g., published International App. No. WO95/11984.)

Thus, depending on the mode of administration, an effective amount administered in a single dose preferably contains from about  $10^6$  to about  $10^{15}$  infectious units. A typical course of treatment would be one such dose per day over a period of five days. As those of skill in the art will appreciate, an effective amount may vary depending on (1) the pathology or other condition to be treated, (2) the status and sensitivity of the patient, and (3) various other factors well known to those of skill in the art, such as the patient's tolerance to other courses of treatment that may have been applied previously. Thus, those of skill in the art may easily and precisely determine effective amounts of the agents/vectors of the present invention which may be administered to a particular patient, based on their understanding of and evaluation of such factors.

The present invention also contemplates methods of ameliorating pathologies characterized by hyperproliferative cells or genetic defects in a subject, by administering to the subject an effective amount of a vector as described herein. Such vectors preferably contain a foreign gene encoding a gene product (e.g. polypeptide or protein) having the ability to ameliorate the pathology, under suitable conditions. As used herein, the term "genetic defect" means any disease, condition or abnormality which results from inherited factors, e.g. Huntington's Disease, Tay-Sachs Disease, or Sickle Cell Disease.

The present invention further provides methods for reducing the proliferation of tumor cells in a subject by introducing into the tumor mass an effective amount of an adenoviral expression vector containing an anti-tumor gene other than a tumor suppressor gene. The anti-tumor gene can encode, for example, thymidine kinase (TK). An effective amount of a therapeutic agent is then administered to the subject; the therapeutic agent, in the presence of the anti-tumor gene, is toxic to the cell.

Using thymidine kinase as exemplary, the therapeutic agent is a thymidine kinase metabolite such as ganciclovir (GCV), 6-methoxypurine arabinonucleoside (araM), or a functional equivalent thereof. Both the thymidine kinase gene and the thymidine kinase metabolite must be used concurrently in order to exert a toxic effect on the host cell. In the presence of the TK gene, GCV is phosphorylated and becomes a potent inhibitor of DNA synthesis, whereas araM is converted to the cytotoxic anabolite araATP. Thus, the precise method of action or synergism is not relevant to therapeutic efficacy; what is relevant is the fact that the concurrent use of appropriate genes and therapeutic agents may effectively ameliorate a specific disease condition.

Another useful example contemplates use of a vector of the present invention which expresses the enzyme cytosine deaminase. Such a vector could be used in conjunction with administration of the drug 5-fluorouracil (Austin and Huber, Mol. Pharm. 43: 380-387 (1993)) or the recently-described *E. coli* Deo gene in combination with 6-methyl-purine-2'-deoxyribonucleoside (Sorscher et al., Gene Therapy 1: 233-238 (1994)).

As with the use of the tumor suppressor genes described previously, the use of other anti-tumor genes, either alone or in combination with the appropriate therapeutic agent, provides a treatment for the uncontrolled cell growth or proliferation characteristic of tumors and malignancies. Thus, the present invention provides therapies to halt the uncontrolled

cellular growth in a patient, thereby alleviating the symptoms or the disease or cachexia present in the patient. The effect of this treatment includes, but is not limited to, prolonged survival time of the patient, reduction in tumor mass or burden, apoptosis of tumor cells, or the reduction in the number of circulating tumor cells. Means of quantifying the beneficial effects of this therapy are well known to those of skill in the art.

The present invention provides a recombinant adenovirus expression vector characterized by the partial or total deletion of one or more adenoviral structural protein genes, such as the gene encoding fiber, which allows the vector to accommodate a therapeutic, foreign nucleic acid sequence encoding a functional foreign polypeptide, protein, or biologically active fragment thereof. For example, such a functional polypeptide moiety may be a suicide gene or a functional equivalent thereof, of which the anti-cancer gene TK is but one example. TK genes, when expressed, produce a gene product which is lethal to the cell, particularly in the presence of GCV. One source of the TK gene is the herpes simplex virus (HSV), albeit other sources are known as well and may be used as taught herein. The TK gene may readily be obtained from HSV by methods well known to those of skill in the art. For example, the plasmid pMLBTK in *E. coli* HB101 (from ATCC #39369) is a source of the HSV-1 TK gene, which may be used as disclosed herein. (See, e.g., published International application No. WO 95/11984, the disclosures of which are incorporated by reference herein.)

A therapeutic gene sequence may be introduced into a tumor mass by combining the adenoviral expression vector with a suitable pharmaceutically acceptable carrier. Introduction can be accomplished, for example, via direct injection of the recombinant Ad vector into the tumor mass. In the specific case of a cancer such as hepatocellular carcinoma (HCC), direct injection into the hepatic artery can be used for delivery, because most HCCs derive their circulation from this artery. Similar techniques of administration may be applied to other specific types of tumors and malignancies, as is known to those of skill in the art.

A method of tumor-specific delivery of a tumor-suppressor gene is accomplished by contacting target tissue in a subject with an effective amount of a recombinant Ad-derived vector of this invention. In the case of anti-tumor therapy, the gene is intended to encode an anti-tumor agent, such as a functional tumor suppressor gene product or suicide gene product. The term "contacting" is intended to encompass any delivery method for the efficient transfer of the vector, such as via intra-tumoral injection.

In another example, adenovirus vectors of the present invention can be used to transfer genes to central nervous system (CNS) tumors *in vivo*. Using stereotactic delivery, Ad-derived vectors can transfer genes into the CNS intended for tumor therapy. For example, Badie, et al. (Neurosurgery 35(5): 910-916 (1994), incorporated by reference herein) reported that 50% and 90% transduction at vector titers of approximately  $10^7$  and  $10^8$  plaque-forming units/ml (pfu/ml) were observed in *in vitro* experiments. In their *in vivo* studies using appropriate animal brain tumor models, titers above  $10^7$  were observed to have a cytopathic effect; more than 50% reduction in tumor cell growth was noted at  $10^8$  pfu/ml; no toxic effects were noted when titers as high as  $10^{10}$  pfu/ml were injected into the brain tissue of subject animals (*Id.*). Thus, the use of titers greater than  $10^7$  pfu/ml appear appropriate when challenging CNS tumors.

The present invention also contemplates methods for determining the efficacy of the within-disclosed therapeutic compositions and methods. One such method for confirming efficacy utilizes the human/SCID (severe combined immunodeficient) mouse model of EBV-induced LPD (lymphoproliferative disease) to ascertain whether EBV-antisense therapeutic nucleotide sequences block tumor formation. (See, e.g., Pisa, et al., Blood 79: 173-179 (1992); Rowe, et al., Curr. Top. Microbiol. Immunol. 166: 325 (1990); and Cannon, et al., J. Clin. Invest. 85: 1333-1337 (1990), the disclosures of which are incorporated by reference herein.)

Finally, the use of Ad vectors of the present invention to prepare medicaments for the treatment, therapy and/or diagnosis of various diseases is also contemplated by this invention. Moreover, other anti-tumor genes may be used in combination with the corresponding therapeutic agent to reduce the proliferation of tumor cells. Such other gene-and-therapeutic-agent combinations are known to those of skill in the art and may be applied as taught herein.

#### F. Construction of Therapeutic Viral Vectors for Gene Delivery

For *in vitro* gene transfer, administration is often accomplished by first isolating a selected cell population from a patient such as lung epithelial cells, lymphocytes and the like followed by *in vitro* gene transfer of the therapeutic compositions of this invention and the replacement of the cells into the patient. *In vivo* therapy is also contemplated, e.g., via the administration of therapeutic compositions of this invention by various delivery means. For

example, aerosol administration and administration via subcutaneous, intravenous, intraperitoneal, intramuscular, ocular means and the like are also within the scope of the present invention.

Other gene-delivery methods are also useful in conjunction with the methods, compositions and constructs of the present invention; see, e.g., published International Application No. WO 95/11984, the disclosures of which are incorporated by reference herein.

Similarly, various non-human animals having inserted therein the vectors or transformed cells of this invention. These "transgenic" animals are made using methods well known to those of skill in the art. For example, see U.S. Patent No. 5,175,384 (the disclosures of which are incorporated by reference herein).

The present invention also contemplates various methods of targeting specific cells -- e.g., cells in a subject in need of diagnosis and/or treatment. As discussed herein, the present invention contemplates that the viral vectors and compositions of the present invention may be directed to specific receptors or cells, for the ultimate purpose of delivering those vectors and compositions to specific cells or cell types. The viral vectors and constructs of the present invention are particularly useful in this regard.

In general, adenovirus attachment and uptake into cells are separate but cooperative events that result from the interaction of distinct viral coat proteins with a receptor for attachment and  $\gamma$  integrin receptors for internalization. Adenovirus attachment to the cell surface via the fiber coat proteins has been discovered to be dissociable and distinct from the subsequent step of internalization, and the present invention is able to take advantage of and function in conjunction with these differing receptors.

#### G. Other Applications

The cell lines, viral vectors and methods of the present invention may also be used for purposes other than the direct administration of therapeutic nucleotide sequences. In one such application, the production of large quantities of biologically active proteins or polypeptides in cells transfected with the within-disclosed viral vectors is contemplated herein. For example, human lymphoblastoid cells may be transfected with an integrative viral vector of the present invention carrying a human hematopoietic growth factor such as the gene for erythropoietin (EPO); cells so transfected are thus able to produce biologically active EPO. (See, e.g., Lopez et al., Gene 148: 285-91 (1994).)

Various other applications and uses of the within-described methods, cell lines, plasmids, vectors, and compositions of the present invention shall become apparent upon closer examination of the Examples that follow.

### EXAMPLES

The following examples are intended to illustrate, but not limit, the present invention. As such, the following description provides details of the manner in which particular embodiments of the present invention may be made and used. This description, while exemplary of the present invention, is not to be construed as specifically limiting the invention. Variations and equivalents, now known or later developed, which would be within the understanding and technical competence of one skilled in this art are to be considered as falling within the scope of this invention.

#### **Example 1**

##### Preparation of Adenovirus Packaging Cell Lines

Cell lines that are commonly used for growing adenovirus are useful as host cells for the preparation of adenovirus packaging cell lines. Preferred cells include 293 cells, an adenovirus-transformed human embryonic kidney cell line obtained from the ATCC, having Accession Number CRL 1573; HeLa, a human epithelial carcinoma cell line (ATCC Accession Number CCL 2); A549, a human lung carcinoma cell line (ATCC Accession Number CCL 1889); and the like epithelial-derived cell lines. As a result of the adenovirus transformation, the 293 cells contain the E1 early region regulatory gene. All cells were maintained in complete DMEM + 10% fetal calf serum unless otherwise noted.

The cell lines of this invention allow for the production and propagation of novel adenovirus-based gene delivery vectors having deletions in preselected gene regions by cellular complementation of adenoviral genes. To provide the desired complementation of such deleted adenoviral genomes in order to generate a novel viral vector of the present invention, plasmid vectors that contain preselected functional units were designed as described herein. Such units include but are not limited to E1 early region, ????? and the viral fiber gene. The preparation of plasmids providing such complementation, thereby being "complementary plasmids or constructs", that are stably inserted into host cell chromosomes are described below.

A. Preparation of an E4-Expressing Plasmid for Complementation of E4-Gene-Deleted Adenoviruses

The viral E4 regulatory region contains a single transcription unit which is alternately spliced to produce several different mRNAs. The E4-expressing plasmid prepared as described herein and used to transfect the 293 cell line contains the entire E4 transcriptional unit as shown in Figure 1. A DNA fragment extending from 175 nucleotides upstream of the E4 transcriptional start site including the natural E4 promoter to 153 nucleotides downstream of the E4 polyadenylation signal including the natural E4 terminator signal, corresponding to nucleotides 32667-35780 of the adenovirus type 5 (hereinafter referred to as Ad5) genome as described in Chroboczek et al. (*Virology*, 186:280-285 (1992), GenBank Accession Number M73260), was amplified from Ad5 genomic DNA, obtained from the ATCC, via the polymerase chain reaction (PCR). Sequences of the primers used were 5'CGGTACACAGAAATTCAGGAGACACAACTCC3' (forward or 5' primer referred to as E4L) (SEQ ID NO 1) and 5'GCCTGGATCCGGGAAGTTACGTAACGTGGGAAAAC3' (SEQ ID NO 2) (backward or 3' primer referred to as E4R). To facilitate cloning of the PCR fragment, these oligonucleotides were designed to create novel sites for the restriction enzymes EcoRI and BamHI, respectively, as indicated with underlined nucleotides. DNA was amplified via PCR using 30 cycles of 92 C for 1 minute, 50 C for 1 minute, and 72 C for 3 minutes resulting in amplified full-length E4 gene products.

The amplified DNA E4 products were then digested with EcoRI and BamHI for cloning into the compatible sites of pBluescript/SK+ by standard techniques to create the plasmid pBS/E4. A 2603 base pair (bp) cassette including the herpes simplex virus thymidine kinase promoter, the hygromycin resistance gene, and the thymidine kinase polyadenylation signal was excised from the plasmid pMEP4 (Invitrogen, San Diego, CA) by digestion with FspI followed by addition of BamHI linkers (5'CGCGGATCCGCG3') (SEQ ID NO 3) for subsequent digestion with BamHI to isolate the hygromycin-containing fragment.

The isolated BamHI-modified fragment was then cloned into the BamHI site of pBS/E4 containing the E4 region to create the plasmid pE4/Hygro containing 8710 bp (Figure 2). The pE4/Hygro plasmid has been deposited with the ATCC as described in Example 3. The complete nucleotide sequence of pE4/Hygro is listed in SEQ ID NO 4. Position number 1 of the linearized vector corresponds to approximately the middle portion of the pBS/SK+



backbone as shown in Figure 2 as a thin line between the 3' BamHI site in the hygromycin insert and the 3' EcoRI site in the E4 insert. The 5' and 3' ends of the E4 gene are located at respective nucleotide positions 3820 and 707 of SEQ ID NO 4 while the 5' and 3' ends of the hygromycin insert are located at respective nucleotide positions 3830 and 6470. In the clone that was selected for use, the E4 and hygromycin resistance genes were divergently transcribed.

B. Preparation of a Fiber-Expressing Plasmid for Complementation of Fiber-Gene-Deleted Adenoviruses

To prepare a fiber-encoding construct, primers were designed to amplify the fiber coding region from Ad5 genomic DNA with the addition of unique BamHI and NotI sites at the 5' and 3' ends of the fragment, respectively. The Ad5 nucleotide sequence is available with the GenBank Accession Number M18369. The 5' and 3' primers had the respective nucleotide sequences of 5' ATGGGATCCAAGATGAAGCGCGCAAGACCG3' (SEQ ID NO 5) and 5' CATAACGCGGCCGCTTCTTTATTCTTGGGC3' (SEQ ID NO 6), where the inserted BamHI and NotI sites are indicated by underlining. The 5' primer also contained a nucleotide substitution 3 nucleotides 5' of the second ATG codon (C to A) that is the initiation site. The nucleotide substitution was included so as to improve the consensus for initiation of fiber protein translation.

The amplified DNA fragment was inserted into the BamHI and NotI sites of pcDNA 3 (Invitrogen) to create the plasmid designated pCDNA3/Fiber having 7148 bp, the plasmid map of which is shown in Figure 3. The parent plasmid contained the CMV promoter, the bovine growth hormone (BHG) terminator and the gene for conferring neomycin resistance. The viral sequence included in this construct corresponds to nucleotides 31040-32791 of the Ad5 genome.

The complete nucleotide sequence of pCDNA3/Fiber is listed in SEQ ID NO 7 where the nucleotide position 1 corresponds to approximately the middle of the pcDNA 3 vector sequence. The 5' and 3' ends of the fiber gene are located at respective nucleotide positions 916 with ATG and 2661 with TAA.

To enhance expression of fiber protein by the constitutive CMV promoter provided by the pcDNA vector, a BglII fragment containing the tripartite leader (TPL) of adenovirus type 2 was excised from pRD112a (Sheay et al., BioTechniques, 15:856-862 (1993) and inserted into

the BamHI site of pCDNA3/Fiber to create the plasmid pCLF having 7469 bp, the plasmid map of which is shown in Figure 4. The adenovirus tripartite leader sequence, present at the 5' end of all major late adenoviral mRNAs as described by Logan et al., Proc. Natl. Acad. Sci., USA, 81:3655-3659 (1984) and Berkner, BioTechniques, 6:616-629 (1988), is encoded by three spatially separated exons corresponding to nucleotide positions 6071-6079 (the 3' end of the first leader segment), 7101-7172 (the entire second leader segment), and 9634-9721 (the third leader segment) in the adenovirus type 2 genome. The tripartite sequence, however, also shows correspondence with the Ad5 leader sequence having three spatially separated exons corresponding to nucleotide positions 6081-6089 (the 3' end of the first leader segment), 7111-7182 (the entire second leader segment), and 9644-9845 (the third leader segment and sequence downstream of that segment). The corresponding cDNA sequence of the tripartite leader sequence present in pCLF is listed in SEQ ID NO 8 bordered by BamHI/BglII 5' and 3' sites at respective nucleotide positions 907-912 to 1228-1233.

The pCLF plasmid has been deposited with the ATCC as described in Example 3. The complete nucleotide sequence of pCLF is listed in SEQ ID NO 8 where the nucleotide position 1 corresponds to approximately the middle of the pcDNA 3 parent vector sequence. The 5' and 3' ends of the Ad5 fiber gene are located at respective nucleotide positions 1237-1239 with ATG and 2980-2982 with TAA. The rest of the vector construct has been previously described above.

#### C. Generation of an Adenovirus Packaging Cell Line Carrying Plasmids Encoding Functional E4 and Fiber Proteins

The 293 cell line was selected for preparing the first adenovirus packaging line as it already contains the E1 gene as prepared by Graham et al., J. Gen. Virol., 36:59-74 (1977) and as further characterized by Spector, Virology, 130:533-538 (1983). Before electroporation, 293 cells were grown in RPMI medium + 10% fetal calf serum. Four x 10<sup>6</sup> cells were electroporated with 20 µg each of pE4/Hygro DNA and pCLF DNA using a BioRad GenePulser and settings of 300 V, 25 µF. DNA for electroporation was prepared using the Qiagen system according to the manufacturer's instructions (Bio-Rad, Richmond, CA).

Following electroporation, cells were split into fresh complete DMEM + 10% fetal calf serum containing 200 µg/ml Hygromycin B (Sigma, St. Louis, MO).

From expanded colonies, genomic DNA was isolated using the "MICROTURBOGEN" system (Invitrogen) according to manufacturer's instructions. The presence of integrated E4 DNA was assessed by PCR using the primer pair E4R and ORF6L (5'TGCTTAAGCGGCCGCGAAGGAGAAGTCC3') (SEQ ID NO 9), the latter of which is a 5' forward primer near adenovirus 5 open reading frame 6. Refer to Figure 1 for position of the primers relative to the E4 genes.

One clone, designated 211, was selected exhibiting altered growth properties relative to that seen in parent cell line 293. The 211 clone contained the expected product, indicating the presence of inserted DNA corresponding to most, if not all, of the E4 fragment contained in the pE4/Hygro plasmid. The 211 cell line has been deposited with the ATCC as described in Example 3. This line was further evaluated by amplification using the primer pair E4L/E4R described above, and a product corresponding to the full-length E4 insert was detected. Genomic Southern blotting was performed on DNA restricted with EcoRI and BamHI. The E4 fragment was then detected at approximately one copy/genome compared to standards with the EcoRI/BamHI E4 fragment as cloned into pBS/E4 for use as a labeled probe with the Genius system according to manufacturer's instructions (Boehringer Mannheim, Indianapolis, IN). In DNA from the 211 cell line, the expected labeled internal fragment pE4/Hygro hybridized with the isolated E4 sequences. In addition, the probe hybridized to a larger fragment which may be the result of a second insertion event (Figure 5).

Although the 211 cell line was not selected by neomycin resistance, thus indicating the absence of fiber gene, to confirm the lack of fiber gene, the 211 cell line was analyzed for expression of fiber protein by indirect immunofluorescence with an anti-fiber polyclonal antibody and a FITC-labeled anti-rabbit IgG (KPL) as secondary. No immunoreactivity was detected. Therefore, to generate 211 clones containing recombinant fiber genes, the 211 clone was expanded by growing in RPMI medium and subjected to additional electroporation with the fiber-encoding pCLF plasmid as described above.

Following electroporation, cells were plated in DMEM + 10% fetal calf serum and colonies were selected with 200 µg/ml G418 (Gibco, Gaithersburg, MD). Positive cell lines remained hygromycin resistant. These candidate sublines of 211 were then screened for fiber protein expression by indirect immunofluorescence as described above. The three sublines screened, 211A, 211B and 211R, along with a number of other sublines, all exhibited nuclear

staining qualitatively comparable to the positive control of 293 cells infected with AdRSV gal (1 pfu/cell) and stained 24 hours post-infection.

Lines positive for nuclear staining in this assay were then subjected to Western blot analysis under denaturing conditions using the same antibody. Several lines in which the antibody detected a protein of the expected molecular weight (62 kd for the Ad5 fiber protein) were selected for further study including 211A, 211B and 211R. The 211A cell line has been deposited with ATCC as described in Example 3.

Western blot analysis using soluble nuclear extracts from these three cell lines and a semipreparative electrophoresis system demonstrated that the fiber protein expressed is in the functional trimeric form characteristic of the native fiber protein as shown in Figure 6. The predicted molecular weight of a trimerized fiber is 186 kd. The lane marked 293 lacks fiber while the sublines contain detectable fiber. Under denaturing conditions, the trimeric form was destroyed resulting in detectable fiber monomers as shown in Figure 6. Those clones containing endogenous E1, newly expressed recombinant E4 and fiber proteins were selected for use in complementing adenovirus gene delivery vectors having the corresponding adenoviral genes deleted as described in Example 2.

D. Preparation of an E1-Expressing Plasmid for Complementation of E1-Gene-Deleted Adenoviruses

In order to prepare adenoviral packaging cell lines other than those based on the E1-gene containing 293 cell line as described in Example 1C above, plasmid vectors containing E1 alone or in various combinations with E4 and fiber genes are constructed as described below.

The region of the adenovirus genome containing the E1a and E1b gene is amplified from viral genomic DNA by PCR as previously described. The primers used are E1L, the 5' or forward primer, and E1R, the 3' or backward primer, having the respective nucleotide sequences 5'CCGAGCTAGCGACTGAAAATGAG3' (SEQ ID NO 10) and 5'CCTCTCGAGAGACAGCAAGACAC3' (SEQ ID NO 11). The E1L and E1R primers include the respective restriction sites NheI and XhoI as indicated by the underlines. The sites are used to clone the amplified E1 gene fragment into the NheI/XhoI sites in pMAM commercially available from Clontech (Palo Alto, CA) to form the plasmid pDEX/E1 having 11152 bp, the plasmid map of which is shown in Figure 7.

The complete nucleotide sequence of pDEX/E1 is listed in SEQ ID NO 12 where the nucleotide position 1 corresponds to approximately 1454 nucleotides from the 3' end of the pMAM backbone vector sequence. The pDEX/E1 plasmid includes nucleotides 552 to 4090 of the adenovirus genome positioned downstream (beginning at nucleotide position 1460 and ending at 4998 in the pDEX/E1 plasmid) of the glucocorticoid-inducible mouse mammary tumor virus (MMTV) promoter of pMAM. The pMAM vector contains the *E. coli gpt* gene that allows stable transfectants to be isolated using hypoxanthine/aminopterin/thymidine (HAT) selection. The pMAM backbone occupies nucleotide positions 1-1454 and 5005-11152 of SEQ ID NO 12.

E. Generation of an Adenovirus Packaging Cell Line Carrying Plasmids Encoding Functional E1, and Fiber Proteins

To create separate adenovirus packaging cell lines equivalent to that of the 211 sublines, 211A, 211B and 211R, as described in Example 1C, alternative cell lines lacking adenoviral genomes are selected for transfection with the plasmid constructs as described below. Acceptable host cells include A549, Hela, Vero and the like cell lines as described in Example 1. The selected cell line is transfected with the separate plasmids, pDEX/E1 and pCLF, respectively for expressing E1, and fiber complementary proteins. Following transfection procedures as previously described, clones containing stable insertions of the two plasmids are isolated by selection with neomycin and HAT. Integration of full-length copy of the E1 gene is assessed by PCR amplification from genomic DNA using the primer set E1L/E1R, as described above. Functional insertion of the fiber gene is assayed by staining with the anti-fiber antibody as previously described.

The resultant stably integrated cell line is then used as a packaging cell system to complement adenoviral gene delivery vectors having the corresponding adenoviral gene deletions as described in Example 2.

F. Preparation of a Plasmid Containing Two or More Adenoviral Genes for Complementing Gene-Deleted Adenoviruses

The methods described in the preceding Examples rely on the use of two plasmids, pE4/Hygro and pCLF, or, pCLF and pDEX/E1 for generating adenoviral cell packaging systems. In alternative embodiments contemplated for use with the methods of this invention, complementing plasmids containing two or more adenoviral genes for expressing of encoded

proteins in various combinations are also prepared as described below. The resultant plasmids are then used in various cell systems with delivery plasmids having the corresponding adenoviral gene deletions. The selection of packaging cell, content of the delivery plasmids and content of the complementing plasmids for use in generating recombinant adenovirus viral vectors of this invention thus depends on whether other adenoviral genes are deleted along with the adenoviral fiber gene, and, if so, which ones.

1. Preparation of a Complementing Plasmid Containing Fiber and E1 Adenoviral Genes

A DNA fragment containing sequences for the CMV promoter, adenovirus tripartite leader, fiber gene and bovine growth hormone terminator is amplified from pCLF prepared in Example 1B using the forward primer 5'GACGGATCGGGAGATCTCC3' (SEQ ID NO 13), that anneals to the nucleotides 1-19 of the pCDNA3 vector backbone in pCLF, and the backward primer 5'CCGCCTCAGAAGCCATAGAGCC3' (SEQ ID NO 14) that anneals to nucleotides 1278-1257 of the pCDNA3 vector backbone. The fragment is amplified as previously described and then cloned into the pDEX/E1 plasmid, prepared in Example 1D. For cloning in the DNA fragment, the pDEX/E1 vector is first digested with NdeI, that cuts at a unique site in the pMAM vector backbone in pDEX/E1, then the ends are repaired by treatment with bacteriophage T4 polymerase and dNTPs.

The resulting plasmid containing E1 and fiber genes, designated pE1/Fiber, provides both dexamethasone-inducible E1 function as described for DEX/E1 and expression of Ad5 fiber protein as described above. A schematic plasmid map of pE1/Fiber, having 14455 bp, is shown in Figure 8.

The complete nucleotide sequence of pE1/Fiber is listed in SEQ ID NO 15 where the nucleotide position 1 corresponds to approximately to 1459 nucleotides from the 3' end of the parent vector pMAM sequence. The 5' and 3 ends of the Ad5 E1 gene are located at respective nucleotide positions 1460 and 4998 followed by pMAM backbone and then separated from the Ad5 fiber from pCLF by the filled-in blunt ended NdeI site. The 5' and 3' ends of the pCLF fiber gene fragment are located at respective nucleotide positions 10922-14223 containing elements as previously described for pCLF.

The resultant pE1/Fiber plasmid is then used to complement one or more delivery plasmids expressing E1 and fiber.

The pE1/Fiber construct is then used to transfect a selected host cell as described in Example 1E to generate stable chromosomal insertions performed as previously described followed by selection on HAT medium. The stable cells are then used as packaging cells as described in Example 2.

2) Preparation of a Complementing Plasmid Containing E4 and Fiber Adenoviral Genes

pCLF prepared as described in Example 1B is partially digested with BglII to cut only at the site in the pCDNA3 backbone. The pE4/Hygro plasmid prepared in Example 1A is digested with BamHI to produce a fragment containing E4. The E4 fragment is then inserted into the BamHI site of pCLF to form plasmid pE4/Fiber. The resultant plasmid provides expression of the fiber gene as described for pCLF and E4 function as described for pE4/Hygro.

A schematic plasmid map of pE4/Fiber, having 10610 bp, is shown in Figure 9. The complete nucleotide sequence of pE4/Fiber is listed in SEQ ID NO 16 where the nucleotide position 1 corresponds to approximately 14 bp from the 3' end of the parent vector pCDNA3 backbone sequence. The 5' and 3' ends of the Ad5 E4 gene are located at respective nucleotide positions 21 and 3149 followed by fused BglII/BamHI sites and pCDNA3 backbone including the CMV promoter again followed by BglII/BamHI sites. The adenovirus leader sequence begins at nucleotide position 4051 and extends to 4366 followed by fused BamHI/BglII sites and the 5' and 3' ends of the fiber gene located at respective nucleotide positions 4372 and 6124.

Stable chromosomal insertions of pE4/Fiber in host cells are obtained as described above.

## Example 2

### Preparation of Adenoviral Gene Delivery Vectors

#### Using Adenoviral Packaging Cell Lines

Adenoviral delivery vectors of this invention are prepared to separately lack the combinations of E1/fiber and E4/fiber. Such vectors are more replication-defective than those previously in use due to the absence of multiple viral genes. A preferred adenoviral delivery vector of this invention that is replication competent but only via a non-fiber means is one that only lacks the fiber gene but contains the remaining functional adenoviral regulatory and structural genes. Furthermore, the adenovirus delivery vectors of this invention have a higher capacity for insertion of foreign DNA.

#### A. Preparation of Adenoviral Gene Delivery Vectors Having Specific Gene Deletions and Methods of Use

To construct the E1/ fiber deleted viral vector containing the LacZ reporter gene construct, two new plasmids were constructed. The plasmid p $\Delta$  E1B $\beta$  gal was constructed as follows. By digestion of pSV $\beta$  gal (ProMega Corp., Madison, WI) with VspI, a DNA fragment containing the SV40 regulatory sequences and the *E. coli* - $\beta$ -galactosidase gene was isolated. The resulting fragment having overhanging ends was then filled in with Klenow fragment of DNA polymerase I in the presence of dNTPs followed by digestion with BamHI. The resulting fragment was cloned into the EcoRV and BamHI sites in the polylinker of p $\Delta$  E1sp1B (Microbix Biosystems, Hamilton, Ontario) to form p $\Delta$  E1B gal that therefore contained the left end of the adenovirus genome with the Ela region replaced by the LacZ cassette (nucleotides 6690 to 4151) of pSV $\beta$  gal. Plasmid DNA was prepared by the alkaline lysis method as described by Birnboim and Doly, Nuc. Acids Res., 7:1513-1523 (1978) from transformed cells used to expand the plasmid. DNA was then purified by CsCl-ethidium bromide density gradient centrifugation.

The second plasmid (pDV44), prepared as described herein, is derived from pBHG10, a vector prepared as described by Bett et al., Proc. Natl. Acad. Sci., USA, 91:8802-8806 (1994) and commercially available from Microbix, which contains an Ad5 genome with the packaging signals at the left end deleted and the E3 region (nucleotides 28133:30818) replaced



by a linker with a unique site for the restriction enzyme *PacI*. An 11.3 kb *BamHI* fragment, which contains the right end of the adenovirus genome, is isolated from pBHG10 and cloned into the *BamHI* site of pBS/SK(+) to create plasmid p11.3 having approximately 14,658 bp. A schematic of the plasmid map is shown in Figure 13. The p11.3 plasmid was then digested with *PacI* and *Sall* to remove the fiber, E4, and inverted terminal repeat (ITR) sequences.

This fragment is replaced with a fragment containing the ITR segments and the E4 gene which is generated by PCR amplification from pBHG10 using the following oligonucleotide sequences (SEQ ID NO 17) (SEQ ID NO 18). These primers incorporate sites for *PacI* and *BamHI*, respectively. Cloning this fragment into the *PacI* and blunt ended *Sall* sites of the p11.3 backbone resulted in a substitution of the fused ITRs, E4 region and fiber gene present in pBHG10, by the ITRs and E4 region alone.

In general, the method for virus production by recombination of plasmids followed by complementation in cell culture involves the isolation of recombinant viruses by cotransfection of any one of the adenovirus packaging cell systems prepared in Example 1, namely 211A, 211B, 211R, A549, Vero cells, and the like, with plasmids carrying sequences corresponding to viral gene delivery vectors.

A selected cell line is plated in dishes and cotransfected with pDV44 and pE1B gal using the calcium phosphate method as described by Bett et al., Proc. Natl. Acad. Sci., USA, 91:8802-8806 (1994). Recombination between the overlapping adenovirus sequences in the two plasmids leads to the creation of a full-length viral chromosome where pDV44 and p $\Delta$ E1B $\beta$  gal recombine to form a recombinant adenovirus vector having multiple deletions. The deletion of E1 and of the fiber gene from the viral chromosome is compensated for by the sequences integrated into the packaging cell genome, and infectious virus particles are produced. The plaques thus generated are isolated and stocks of the recombinant virus are produced by standard methods.

In a preferred embodiment of this invention, a delivery plasmid is prepared that does not require the above-described recombination events to prepare a therapeutic viral vector having a fiber gene deletion. A single delivery plasmid containing all the adenoviral genome

necessary for packaging but lacking the fiber gene is prepared from plasmid pFG140 containing full-length Ad85 that is commercially available from Microbix. The resultant delivery plasmid referred to as pFG140-f is then used with pCLF stably integrated cells as described above to prepare a therapeutic viral vector lacking fiber. In a preferred aspect of this invention, the fiber gene is replaced with a therapeutic gene of interest for preparing a therapeutic delivery adenoviral vector.

Vectors for the delivery of any desired therapeutic gene are prepared by cloning the gene of interest into the multiple cloning sites in the polylinker of commercially available pE1sp1B (Microbix Biosystems), in an analogous manner as performed for preparing pE1B gal as described above. The same cotransfection and recombination procedure is then followed as described herein to obtain viral gene delivery vectors.

The recombinant viruses thus produced are used as gene delivery tools both in cultured cells and *in vivo*. For studies of the effectiveness and relative immunogenicity of multiply-deleted vectors, virus particles are produced by growth in the packaging lines described in Example 1 and are purified by CsCl gradient centrifugation. Following titering, virus particles are administered to mice via systemic or local injection or by aerosol delivery to lung. The LacZ reporter gene allows the number and type of cells which are successfully transduced to be evaluated. The duration of transgene expression is evaluated in order to determine the long-term effectiveness of treatment with multiply-deleted recombinant adenoviruses relative to the standard technologies which have been used in clinical trials to date. The immune response to the improved vectors described here is determined by assessing parameters such as inflammation, production of cytotoxic T lymphocytes directed against the vector, and the nature and magnitude of the antibody response directed against viral proteins.

Versions of the vectors which contain therapeutic genes such as CFTR for treatment of cystic fibrosis or tumor suppressor genes for cancer treatment are evaluated in the animal system for safety and efficiency of gene transfer and expression. Following this evaluation, they are used as experimental therapeutic agents in human clinical trials.

**B. Retargeting of Adenoviral Gene Delivery Vectors by Producing Viral Particles Containing Different or Altered Fiber Proteins**

As the specificity of adenovirus binding to target cells is largely determined by the fiber protein, viral particles that incorporate modified fiber proteins or fiber proteins from different

adenoviral serotypes (pseudotyped vectors) have different specificities. Thus, the expression of the native Ad5 fiber protein in adenovirus packaging cells as described above is also applicable to production of different fiber proteins.

In one aspect of invention, chimeric fiber proteins are produced according to the methods of Stevenson et al., *J. Virol.*, 69:2850-2857 (1995). The authors showed that the determinants for fiber receptor binding activity are located in the head domain of the fiber and that isolated head domain is capable of trimerization and binding to cellular receptors. The head domains of adenovirus type 3 (Ad3) and Ad5 were exchanged in order to produce chimeric fiber proteins. Similar constructs for encoding chimeric fiber proteins for use in the methods of this invention are contemplated. Thus, instead of the using the intact Ad5 fiber-encoding construct prepared in Example 1 as a complementing viral vector in adenoviral packaging cells, the constructs described herein are used to transfect cells along with E4 and/or E1-encoding constructs.

Briefly, full-length Ad5 and Ad3 were amplified from purified adenovirus genomic DNA as a template. The Ad5 and Ad3 nucleotides sequences are available with the respective GenBank Accession Numbers M18369 and M12411. Oligonucleotide primers are designed to amplify the entire coding sequence of the full-length fiber genes, starting from the start codon, ATG, and ending with the termination codon TAA. For cloning purposes, the 5' and 3' primers contain the respective restriction sites BamHI and NotI for cloning into pcDNA plasmid as described in Example 1A. PCR is performed as described above.

The resultant products are then used to construct chimeric fiber constructs by PCR gene overlap extension, as described by Horton et al., *BioTechniques*, 8:525-535 (1990). The Ad5 fiber tail and shaft regions (5TS; the nucleotide region encoding amino acid residue positions 1 to 403) are connected to the Ad3 fiber head region (3H; the nucleotide region encoding amino acid residue positions 136 to 319) to form the 5TS3H fiber chimera. Conversely, the Ad3 fiber tail and shaft regions (3TS; the nucleotide region encoding amino acid residues positions 1 to 135) are connected to the Ad5 fiber head region (5H; the nucleotide region encoding the amino acid residue positions 404 to 581) to form the 3TS5H fiber chimera. The fusions are made at the conserved TLWT (SEQ ID NO 21) sequence at the fiber shaft-head junction.

The resultant chimeric fiber PCR products are then digested with BamHI and NotI for separate directional ligation into a similarly digested pcDNA vector. The Ad2 leader sequence is then subcloned into the BamHI as described in Example 1A for preparing an expression vector for subsequent transfection into 211 cells as described above or into the alternative packaging cell systems as previously described. The resultant chimeric fiber construct-containing adenoviral packaging cell lines are then used to complement adenoviral delivery vectors as previously described. Other fiber chimeric constructs are obtained using a similar approach with the various adenovirus serotypes known.

In an alternative embodiment, the methods of this invention contemplate the use of the modified proteins including novel epitopes as described by Michael et al., Gene Therapy, 2:660-668 (1995) and in International Publication WO 95/26412, the disclosures of which are incorporated by reference herein. Both publications describe the construction of a cell-type specific therapeutic viral vector having a new binding specificity incorporated into the virus concurrent with the destruction of the endogenous viral binding specificity. In particular, the authors described the production of an adenoviral vector encoding a gastrin releasing peptide (GRP) at the 3' end of the coding sequence of the Ad5 fiber gene. The resulting fiber-GRP fusion protein was expressed and shown to assemble functional fiber trimers that were correctly transported to the nucleus of HeLa cells following synthesis.

Based on the teachings in the paper and International Publication, similar constructs are contemplated for use in the complementing adenoviral packaging cell systems of this invention for generating new adenoviral gene delivery vectors that are replication-deficient and less immunogenic. Heterologous ligands contemplated for use herein to redirect fiber specificity range from as few as 10 amino acids in size to large globular structures, some of which necessitate the addition of a spacer region so as to reduce or preclude steric hindrance of the heterologous ligand with the fiber or prevent trimerization of the fiber protein. The ligands are inserted at the end or within the linker region. Preferred ligands include those that target specific cell receptors or those that are used for coupling to other moieties such as biotin and avidin. The types of cell signaling as a result of binding by a ligand is dependent upon the specificity of that ligand; i.e. receptor internalization or lack thereof.

A preferred spacer includes a short 12 amino acid peptide linker composed of a series of serines and alanine flanked by a proline residue at each end. One of ordinary skill in the art

is familiar with the preparation of linkers to accomplish sufficient protein presentation and for altering the binding specificity of the fiber protein without compromising the cellular events that follow viral internalization. Moreover, within the context of this invention, preparation of modified fibers having ligands positioned internally within the fiber protein, at the amino terminus and at the carboxy terminus as described below are contemplated for use with the methods described herein.

The preparation of a fiber having a heterologous binding ligand is prepared essentially as described in the above-cited paper. Briefly, for the ligand of choice, site-directed mutagenesis is used to insert the coding sequence for a linker into the NotI site at the 3' end of the Ad5 fiber construct in pCLF as prepared in Example 1. The 3' or antisense oligonucleotide encodes a preferred linker sequence of ProSerAlaSerAlaSerAlaSerAlaProGlySer (SEQ ID NO 22) followed by a unique restriction site and two stop codons, respectively, to allow the insertion of a coding sequence for a selected heterologous ligand and to ensure proper translation termination. The 3' end of the antisense oligonucleotide includes sequences that overlap with vector sequence into which the oligonucleotide is inserted via site-directed mutagenesis. Following mutagenesis of the pCLF sequence adding the linker and stop codon sequences, a nucleotide sequence encoding a preselected ligand is obtained, linkers corresponding to the unique restriction site are attached and then the sequence is cloned into linearized corresponding restriction site.

Into the resultant pCLF vector containing a Ad5 fiber gene sequence with 3' nucleotides encoding a linker and a ligand, the Ad2 leader sequence is inserted as previously described. The resultant fiber-ligand construct is then used to transfect 211 or the alternative cell packaging systems previously described to produce complementing viral vector packaging systems for use with the methods of this invention.

In a further embodiment, fiber proteins encoded by fiber genes isolated from different adenoviral serotypes are used intact for transfection into 211 or an alternative cell packaging system as previously described.

A gene encoding the fiber protein of interest is first cloned to create a plasmid analogous to pCLF, and stable cell lines producing the fiber protein are generated as described above for Ad5 fiber. The adenovirus vector described which lacks the fiber gene is then propagated in the cell line producing the fiber protein relevant for the purpose at hand. As the

only fiber gene present is the one in the packaging cells, the adenoviruses produced contain only the fiber protein of interest and therefore have the binding specificity conferred by the complementing protein. Such viral particles are used in studies such as those described above to determine their properties in experimental animal systems.

C. Targeted Gene Delivery Using Viral Vector Particles Lacking Fiber Protein

An alternative mode of entry for adenoviral infection of hematopoietic cells has been described by Huang, et al., J. Virol., 69:2257-2263 (1995) which does not involve the fiber protein-host cell receptor interaction. As infection of most other cell types does require the presence of fiber protein, vector particles which lack fiber may preferentially infect hematopoietic cells, such as monocytes or macrophages.

To produce a fiber-free adenovirus vector particle, a vector lacking the fiber gene as described above in Example 2A but containing a gene of interest for delivery is amplified by growth in cells which do not produce a fiber protein, such as the 211 cells prepared in Example 1, thereby producing large numbers of particles lacking fiber protein. The recovered fiber-free viral particles are then used to deliver the inserted gene of interest following the methods of this invention via targeting mechanisms provided by other regions of the adenoviral vector, i.e., via the native penton base.

**Example 3**

Deposit of Materials

The following cell lines and plasmids have been deposited on September 25, 1996, with the American Type Culture Collection, 1301 Parklawn Drive, Rockville, MD, USA (ATCC):

<u>Material</u>	<u>ATCC Accession No.</u>
Plasmid pE4/Hygro	97739
Plasmid pCLF	97737
211 Cell Line	CRL-12193
211A Cell Line	CRL-12194

The foregoing written specification is considered to be sufficient to enable one skilled in the art to practice the invention. The present invention is not to be limited in scope by the

cell lines and plasmids deposited, since the deposited embodiment is intended as a single illustration of one aspect of the invention and any cell lines or plasmid vectors that are functionally equivalent are within the scope of this invention.

The foregoing specification, including the specific embodiments and examples, is intended to be illustrative of the present invention and is not to be taken as limiting. Numerous other variations and modifications can be effected without departing from the true spirit and scope of the present invention.

#### Example 4

The native fiber protein is a homotrimer { Henry L.J. et al 1994 Characterization of the knob domain of the adenivirus Type 5 fiber protein expressed in *Escherichia coli* J. Virol 68:5239-5246 }, and trimerization is essential for assembly of the penton/fiber complex { Novelli A et al 1991 Assembly of adenovirus type 2 fiber synthesized in cell-free translation system. J. Biol. Chem 266:9299-9303 }. To assess the multimeric structure of the recombinant fiber protein produced by the cell lines, cells were labeled with 50  $\mu\text{Ci/ml}$  [ $^{35}\text{S}$ ] Translabel (ICN) for two hours at 37 ° C, lysed in RIPA buffer, and fiber protein was immunoprecipitated as described { Harlow E et al 1988 Antibodies. Cold Spring Harbour Laboratory, cold Spring Harbour}. Immune complexes were collected on Protein A-Sepharose beads (Pierce), extensively washed with RIPA buffer, and incubated at room temperature in 0.1 M triethylamine, pH 11.5 to release bound fiber protein. A portion of the precipitated fiber was electrophoresed on a 8% SDS-PAGE gel under denaturing (1% SDS in loading buffer, samples boiled for 5 minutes) or semi-native (0.1% SDS in loading buffer, samples not heated) conditions.

As seen in Fig. 13, lines 211A, 211B, and 211R, but not the control 293 cells, expressed an immunologically reactive protein which migrated at the predicted molecular weight for trimer (186 kD) under seminative conditions and for monomer (62 kD) under denaturing conditions. The behavior of the precipitated fiber was indistinguishable from that of purified baculovirus-produced recombinant Ad2

fiber { Wickham T et al 1993 Cell 73:309-319} (the 58 kD Ad2 and 62 kD Ad5 fibers have very similar mobilities under these conditions).

To determine whether the fiber-expressing lines could support the growth of a fiber-defective adenovirus, we performed one-step growth experiments using the temperature-sensitive fiber mutant Ad H5ts142 (the gift of Harold Ginsberg). At the restrictive temperature (39.5 °C), this mutant produces an underglycosylated fiber protein which is not incorporated into mature virions { Chee-Sheung C. C et al 1982 J. Virol 42: 932-950 }. This results in the accumulation of non-infectious viral particles. We asked whether the recombinant fiber protein expressed by our cell lines could complement the H5ts142 defect and rescue viral growth.

Cell lines 293, 211A, 211B and 211R (2 x 10<sup>6</sup> cells/sample) were infected with H5ts142 at 10 pfu/cell. 48 hours later, cells were detached with 25 mM EDTA and virus was harvested by four rapid freeze-thaw cycles. Debris was removed by a 10 minute spin at 1500 x g, and viral titers determined by fluorescent focus assay { Thiel J.F et al 1967 Proc. Soc. Exp. Biol. Med. 125:892-895 } on SW480 cells with a polyclonal anti-penton base Ab { Wickham T et al 1993 Cell 73:309-319}. As shown in Fig. 14, the fiber mutant virus replicated to high titers in 293 cells at 32.5° C (the permissive temperature), but to a much lower extent at the restrictive temperature of 39.5° C. The fiber-producing packaging lines 211A, 211B, or 211R supported virus production at 39.5° C to levels within two- to three-fold of those seen at the permissive temperature in 293 cells, indicating that these cells provided partial complementation of the fiber defect.

Interestingly, virus yields from the fiber-producing cell lines were also somewhat higher than those from 293 cells at 32.5° C (the 'permissive' temperature). This suggests that fiber produced by the ts142 virus may be partially defective even at the permissive temperature. Alternatively, a non-specific increase in adenoviral titer could result when viruses are grown in our packaging cells, by a mechanism not involving fiber complementation. However, we have found that viruses with wild type fiber genes (such as Ad.RSVbgal) replicate to identical levels either in our packaging lines or in 293



cells (data not shown). Taken together, these results demonstrate that the observed increase in H5ts142 growth is due to specific complementation of the fiber mutation.

Even in the fiber-expressing cell lines, the fiber mutant grows to higher titers at 32 °C than at 39.5 °C. This incomplete complementation may be due to the packaging lines' expression of fiber at a level somewhat below that seen in a wild-type infection (Fig. 16). A recent study reported an E4-deleted vector which coincidentally reduced fiber protein expression, resulting in a large reduction in the titer of virus produced { Brough et al 1996 J. Virol. 70: 6497-6501 }. Another possibility is that the defective ts142 fiber protein produced at the restrictive temperature might form complexes with some of the wild type protein produced by the cells and prevent its assembly into particles.

Although the fiber proteins of different Ad serotypes differ in the length of their shaft domains and in their receptor-binding knob domains, the N-terminal regions responsible for interaction with the viral penton base are highly conserved { Amberg N et al 1997 Virology 227:239-244 } (Fig. 15A). This suggests that fibers from many viral serotypes, with their different cell-binding specificities, may be amenable for use in producing gene delivery vectors.

We asked whether the recombinant Ad5 fiber produced by our packaging cells could be incorporated into particles of another adenovirus serotype. Adenovirus type 3 was grown either in fiber-producing cell lines or in 293 cells. Viral particles were purified by two sequential centrifugations (3 h at 111,000 x g) on preformed 15-40% CsCl gradients to remove soluble cellular proteins and then dialyzed extensively against 10 mM Tris-HCl, pH 8.1, 150 mM NaCl, 10% glycerol. Ad5 fiber protein was detected by immunoblotting using the polyclonal anti-fiber serum, followed by detection with a horseradish peroxidase-conjugated goat anti-rabbit antibody (Kirkegaard and Perry Laboratories) and the ECL chemiluminescence substrate (Amersham). The purified Ad3 particles contained Ad5 fiber protein after a single passage through a fiber-expressing cell line but not after passage through 293 cells (Fig. 15B). Previous work has demonstrated that Ad2 fiber is capable of interacting *in vitro* with Ad3 penton base { Fender et al 1997 Nature Biotech. 15:52-56 }, and our result

demonstrates that the type 5 fiber protein produced by the cells is capable of assembling into complete Ad3 particles.

A vector based on Ad5 but containing the gene for the Ad7 fiber protein has been described { Gall J. et al 1996 J. Virol. 70:2116-2123 }, as well as Ads containing chimeric fiber genes { Krasnykh V. N et al J. Virol. 70:6839-6846 }. Addition of a short peptide linker to the fiber in order to confer binding to a different cellular protein has also been reported { 8188 }. By using packaging technology such as that presented here, Ad vectors equipped with different fiber proteins may be produced simply by growth in cells expressing the fiber of interest, without the time-consuming step of generating a new vector genome for each application.

Replacing or modifying the fiber gene in the vector chromosome would also require that the new fiber protein bind a receptor on the surface of the cells it is to be grown. The packaging cell approach will allow the generation of Ad particles containing a fiber which can no longer bind to its host cells, by a single round of growth in cells expressing the desired fiber gene. This will greatly expand the repertoire of fiber proteins which can be incorporated into particles, as well as simplifying the process of retargeting gene delivery vectors.

Finally, a novel fiber-independent pathway of infection has recently been described in hematopoietic cells, in which penton base provides the initial virus-cell interaction by binding to integrin  $\alpha_m\beta_2$  { Huang S. et al 1996 J. Virol 70: 4502-4508 }. This suggests that viral particles lacking fiber protein may be useful in targeting gene delivery to specific cell types via this pathway.

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANT: Nemerow, Glen R.  
Von Seggern, Daniel J.
- (ii) TITLE OF INVENTION: PACKAGING CELL LINES, ADENOVIRUS  
VECTORS, AND METHODS OF USING SAME
- (iii) NUMBER OF SEQUENCES: 20
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: THE SCRIPPS RESEARCH INSTITUTE
  - (B) STREET: 10550 North Torrey Pines Road
  - (C) CITY: La Jolla
  - (D) STATE: California
  - (E) COUNTRY: United States
  - (F) ZIP: 92037
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Floppy disk
  - (B) COMPUTER: IBM PC compatible
  - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
  - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER: US
  - (B) FILING DATE: 25-SEP-1996
  - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: Logan, April C.
  - (B) REGISTRATION NUMBER: 33,950
  - (C) REFERENCE/DOCKET NUMBER: TSRI 554.0
- (ix) TELECOMMUNICATION INFORMATION:
  - (A) TELEPHONE: (619) 554-2937

- 61 -

(B) TELEFAX: (619) 554-6312

## (2) INFORMATION FOR SEQ ID NO:1:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 30 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CGGTACACAG AATTCAGGAG ACACAACTCC

30

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 35 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

GCCTGGATCC GGGAAGTTAC GTAACGTGGG AAAAC  
35

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CGCGGATCCG CG  
12

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 8710 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: circular

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CACCTAAATT GTAAGCGTTA ATATTTTGTT AAAATTCGCG TTAAATTTTT GTTAAATCAG  
60

CTCATTTTTT AACCAATAGG CCGAAATCGG CAAAATCCCT TATAAATCAA AAGAATAGAC  
120

CGAGATAGGG TTGAGTGTTG TTCCAGTTTG GAACAAGAGT CCACTATTAA AGAACGTGGA  
180

CTCCAACGTC AAAGGGCGAA AAACCGTCTA TCAGGGCGAT GGCCCCACTAC GTGAACCATC  
240

ACCCTAATCA AGTTTTTTGG GGTCGAGGTG CCGTAAAGCA CTAAATCGGA ACCCTAAAGG  
300

GAGCCCCCGA TTTAGAGCTT GACGGGGAAA GCCGGCGAAC GTGGCGAGAA AGGAAGGGAA  
360

GAAAGCGAAA GGAGCGGGCG CTAGGGCGCT GGCAAGTGTA GCGGTCACGC TGCGCGTAAC  
420

CACCACACCC GCCGCGCTTA ATGCGCCGCT ACAGGGCGCG TCCCATTTCG CATTTCAGGCT  
480

GCGCAACTGT TGGGAAGGGC GATCGGTGCG GGCCTCTTCG CTATTACGCC AGCTGGCGAA  
540

AGGGGGATGT GCTGCAAGGC GATTAAGTTG GGTAACGCCA GGGTTTTCCC AGTCACGACG  
600

TTGTAAAACG ACGGCCAGTG AATTGTAATA CGACTCACTA TAGGGCGAAT TGGGTACCGG  
660

GCCCCCCTC GAGGTCGACG GTATCGATAA GCTTGATATC GAATTCAGGA GACACAACCTC  
720

CAAGTGCATA CTCTATGTCA TTTTCATGGG ACTGGTCTGG CCACAACCTAC ATTAATGAAA  
780

TATTTGCCAC ATCCTCTTAC ACTTTTTCAT ACATTGCCCA AGAATAAAGA ATCGTTTGTG  
840

TTATGTTTCA ACGTGTTTAT TTTTCAATTG CAGAAAATTT CAAGTCATTT TTCATTTCAGT  
900

AGTATAGCCC CACCACCACA TAGCTTATAC AGATCACCGT ACCTTAATCA AACTCACAGA  
960

ACCCTAGTAT TCAACCTGCC ACCTCCCTCC CAACACACAG AGTACACAGT CCTTTCTCCC  
1020

CGGCTGGCCT TAAAAAGCAT CATATCATGG GTAACAGACA TATTCTTAGG TGTTATATTC  
1080

CACACGGTTT CCTGTGAGC CAAACGCTCA TCAGTGATAT TAATAAACTC CCCGGGCAGC  
1140

TCACTTAAGT TCATGTCGCT GTCCAGCTGC TGAGCCACAG GCTGCTGTCC AACTTGCGGT  
1200

TGCTTAACGG GCGGCGAAGG AGAAGTCCAC GCCTACATGG GGGTAGAGTC ATAATCGTGC  
1260

ATCAGGATAG GCGGGTGGTG CTGCAGCAGC GCGCGAATAA ACTGCTGCCG CCGCCGCTCC  
1320

GTCCTGCAGG AATACAACAT GGCAGTGGTC TCCTCAGCGA TGATTTCGCAC CGCCCCGAGC  
1380

ATAAGGCGCC TTGTCCTCCG GGCACAGCAG CGCACCTGA TCTCACTTAA ATCAGCACAG  
1440

TAAGTGCAGC ACAGCACCAC AATATTGTTC AAAATCCCAC AGTGCAAGGC GCTGTATCCA  
1500

AAGCTCATGG CGGGGACCAC AGAACCCACG TGGCCATCAT ACCACAAGCG CAGGTAGATT  
1560

AAGTGGCGAC CCCTCATAAA CACGCTGGAC ATAAACATTA CCTCTTTTGG CATGTTGTAA  
1620

TTCACCACCT CCCGGTACCA TATAAACCTC TGATTAAACA TGGCGCCATC CACCACCATC  
1680

CTAAACCAGC TGGCCAAAAC CTGCCC GCCG GCTATACACT GCAGGGAACC GGGACTGGAA  
1740

CAATGACAGT GGAGAGCCCA GGACTCGTAA CCATGGATCA TCATGCTCGT CATGATATCA  
1800

ATGTTGGCAC AACACAGGCA CACGTGCATA CACTTCCTCA GGATTACAAG CTCCTCCCGC  
1860

GTTAGAACCA TATCCCAGGG AACAACCCAT TCCTGAATCA GCGTAAATCC CACACTGCAG  
1920

GGAAGACCTC GCACGTAAC CACGTTGTGC ATTGTCAAAG TGTTACATTC GGGCAGCAGC  
1980

GGATGATCCT CCAGTATGGT AGCGCGGGTT TCTGTCTCAA AAGGAGGTAG ACGATCCCTA  
2040

CTGTACGGAG TGCGCCGAGA CAACCGAGAT CGTGTTGGTC GTAGTGTCAT GCCAAATGGA  
2100

ACGCCGGACG TAGTCATATT TCCTGAAGCA AAACCAGGTG CGGGCGTGAC AACAGATCT  
2160

GCGTCTCCGG TCTCGCCGCT TAGATCGCTC TGTGTAGTAG TTGTAGTATA TCCACTCTCT  
2220

CAAAGCATCC AGGCGCCCCC TGGCTTCGGG TTCTATGTAA ACTCCTTCAT GCGCCGCTGC  
2280



CCTGATAACA TCCACCACCG CAGAATAAGC CACACCCAGC CAACCTACAC ATTCTGTTCTG  
2340

CGAGTCACAC ACGGGAGGAG CGGGAAGAGC TGGAAGAACC ATGTTTTTTT TTTTATTCCA  
2400

AAAGATTATC CAAAACCTCA AAATGAAGAT CTATTAAGTG AACGCGCTCC CCTCCGGTGG  
2460

CGTGGTCAAA CTCTACAGCC AAAGAACAGA TAATGGCATT TGTAAGATGT TGCACAATGG  
2520

CTTCCAAAAG GCAAACGGCC CTCACGTCCA AGTGGACGTA AAGGCTAAAC CCTTCAGGGT  
2580

GAATCTCCTC TATAAACATT CCAGCACCTT CAACCATGCC CAAATAATTC TCATCTCGCC  
2640

ACCTTCTCAA TATATCTCTA AGCAAATCCC GAATATTAAG TCCGGCCATT GTAAAAATCT  
2700

GCTCCAGAGC GCCCTCCACC TTCAGCCTCA AGCAGCGAAT CATGATTGCA AAAATTCAGG  
2760

TTCCTCACAG ACCTGTATAA GATTCAAAAG CGGAACATTA ACAAAAATAC CGCGATCCCCG  
2820

TAGGTCCCTT CGCAGGGCCA GCTGAACATA ATCGTGCAGG TCTGCACGGA CCAGCGCGGC  
2880

CACTTCCCCG CCAGGAACCT TGACAAAAGA ACCCACACTG ATTATGACAC GCATACTCGG  
2940

AGCTATGCTA ACCAGCGTAG CCCCAGTGTA AGCTTTGTTG CATGGGCGGC GATATAAAAT  
3000

GCAAGGTGCT GCTCAAAAAA TCAGGCAAAG CCTCGCGCAA AAAAGAAAGC ACATCGTAGT  
3060

CATGCTCATG CAGATAAAGG CAGGTAAGCT CCGGAACCAC CACAGAAAAA GACACCATT  
3120

TTCTCTCAAA CATGTCTGCG GGTTTCTGCA TAAACACAAA ATAAAATAAC AAAAAACAT  
3180

TTAAACATTA GAAGCCTGTC TTACAACAGG AAAACAACC CTTATAAGCA TAAGACGGAC  
3240

TACGGCCATG CCGGCGTGAC CGTAAAAAAA CTGGTCACCG TGATTAAAAA GCACCACCGA  
3300

CAGCTCCTCG GTCATGTCCG GAGTCATAAT GTAAGACTCG GTAAACACAT CAGGTTGATT  
3360

CATCGGTCAG TGCTAAAAAG CGACCGAAAT AGCCCGGGGG AATACATACC CGCAGGCGTA  
3420

GAGACAACAT TACAGCCCCC ATAGGAGGTA TAACAAAATT AATAGGAGAG AAAAACACAT  
3480

AAACACCTGA AAAACCCTCC TGCCTAGGCA AAATAGCACC CTCCCGCTCC AGAACAACAT  
3540

ACAGCGCTTC ACAGCGGCAG CCTAACAGTC AGCCTTACCA GTAAAAAGA AAACCTATTA  
3600

AAAAACACC ACTCGACAG GCACCAGCTC AATCAGTCAC AGTGTAAGAA AGGGCCAAGT  
3660

GCAGAGCGAG TATATATAGG ACTAAAAAAT GACGTAACGG TTAAAGTCCA CAAAAACAC  
3720

CCAGAAAACC GCACGCGAAC CTACGCCAG AAACGAAAGC CAAAAACCC ACAACTTCCT  
3780

CAAATCGTCA CTTCCGTTTT CCCACGTTAC GTAACCTCCC GGATCCGCGG CATTACAGT  
3840

TCTCCGCAAG AATTGATTGG CTCCAATTCT TGGAGTGGTG AATCCGTTAG CGAGGTGCCC  
3900

CCGGCTTCCA TTCAGGTCGA GGTGGCCCGG CTCCATGCAC CGCGACGCAA CGCGGGGAGG  
3960

CAGACAAGGT ATAGGGCGGC GCCTACAATC CATGCCAACC CGTTCCATGT GCTCGCCGAG  
4020

GCGGCATAAA TCGCCGTGAC GATCAGCGGT CCAGTGATCG AAGTTAGGCT GGTAAGAGCC  
4080

GCGAGCGATC CTTGAAGCTG TCCCTGATGG TCGTCATCTA CCTGCCTGGA CAGCATGGCC  
4140

TGCAACGCGG GCATCCCGAT GCCGCCGGAA GCGAGAAGAA TCATAATGGG GAAGGCCATC  
4200

CAGCCTCGCG TCGCGAACGC CAGCAAGACG TAGCCCAGCG CGTCGGCCGC CATGCCCTGC  
4260

TTCATCCCCG TGGCCCGTTG CTCGCGTTTG CTGGCGGTGT CCCCGBAAGA AATATATTTG  
4320

CATGTCTTTA GTTCTATGAT GACACAAACC CCGCCCAGCG TCTTGTCATT GGCGAATTCC  
4380

AACACGCAGA TGCAGTCGGG GCGGCGCGGT CCCAGGTCCA CTTCGCATAT TAAGGTGACG  
4440

CGTGTGGCCT CGAACACCGA GCGACCCTGC AGCGACCCGC TTAACAGCGT CAACAGCGTG  
4500

CCGCAGATCC CGGGCAATGA GATATGAAAA AGCCTGAACT CACCGCGACG TCTGTGCGAG  
4560

AGTTTCTGAT CGAAAAGTTC GACAGCGTCT CCGACCTGAT GCAGCTCTCG GAGGGCGAAG  
4620

AATCTCGTGC TTTCAGCTTC GATGTAGGAG GCGTGGATA TGTCTGCGG GTAAATAGCT  
4680

GCGCCGATGG TTTCTACAAA GATCGTTATG TTTATCGGCA CTTTGCATCG GCCGCGCTCC  
4740

CGATTCCGGA AGTGCTTGAC ATTGGGGAAT TCAGCGAGAG CCTGACCTAT TGCATCTCCC  
4800

GCCGTGCACA GGGTGTACAG TTGCAAGACC TGCCTGAAAC CGAACTGCCC GCTGTTCTGC  
4860

AGCCGGTCGC GGAGGCCATG GATGCGATCG CTGCGGCCGA TCTTAGCCAG ACGAGCGGGT  
4920

TCGGCCCATT CGGACCGCAA GGAATCGGTC AATACACTAC ATGGCGTGAT TTCATATGCG  
4980

CGATTGCTGA TCCCCATGTG TATCACTGGC AAATGTGAT GGACGACACC GTCAGTGCGT  
5040

CCGTGCGCA GGCTCTCGAT GAGCTGATGC TTTGGGCCGA GGACTGCCCC GAAGTCCGGC  
5100

ACCTCGTGCA CGCGGATTTC GGCTCCAACA ATGTCCTGAC GGACAATGGC CGCATAACAG  
5160

CGGTCATTGA CTGGAGCGAG GCGATGTTGC GGGATTCCCA ATACGAGGTC GCCAACATCT  
5220

TCTTCTGGAG GCCGTGGTTG GCTTGTATGG AGCAGCAGAC GCGCTACTTC GAGCGGAGGC  
5280

ATCCGGAGCT TGCAGGATCG CCGCGGCTCC GGGCGTATAT GCTCCGCATT GGTCTTGACC  
5340

AACTCTATCA GAGCTTGGTT GACGGCAATT TCGATGATGC AGCTTGGGCG CAGGGTCGAT  
5400

GCGACGCAAT CGTCCGATCC GGAGCCGGGA CTGTCGGGCG TACACAAATC GCCCGCAGAA  
5460

GCGCGGCCGT CTGGACCGAT GGCTGTGTAG AAGTACTCGC CGATAGTGGA AACCGACGCC  
5520

CCAGCACTCG TCCGAGGGCA AAGGAATAGG GGAGATGGGG GAGGCTAACT GAAACACGGA  
5580

AGGAGACAAT ACCGGAAGGA ACCCGCGCTA TGACGGCAAT AAAAAGACAG AATAAAACGC  
5640

ACGGGTGTTG GGTCGTTTGT TCATAAACGC GGGGTTCGGT CCCAGGGCTG GCACTCTGTC  
5700

GATACCCAC CGAGACCCCA TTGGGGCCAA TACGCCGCG TTTCTTCCTT TTCCCCACCC  
5760

CACCCCCCAA GTTCGGGTGA AGGCCAGGG CTCGCAGCCA ACGTCGGGGC GGCAGGCCCT  
5820

GCCATAGCCA CTGGCCCCGT GGGTTAGGGA CGGGTCCCC CATGGGGAAT GGTTTATGGT  
5880

TCGTGGGGGT TATTATTTTG GCGTTGCGT GGGGTCTGGT CCACGACTGG ACTGAGCAGA  
5940

CAGACCCATG GTTTTGGAT GGCCTGGGCA TGGACCGCAT GTACTGGCGC GACACGAACA  
6000

CCGGGCGTCT GTGGCTGCCA AACACCCCCG ACCCCAAAA ACCACGCGC GGATTTCTGG  
6060

CGCCCAGTGC CGTCGACCGG TCATGGCTGC GCCCGACAC CCGCCAACAC CCGCTGACGC  
6120

GCCCTGACGG GCTTGTCTGC TCCCGGCATC CGCTTACAGA CAAGCTGTGA CCGTCTCCGG  
6180

GAGCTGCATG TGTCAGAGGT TTTCACCGTC ATCACCGAAA CGCGCGAGGC AGCCGGATCA  
6240

TAATCAGCCA TACCACATTT GTAGAGGTTT TACTTGCTTT AAAAAACCTC CCCACCTCCC  
6300

CCTGAACCTG AAACATAAAA TGAATGCAAT TGTGTGTGTT AACTTGTTTA TTGCAGCTTA  
6360

TAATGGTTAC AAATAAAGCA ATAGCATCAC AAATTCACA AATAAAGCAT TTTTTCCTACT  
6420

GCATTCTAGT TGTGGTTTGT CCAAACATCAT CAATGTATCT TATCATGTCT GGATCCACTA  
6480

GTTCTAGAGC GGCCGCCACC GCGGTGGAGC TCCAGCTTTT GTTCCCTTTA GTGAGGGTTA  
6540

ATTTCGAGCT TGGCGTAATC ATGGTCATAG CTGTTTCCTG TGTGAAATTG TTATCCGCTC  
6600

ACAATTCCAC ACAACATACG AGCCGGAAGC ATAAAGTGTA AAGCCTGGGG TGCCTAATGA  
6660

GTGAGCTAAC TCACATTAAT TGCGTTGCGC TCACTGCCCCG CTTTCCAGTC GGGAAACCTG  
6720

TCGTGCCAGC TGCATTAATG AATCGGCCAA CGCGCGGGGA GAGGCGGTTT GCGTATTGGG  
6780

CGCTCTTCCG CTTCCCTCGCT CACTGACTCG CTGCGCTCGG TCGTTCGGCT GCGGCGAGCG  
6840

GTATCAGCTC ACTCAAAGGC GGTAATACGG TTATCCACAG AATCAGGGGA TAACGCAGGA  
6900

AAGAACATGT GAGCAAAAGG CCAGCAAAAG GCCAGGAACC GTAAAAAGGC CGCGTTGCTG  
6960

GCGTTTTTCC ATAGGCTCCG CCCCCCTGAC GAGCATCACA AAAATCGACG CTCAAGTCAG  
7020

AGGTGGCGAA ACCCGACAGG ACTATAAAGA TACCAGGCGT TTCCCCCTGG AAGCTCCCTC  
7080

GTGCGCTCTC CTGTTCCGAC CCTGCCGCTT ACCGGATACC TGTCCGCCTT TCTCCCTTCG  
7140

GGAAGCGTGG CGCTTTCTCA TAGCTCACGC TGTAGGTATC TCAGTTCGGT GTAGGTCGTT  
7200

CGCTCCAAGC TGGGCTGTGT GCACGAACCC CCCGTTTCAGC CCGACCGCTG CGCCTTATCC  
7260

GGTAACTATC GTCTTGAGTC CAACCCGGTA AGACACGACT TATCGCCACT GGCAGCAGCC  
7320

ACTGGTAACA GGATTAGCAG AGCGAGGTAT GTAGGCGGTG CTACAGAGTT CTTGAAGTGG  
7380

TGGCCTAACT ACGGCTACAC TAGAAGGACA GTATTTGGTA TCTGCGCTCT GCTGAAGCCA  
7440

GTTACCTTCG GAAAAAGAGT TGGTAGCTCT TGATCCGGCA AACAAACCAC CGCTGGTAGC  
7500

GGTGGTTTTT TTGTTTGCAA GCAGCAGATT ACGCGCAGAA AAAAAGGATC TCAAGAAGAT  
7560

CCTTTGATCT TTTCTACGGG GTCTGACGCT CAGTGAACG AAAACTCACG TTAAGGGATT  
7620

TTGGTCATGA GATTATCAAA AAGGATCTTC ACCTAGATCC TTTTAAATTA AAAATGAAGT  
7680

TTTAAATCAA TCTAAAGTAT ATATGAGTAA ACTTGGTCTG ACAGTTACCA ATGCTTAATC  
7740

AGTGAGGCAC CTATCTCAGC GATCTGTCTA TTTCGTTTAT CCATAGTTGC CTGACTCCCC  
7800

GTCGTGTAGA TAACTACGAT ACGGGAGGGC TTACCATCTG GCCCCAGTGC TGCAATGATA  
7860

CCGCGAGACC CACGCTCACC GGCTCCAGAT TTATCAGCAA TAAACCAGCC AGCCGGAAGG  
7920

GCCGAGCGCA GAAGTGGTCC TGCAACTTTA TCCGCCTCCA TCCAGTCTAT TAATTGTTGC  
7980

CGGGAAGCTA GAGTAAGTAG TTCGCCAGTT AATAGTTTGC GCAACGTTGT TGCCATTGCT  
8040

ACAGGCATCG TGGTGTACAG CTCGTCGTTT GGTATGGCTT CATTCAGCTC CGGTTCCCAA  
8100

CGATCAAGGC GAGTTACATG ATCCCCCATG TTGTGCAAAA AAGCGGTTAG CTCCTTCGGT  
8160

CCTCCGATCG TTGTCAGAAG TAAGTTGGCC GCAGTGTTAT CACTCATGGT TATGGCAGCA  
8220

CTGCATAATT CTCTTACTGT CATGCCATCC GTAAGATGCT TTTCTGTGAC TGGTGAGTAC  
8280

TCAACCAAGT CATTCTGAGA ATAGTGTATG CGGCGACCGA GTTGCTCTTG CCCGGCGTCA  
8340

ATACGGGATA ATACCGCGCC ACATAGCAGA ACTTTAAAAG TGCTCATCAT TGGAAAACGT  
8400

TCTTCGGGGC GAAAACTCTC AAGGATCTTA CCGCTGTTGA GATCCAGTTC GATGTAACCC  
8460

ACTCGTGCAC CCAACTGATC TTCAGCATCT TTTACTTTCA CCAGCGTTTC TGGGTGAGCA  
8520



- 74 -

AAAACAGGAA GGCAAAATGC CGCAAAAAG GGAATAAGGG CGACACGGAA ATGTTGAATA  
8580

CTCATACTCT TCCTTTTTC AATTATTGA AGCATTTATC AGGGTTATTG TCTCATGAGC  
8640

GGATACATAT TTGAATGTAT TTAGAAAAAT AAACAAATAG GGGTTCCGCG CACATTTCCT  
8700

CGAAAAGTGC  
8710

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

ATGGGATCCA AGATGAAGCG CGCAAGACCG  
30

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single

- 75 -

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

CATAACGCGG CCGCTTCTTT ATTCTTGGGC

30

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 7148 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: circular

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

GACGGATCGG GAGATCTCCC GATCCCCTAT GGTGACTCT CAGTACAATC TGCTCTGATG

60

CCGCATAGTT AAGCCAGTAT CTGCTCCCTG CTTGTGTGTT GGAGGTCGCT GAGTAGTGCG

120

CGAGCAAAAT TTAAGCTACA ACAAGGCAAG GCTTGACCGA CAATTGCATG AAGAATCTGC

180

TTAGGGTTAG GCGTTTTGCG CTGCTTCGCG ATGTACGGGC CAGATATACG CGTTGACATT  
240

GATTATTGAC TAGTTATTAA TAGTAATCAA TTACGGGGTC ATTAGTTCAT AGCCCATATA  
300

TGGAGTTCCG CGTTACATAA CTTACGGTAA ATGGCCCGCC TGGCTGACCG CCCAACGACC  
360

CCCGCCCATT GACGTCAATA ATGACGTATG TTCCCATAGT AACGCCAATA GGGACTTTCC  
420

ATTGACGTCA ATGGGTGGAC TATTTACGGT AAACGTGCCA CTTGGCAGTA CATCAAGTGT  
480

ATCATATGCC AAGTACGCCC CCTATTGACG TCAATGACGG TAAATGGCCC GCCTGGCATT  
540

ATGCCCAGTA CATGACCTTA TGGGACTTTC CTACTTGGCA GTACATCTAC GTATTAGTCA  
600

TCGCTATTAC CATGGTGATG CGGTTTTGGC AGTACATCAA TGGGCGTGGA TAGCGGTTTG  
660

ACTCACGGGG ATTTCCAAGT CTCCACCCCA TTGACGTCAA TGGGAGTTTG TTTTGGCACC  
720

AAAATCAACG GGAAGTTTCCA AAATGTCGTA ACAACTCCGC CCCATTGACG CAAATGGGCG  
780

GTAGGCGTGT ACGGTGGGAG GTCTATATAA GCAGAGCTCT CTGGCTAACT AGAGAACCCA  
840

CTGCTTACTG GCTTATCGAA ATTAATACGA CTCACTATAG GGAGACCCAA GCTTGGTACC  
900

GAGCTCGGAT CCAAGATGAA GCGCGCAAGA CCGTCTGAAG ATACCTTCAA CCCCCTGTAT  
960

CCATATGACA CGGAAACCGG TCCTCCAACT GTGCCTTTTC TTA CTCTCC CTTTGTATCC  
1020

CCCAATGGGT TTCAAGAGAG TCCCCCTGGG GTACTCTCTT TGCGCCTATC CGAACCTCTA  
1080

GTTACCTCCA ATGGCATGCT TGCGCTCAAA ATGGGCAACG GCCTCTCTCT GGACGAGGCC  
1140

GGCAACCTTA CCTCCCCAAA TGTAACCACT GTGAGCCCAC CTCTCAAAAA AACCAAGTCA  
1200

AACATAAACC TGGAAATATC TGCACCCCTC ACAGTTACCT CAGAAGCCCT AACTGTGGCT  
1260

GCCGCCGCAC CTCTAATGGT CGCGGGCAAC AACTCACCAC TGCAATCACA GGCCCCGCTA  
1320

ACCGTGACAG ACTCCAAACT TAGCATTGCC ACCCAAGGAC CCCTCACAGT GTCAGAAGGA  
1380

AAGCTAGCCC TGCAAACATC AGGCCCCCTC ACCACCACCG ATAGCAGTAC CCTTACTATC  
1440

ACTGCCTCAC CCCCTCTAAC TACTGCCACT GGTAGCTTGG GCATTGACTT GAAAGAGCCC  
1500

ATTTATACAC AAAATGGAAA ACTAGGACTA AAGTACGGGG CTCCTTTGCA TGTAACAGAC  
1560

GACCTAAACA CTTTGACCGT AGCAACTGGT CCAGGTGTGA CTATTAATAA TACTTCCTTG  
1620

CAAATAAAG TTA CTGGAGC CTTGGGTTTT GATTACACAAG GCAATATGCA ACTTAATGTA  
1680

GCAGGAGGAC TAAGGATTGA TTCTCAAAAC AGACGCCTTA TACTTGATGT TAGTTATCCG  
1740

TTTGATGCTC AAAACCAACT AAATCTAAGA CTAGGACAGG GCCCTCTTTT TATAAACTCA  
1800

GGCCACAAC TGGATATTAA CTACAACAAA GGCCTTTACT TGTTTACAGC TTCAAACAAT  
1860

TCCAAAAAGC TTGAGGTTAA CCTAAGCACT GCCAAGGGGT TGATGTTTGA CGCTACAGCC  
1920

ATAGCCATTA ATGCAGGAGA TGGGCTTGAA TTTGGTTCAC CTAATGCACC AAACACAAAT  
1980

CCCCTCAAAA CAAAAATTGG CCATGGCCTA GAATTTGATT CAAACAAGGC TATGGTTCCT  
2040

AAACTAGGAA CTGGCCTTAG TTTTGACAGC ACAGGTGCCA TTACAGTAGG AAACAAAAAT  
2100

AATGATAAGC TAACTTTGTG GACCACACCA GCTCCATCTC CTAAGTGTAG ACTAAATGCA  
2160

GAGAAAGATG CTAACTCAC TTTGGTCTTA ACAAATGTG GCAGTCAAAT ACTTGCTACA  
2220

GTTTCAGTTT TGGCTGTAA AGGCAGTTG GCTCCAATAT CTGGAACAGT TCAAAGTGCT  
2280

CATCTTATTA TAAGATTTGA CGAAAATGGA GTGCTACTAA ACAATTCCTT CCTGGACCCA  
2340

GAATATTGGA ACTTTAGAAA TGGAGATCTT ACTGAAGGCA CAGCCTATAC AAACGCTGTT  
2400

GGATTTATGC CTAACCTATC AGCTTATCCA AAATCTCAG GTAAACTGC CAAAAGTAAC  
2460

ATTGTCAGTC AAGTTTACTT AAACGGAGAC AAAACTAAAC CTGTAACACT AACCATTACA  
2520

CTAAACGGTA CACAGGAAAC AGGAGACACA ACTCCAAGTG CATACTCTAT GTCATTTTCA  
2580

TGGGACTGGT CTGGCCACAA CTACATTAAT GAAATATTTG CCACATCCTC TTACACTTTT  
2640

TCATACATTG CCCAAGAATA AAGAAGCGGC CGCTCGAGCA TGCATCTAGA GGGCCCTATT  
2700

CTATAGTGTC ACCTAAATGC TAGAGCTCGC TGATCAGCCT CGACTGTGCC TTCTAGTTGC  
2760

CAGCCATCTG TTGTTTGCCC CTCCCCCGTG CCTTCCTTGA CCCTGGAAGG TGCCACTCCC  
2820

ACTGTCCTTT CCTAATAAAA TGAGGAAATT GCATCGCATT GTCTGAGTAG GTGTCATTCT  
2880

ATTCTGGGGG GTGGGGTGGG GCAGGACAGC AAGGGGGAGG ATTGGGAAGA CAATAGCAGG  
2940

CATGCTGGGG ATGCGGTGGG CTCTATGGCT TCTGAGGCGG AAAGAACCAG CTGGGGCTCT  
3000

AGGGGGTATC CCCACGCGCC CTGTAGCGGC GCATTAAGCG CGGCGGGTGT GGTGGTTACG  
3060

CGCAGCGTGA CCGCTACACT TGCCAGCGCC CTAGCGCCCG CTCCTTTCGC TTTCTTCCCT  
3120

TCCTTTCTCG CCACGTTTCGC CGGCTTTCCC CGTCAAGCTC TAAATCGGGG CATCCCTTTA  
3180

GGGTTCCGAT TTAGTGCTTT ACGGCACCTC GACCCCCAAA AACTTGATTA GGGTGATGGT  
3240

TCACGTAGTG GGCCATCGCC CTGATAGACG GTTTTTCGCC CTTTGACGTT GGAGTCCACG  
3300

TTCTTTAATA GTGGACTCTT GTTCCAAACT GGAACAACAC TCAACCCTAT CTCGGTCTAT  
3360

TCTTTTGATT TATAAGGGAT TTTGGGGATT TCGGCCTATT GGTAAAAAA TGAGCTGATT  
3420

TAACAAAAAT TTAACGCGAA TTAATTCTGT GGAATGTGTG TCAGTTAGGG TGTGGAAAGT  
3480

CCCCAGGCTC CCCAGGCAGG CAGAAGTATG CAAAGCATGC ATCTCAATTA GTCAGCAACC  
3540

AGGTGTGGAA AGTCCCCAGG CTCCCCAGCA GGCAGAAGTA TGCAAAGCAT GCATCTCAAT  
3600

TAGTCAGCAA CCATAGTCCC GCCCCTAACT CCGCCCCATCC CGCCCCTAAC TCCGCCCAGT  
3660

TCCGCCCATT CTCCGCCCCA TGGCTGACTA ATTTTTTTTA TTTATGCAGA GGCCGAGGCC  
3720

GCCTCTGCCT CTGAGCTATT CCAGAAGTAG TGAGGAGGCT TTTTGGAGG CCTAGGCTTT  
3780

TGCAAAAAGC TCCCGGGAGC TTGTATATCC ATTTTCGGAT CTGATCAAGA GACAGGATGA  
3840

GGATCGTTTC GCATGATTGA ACAAGATGGA TTGCACGCAG GTTCTCCGGC CGCTTGGGTG  
3900

GAGAGGCTAT TCGGCTATGA CTGGGCACAA CAGACAATCG GCTGCTCTGA TGCCGCCGTG  
3960

TTCCGGCTGT CAGCGCAGGG GCGCCCGGTT CTTTTGTCA AGACCGACCT GTCCGGTGCC  
4020

CTGAATGAAC TGCAGGACGA GGCAGCGCGG CTATCGTGCC TGGCCACGAC GGGCGTTCCT  
4080

TGCGCAGCTG TGCTCGACGT TGTCAGTAA GCGGGAAGGG ACTGGCTGCT ATTGGGCGAA  
4140

GTGCCGGGGC AGGATCTCCT GTCATCTCAC CTTGCTCCTG CCGAGAAAGT ATCCATCATG  
4200

GCTGATGCAA TGCGGCGGCT GCATACGCTT GATCCGGCTA CCTGCCCATT CGACCACCAA  
4260

GCGAAACATC GCATCGAGCG AGCACGTACT CGGATGGAAG CCGGTCTTGT CGATCAGGAT  
4320

GATCTGGACG AAGAGCATCA GGGGCTCGCG CCAGCCGAAC TGTTCGCCAG GCTCAAGGCG  
4380

CGCATGCCCC ACGGCGAGGA TCTCGTCGTG ACCCATGGCG ATGCCTGCTT GCCGAATATC  
4440

ATGGTGGAAA ATGGCCGCTT TTCTGGATTC ATCGACTGTG GCCGGCTGGG TGTGGCGGAC  
4500

CGCTATCAGG ACATAGCGTT GGCTACCCGT GATATTGCTG AAGAGCTTGG CGGCGAATGG  
4560

GCTGACCGCT TCCTCGTGCT TTACGGTATC GCCGCTCCCG ATTCGCAGCG CATCGCCTTC  
4620

TATCGCCTTC TTGACGAGTT CTTCTGAGCG GGACTCTGGG GTTCGAAATG ACCGACCAAG  
4680

CGACGCCCAA CCTGCCATCA CGAGATTTCG ATTCCACCGC CGCCTTCTAT GAAAGGTTGG  
4740

GCTTCGGAAT CGTTTTCCGG GACGCCGGCT GGATGATCCT CCAGCGCGGG GATCTCATGC  
4800

TGGAGTTCTT CGCCCACCCC AACTTGTTTA TTGCAGCTTA TAATGGTTAC AAATAAAGCA  
4860



ATAGCATCAC AAATTTACACA AATAAAGCAT TTTTTTCACT GCATTCTAGT TGTGGTTTGT  
4920

CCAAACTCAT CAATGTATCT TATCATGTCT GTATACCGTC GACCTCTAGC TAGAGCTTGG  
4980

CGTAATCATG GTCATAGCTG TTTCTGTGT GAAATTGTTA TCCGCTCACA ATTCCACACA  
5040

ACATACGAGC CGGAAGCATA AAGTGTAAG CCTGGGGTGC CTAATGAGTG AGCTAACTCA  
5100

CATTAATTGC GTTGCGCTCA CTGCCCCTT TCCAGTCGGG AAACCTGTCTG TGCCAGCTGC  
5160

ATTAATGAAT CGGCCAACGC GCGGGGAGAG GCGGTTTGCG TATTGGGCGC TCTTCCGCTT  
5220

CCTCGCTCAC TGA CTGCTG CGCTCGGTCTG TTCGGCTGCG GCGAGCGGTA TCAGCTCACT  
5280

CAAAGGCGGT AATACGGTTA TCCACAGAAT CAGGGGATAA CGCAGGAAAG AACATGTGAG  
5340

CAAAAGGCCA GCAAAGGCC AGGAACCGTA AAAAGGCCGC GTTGCTGGCG TTTTCCATA  
5400

GGCTCCGCCC CCTGACGAG CATCACAAA ATCGACGCTC AAGTCAGAGG TGGCGAAACC  
5460

CGACAGGACT ATAAAGATAC CAGGCGTTTC CCCCTGGAAG CTCCTCGTG CGCTCTCCTG  
5520

TTCCGACCCT GCCGCTTACC GGATACCTGT CCGCCTTTCT CCCTTCGGGA AGCGTGGCGC  
5580

TTTCTCAATG CTCACGCTGT AGGTATCTCA GTTCGGTGTA GGTCGTTCTG TCCAAGCTGG  
5640

GCTGTGTGCA CGAACCCCCC GTTCAGCCCC ACCGCTGCGC CTTATCCGGT AACTATCGTC  
5700

TTGAGTCCAA CCCGGTAAGA CACGACTTAT CGCCACTGGC AGCAGCCACT GGTAACAGGA  
5760

TTAGCAGAGC GAGGTATGTA GCGGGTGCTA CAGAGTTCTT GAAGTGGTGG CCTAACTACG  
5820

GCTACACTAG AAGGACAGTA TTTGGTATCT GCGCTCTGCT GAAGCCAGTT ACCTTCGGAA  
5880

AAAGAGTTGG TAGCTCTTGA TCCGGCAAAC AAACCACCGC TGGTAGCGGT GGTTTTTTTG  
5940

TTTGCAAGCA GCAGATTACG CGCAGAAAAA AAGGATCTCA AGAAGATCCT TTGATCTTTT  
6000

CTACGGGGTC TGACGCTCAG TGGAACGAAA ACTCACGTTA AGGGATTTTG GTCATGAGAT  
6060

TATCAAAAAG GATCTTCACC TAGATCCTTT TAAATTAAAA ATGAAGTTTT AAATCAATCT  
6120

AAAGTATATA TGAGTAACT TGGTCTGACA GTTACCAATG CTTAATCAGT GAGGCACCTA  
6180

TCTCAGCGAT CTGTCTATTT CGTTCATCCA TAGTTGCCTG ACTCCCCGTC GTGTAGATAA  
6240

CTACGATACG GGAGGGCTTA CCATCTGGCC CCAGTGCTGC AATGATACCG CGAGACCCAC  
6300

GCTCACCGGC TCCAGATTTA TCAGCAATAA ACCAGCCAGC CGGAAGGGCC GAGCGCAGAA  
6360

GTGGTCCTGC AACTTTATCC GCCTCCATCC AGTCTATTAA TTGTTGCCGG GAAGCTAGAG  
6420

TAAGTAGTTC GCCAGTTAAT AGTTTGCGCA ACGTTGTTGC CATTGCTACA GGCATCGTGG  
6480

TGTCACGCTC GTCGTTTGGT ATGGCTTCAT TCAGCTCCGG TTCCCAACGA TCAAGGCGAG  
6540

TTACATGATC CCCCATGTTG TGCAAAAAAG CGGTTAGCTC CTTCCGGTCCT CCGATCGTTG  
6600

TCAGAAGTAA GTTGGCCGCA GTGTTATCAC TCATGGTTAT GGCAGCACTG CATAATTCTC  
6660

TTACTGTCAT GCCATCCGTA AGATGCTTTT CTGTGACTGG TGAGTACTCA ACCAAGTCAT  
6720

TCTGAGAATA GTGTATGCGG CGACCGAGTT GCTCTTGCCC GCGTCAATA CGGGATAATA  
6780

CCGCGCCACA TAGCAGAACT TTAAAAGTGC TCATCATTGG AAAACGTTCT TCGGGGCGAA  
6840

AACTCTCAAG GATCTTACCG CTGTTGAGAT CCAGTTCGAT GTAACCCACT CGTGCACCCA  
6900

ACTGATCTTC AGCATCTTTT ACTTTCACCA GCGTTTCTGG GTGAGCAAAA ACAGGAAGGC  
6960

AAAATGCCGC AAAAAAGGGA ATAAGGGCGA CACGGAAATG TTGAATACTC ATACTCTTCC  
7020

TTTTTCAATA TTATTGAAGC ATTTATCAGG GTTATTGTCT CATGAGCGGA TACATATTTG  
7080

AATGTATTTA GAAAAATAAA CAAATAGGGG TTCCGCGCAC ATTTCCCCGA AAAGTGCCAC  
7140

CTGACGTC  
7148

## (2) INFORMATION FOR SEQ ID NO:8:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7469 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: circular

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

GACGGATCGG GAGATCTCCC GATCCCCTAT GGTCGACTCT CAGTACAATC TGCTCTGATG  
60

CCGCATAGTT AAGCCAGTAT CTGCTCCCTG CTTGTGTGTT GGAGGTCGCT GAGTAGTGCG  
120

CGAGCAAAAT TTAAGCTACA ACAAGGCAAG GCTTGACCGA CAATTGCATG AAGAATCTGC  
180

TTAGGGTTAG GCGTTTTGCG CTGCTTCGCG ATGTACGGGC CAGATATACG CGTTGACATT  
240

GATTATTGAC TAGTTATTAA TAGTAATCAA TTACGGGGTC ATTAGTTCAT AGCCCATATA  
300

TGGAGTCCG CGTTACATAA CTTACGGTAA ATGGCCCGCC TGGCTGACCG CCCAACGACC  
360

CCCGCCCAT T GACGTCAATA ATGACGTATG TTCCCATAGT AACGCCAATA GGGACTTTCC  
420

ATTGACGTCA ATGGGTGGAC TATTTACGGT AAAGTGGCCA CTTGGCAGTA CATCAAGTGT  
480

ATCATATGCC AAGTACGCCC CCTATTGACG TCAATGACGG TAAATGGCCC GCCTGGCATT  
540

ATGCCCAGTA CATGACCTTA TGGGACTTTC CTAATTGGCA GTACATCTAC GTATTAGTCA  
600

TCGCTATTAC CATGGTGATG CGGTTTTGGC AGTACATCAA TGGGCGTGGA TAGCGGTTTG  
660

ACTCACGGGG ATTTCCAAGT CTCCACCCCA TTGACGTCAA TGGGAGTTTG TTTTGGCACC  
720

AAAATCAACG GGACTTTCCA AAATGTCGTA ACAACTCCGC CCCATTGACG CAAATGGGCG  
780

GTAGGCGTGT ACGGTGGGAG GTCTATATAA GCAGAGCTCT CTGGCTAACT AGAGAACCCA  
840

CTGCTTACTG GCTTATCGAA ATTAATACGA CTCACTATAG GGAGACCCAA GCTTGGTACC  
900

GAGCTCGGAT CTGAATTCGA GCTCGCTGTT GGGCTCGCGG TTGAGGACAA ACTCTTCGCG  
960

GTCTTTCCAG TACTCTTGGA TCGGAAACCC GTCGGCCTCC GAACGGTACT CCGCCACCGA  
1020

GGGACCTGAG CGAGTCCGCA TCGACCGGAT CGGAAAACCT CTCGAGAAAG GCGTCTAACC  
1080

AGTCACAGTC GCAAGGTAGG CTGAGCACCG TGGCGGGCGG CAGCGGGTGG CGGTCGGGGT  
1140

TGTTTCTGGC GGAGGTGCTG CTGATGATGT AATTAAAGTA GGCGGTCTTG AGACGGCGGA  
1200

TGGTCGAGGT GAGGTGTGGC AGGCTTGAGA TCCAAGATGA AGCGCGCAAG ACCGTCTGAA  
1260

GATACCTTCA ACCCCGTGTA TCCATATGAC ACGGAAACCG GTCCTCCAAC TGTGCCTTTT  
1320

CTTACTCCTC CCTTTGTATC CCCCAATGGG TTTCAAGAGA GTCCCCCTGG GGTACTCTCT  
1380

TTGCGCCTAT CCGAACCTCT AGTTACCTCC AATGGCATGC TTGCGCTCAA AATGGGCAAC  
1440

GGCCTCTCTC TGGACGAGGC CGGCAACCTT ACCTCCCAAA ATGTAACCAC TGTGAGCCCA  
1500

CCTCTCAAAA AAACCAAGTC AAACATAAAC CTGGAAATAT CTGCACCCCT CACAGTTACC  
1560

TCAGAAGCCC TAACTGTGGC TGCCGCCGCA CCTCTAATGG TCGCGGGCAA CACACTCACC  
1620

ATGCAATCAC AGGCCCCGCT AACCGTGCAC GACTCCAAAC TTAGCATTGC CACCCAAGGA  
1680

CCCCTCACAG TGTCAGAAGG AAAGCTAGCC CTGCAAACAT CAGGCCCCCT CACCACCACC  
1740

GATAGCAGTA CCCTTACTAT CACTGCCTCA CCCCCTCTAA CTACTGCCAC TGGTAGCTTG  
1800

GGCATTGACT TGAAAGAGCC CATTTATACA CAAAATGGAA AACTAGGACT AAAGTACGGG  
1860

GCTCCTTTGC ATGTAACAGA CGACCTAAAC ACTTTGACCG TAGCAACTGG TCCAGGTGTG  
1920

ACTATTAATA ATACTTCCTT GCAAACATAA GTTACTGGAG CCTTGGGTTT TGATTCACAA  
1980

GGCAATATGC AACTTAATGT AGCAGGAGGA CTAAGGATTG ATTCTCAAAA CAGACGCCTT  
2040

ATACTTGATG TTAGTTATCC GTTTGATGCT CAAAACCAAC TAAATCTAAG ACTAGGACAG  
2100

GGCCCTCTTT TTATAAACTC AGCCCACAAC TTGGATATTA ACTACAACAA AGGCCTTTAC  
2160

TTGTTTACAG CTTCAAACAA TTCCAAAAAG CTTGAGGTTA ACCTAAGCAC TGCCAAGGGG  
2220

TTGATGTTTG ACGCTACAGC CATAGCCATT AATGCAGGAG ATGGGCTTGA ATTTGGTTCA  
2280

CCTAATGCAC CAAACACAAA TCCCCTCAA AAAAAATTG GCCATGGCCT AGAATTTGAT  
2340

TCAAACAAGG CTATGGTTCC TAACTAGGA ACTGGCCTTA GTTTTGACAG CACAGGTGCC  
2400

ATTACAGTAG GAAACAAAAA TAATGATAAG CTAACTTTGT GGACCACACC AGCTCCATCT  
2460

CCTAACTGTA GACTAAATGC AGAGAAAGAT GCTAACTCA CTTTGGTCTT AACAAAATGT  
2520

GGCAGTCAAA TACTTGCTAC AGTTTCAGTT TTGGCTGTTA AAGGCAGTTT GGCTCCAATA  
2580

TCTGGAACAG TTCAAAGTGC TCATCTTATT ATAAGATTTG ACGAAAATGG AGTGCTACTA  
2640

AACAATTCCT TCCTGGACCC AGAATATTGG AACTTTAGAA ATGGAGATCT TACTGAAGGC  
2700

ACAGCCTATA CAAACGCTGT TGGATTTATG CCTAACCTAT CAGCTTATCC AAAATCTCAC  
2760

GGTAAACTG CCAAAAGTAA CATTGTCAGT CAAGTTTACT TAAACGGAGA CAAAACTAAA  
2820

CCTGTAACAC TAACCATTAC ACTAAACGGT ACACAGGAAA CAGGAGACAC AACTCCAAGT  
2880

GCATACTCTA TGTCAATTTTC ATGGGACTGG TCTGGCCACA ACTACATTAA TGAAATATTT  
2940

GCCACATCCT CTTACACTTT TTCATACATT GCCCAAGAAT AAAGAAGCGG CCGCTCGAGC  
3000

ATGCATCTAG AGGGCCCTAT TCTATAGTGT CACCTAAATG CTAGAGCTCG CTGATCAGCC  
3060

TCGACTGTGC CTTCTAGTTG CCAGCCATCT GTTGTTTGCC CCTCCCCCGT GCCTTCCTTG  
3120

ACCCTGGAAG GTGCCACTCC CACTGTCCTT TCCTAATAAA ATGAGGAAAT TGCATCGCAT  
3180

TGTCTGAGTA GGTGTCATTC TATTCTGGGG GGTGGGGTGG GGCAGGACAG CAAGGGGGAG  
3240

GATTGGGAAG ACAATAGCAG GCATGCTGGG GATGCGGTGG GCTCTATGGC TTCTGAGGCG  
3300

GAAAGAACCA GCTGGGGCTC TAGGGGGTAT CCCCACGCGC CCTGTAGCGG CGCATTAAAGC  
3360

GCGGCGGGTG TGGTGGTTAC GCGCAGCGTG ACCGCTACAC TTGCCAGCGC CCTAGCGCCC  
3420

GCTCCTTTTCG CTTTCTTCCC TTCCTTTCTC GCCACGTTTC CCGGCTTTCC CCGTCAAGCT  
3480

CTAAATCGGG GCATCCCTTT AGGGTTCCGA TTTAGTGCTT TACGGCACCT CGACCCCAAA  
3540



AAACTTGATT AGGGTGATGG TTCACGTAGT GGGCCATCGC CCTGATAGAC GGTTTTTCGC  
3600

CCTTTGACGT TGGAGTCCAC GTTCTTTAAT AGTGGACTCT TGTTCCAAAC TGGAACAACA  
3660

CTCAACCCTA TCTCGGTCTA TTCTTTTGAT TTATAAGGGA TTTTGGGGAT TTCGGCCTAT  
3720

TGGTTAAAAA ATGAGCTGAT TTAACAAAAA TTAAACGCGA ATTAATTCTG TGGAATGTGT  
3780

GTCAGTTAGG GTGTGGAAAG TCCCCAGGCT CCCCAGGCAG GCAGAAGTAT GCAAAGCATG  
3840

CATCTCAATT AGTCAGCAAC CAGGTGTGGA AAGTCCCCAG GCTCCCCAGC AGGCAGAAGT  
3900

ATGCAAAGCA TGCATCTCAA TTAGTCAGCA ACCATAGTCC CGCCCCCTAAC TCCGCCCCATC  
3960

CCGCCCCCTAA CTCCGCCCAG TTCCGCCCCAT TCTCCGCCCC ATGGCTGACT AATTTTTTTTT  
4020

ATTTATGCAG AGGCCGAGGC CGCCTCTGCC TCTGAGCTAT TCCAGAAGTA GTGAGGAGGC  
4080

TTTTTTGGAG GCCTAGGCTT TTGCAAAAAG CTCCCGGGAG CTTGTATATC CATTTTCGGA  
4140

TCTGATCAAG AGACAGGATG AGGATCGTTT CGCATGATTG AACAAGATGG ATTGCACGCA  
4200

GGTTCTCCGG CCGCTTGGGT GGAGAGGCTA TTCGGCTATG ACTGGGCACA ACAGACAATC  
4260

GGCTGCTCTG ATGCCGCCGT GTTCCGGCTG TCAGCGCAGG GGCGCCCCGGT TCTTTTTGTC  
4320

AAGACCGACC TGTCCGGTGC CCTGAATGAA CTGCAGGACG AGGCAGCGCG GCTATCGTGG  
4380

CTGGCCACGA CGGGCGTTCC TTGCGCAGCT GTGCTCGACG TTGTCACTGA AGCGGGAAGG  
4440

GACTGGCTGC TATTGGGCGA AGTGCCGGGG CAGGATCTCC TGTCATCTCA CCTTGCTCCT  
4500

GCCGAGAAAG TATCCATCAT GGCTGATGCA ATGCGGCGGC TGCATACGCT TGATCCGGCT  
4560

ACCTGCCCAT TCGACCACCA AGCGAAACAT CGCATCGAGC GAGCACGTAC TCGGATGGAA  
4620

GCCGGTCTTG TCGATCAGGA TGATCTGGAC GAAGAGCATC AGGGGCTCGC GCCAGCCGAA  
4680

CTGTTGCCCA GGCTCAAGGC GCGCATGCCC GACGGCGAGG ATCTCGTCGT GACCCATGGC  
4740

GATGCCTGCT TGCCGAATAT CATGGTGGAA AATGGCCGCT TTTCTGGATT CATCGACTGT  
4800

GGCCGGCTGG GTGTGGCGGA CCGCTATCAG GACATAGCGT TGGCTACCCG TGATATTGCT  
4860

GAAGAGCTTG GCGGCGAATG GGCTGACCGC TTCCTCGTGC TTTACGGTAT CGCCGCTCCC  
4920

GATTTCGCAGC GCATCGCCTT CTATCGCCTT CTTGACGAGT TCTTCTGAGC GGGACTCTGG  
4980

GGTTCGAAAT GACCGACCAA GCGACGCCCA ACCTGCCATC ACGAGATTTC GATTCCACCG  
5040

CCGCCTTCTA TGAAAGGTTG GGCTTCGGAA TCGTTTTCCG GGACGCCGGC TGGATGATCC  
5100

TCCAGCGCGG GGATCTCATG CTGGAGTTCT TCGCCCACCC CAACTTGTTT ATTGCAGCTT  
5160

ATAATGGTTA CAAATAAAGC AATAGCATCA CAAATTTAC AAATAAAGCA TTTTTTTCAC  
5220

TGCATTCTAG TTGTGGTTTG TCCAAACTCA TCAATGTATC TTATCATGTC TGTATACCGT  
5280

CGACCTCTAG CTAGAGCTTG GCGTAATCAT GGTCATAGCT GTTTCCTGTG TGAAATTGTT  
5340

ATCCGCTCAC AATTCACAC AACATACGAG CCGGAAGCAT AAAGTGTAAG GCCTGGGGTG  
5400

CCTAATGAGT GAGCTAACTC ACATTAATTG CGTTGCGCTC ACTGCCCCTG TTCCAGTCGG  
5460

GAAACCTGTC GTGCCAGCTG CATTAATGAA TCGGCCAACG CGCGGGGAGA GGCGGTTTGC  
5520

GTATTGGGCG CTCTTCCGCT TCCTCGCTCA CTGACTCGCT GCGCTCGGTC GTTCGGCTGC  
5580

GGCGAGCGGT ATCAGCTCAC TCAAAGGCGG TAATACGGTT ATCCACAGAA TCAGGGGATA  
5640

ACGCAGGAAA GAACATGTGA GCAAAGGCC AGCAAAGGC CAGGAACCGT AAAAAGGCCG  
5700

CGTTGCTGGC GTTTTTCCAT AGGCTCCGCC CCCCTGACGA GCATCACAAA AATCGACGCT  
5760

CAAGTCAGAG GTGGCGAAAC CCGACAGGAC TATAAAGATA CCAGGCGTTT CCCCTGGAA  
5820

GCTCCCTCGT GCGCTCTCCT GTTCCGACCC TGCCGCTTAC CGGATACCTG TCCGCCTTTC  
5880

TCCCTTCGGG AAGCGTGGCG CTTTCTCAAT GCTCACGCTG TAGGTATCTC AGTTCGGTGT  
5940

AGGTCGTTCG CTCCAAGCTG GGCTGTGTGC ACGAACCCCC CGTTCAGCCC GACCGCTGCG  
6000

CCTTATCCGG TAACTATCGT CTTGAGTCCA ACCCGGTAAG ACACGACTTA TCGCCACTGG  
6060

CAGCAGCCAC TGGTAACAGG ATTAGCAGAG CGAGGTATGT AGGCGGTGCT ACAGAGTTCT  
6120

TGAAGTGGTG GCCTAACTAC GGCTACACTA GAAGGACAGT ATTTGGTATC TCGCTCTGC  
6180

TGAAGCCAGT TACCTTCGGA AAAAGAGTTG GTAGCTCTTG ATCCGGCAAA CAAACCACCG  
6240

CTGGTAGCGG TGGTTTTTTT GTTTGCAAGC AGCAGATTAC GCGCAGAAAA AAAGGATCTC  
6300

AAGAAGATCC TTTGATCTTT TCTACGGGGT CTGACGCTCA GTGGAACGAA AACTCACGTT  
6360

AAGGGATTTT GGTCATGAGA TTATCAAAAA GGATCTTCAC CTAGATCCTT TTAAATTAAA  
6420

AATGAAGTTT TAAATCAATC TAAAGTATAT ATGAGTAAAC TTGGTCTGAC AGTTACCAAT  
6480

GCTTAATCAG TGAGGCACCT ATCTCAGCGA TCTGTCTATT TCGTTCATCC ATAGTTGCCT  
6540

GACTCCCCGT CGTGTAGATA ACTACGATAC GGGAGGGCTT ACCATCTGGC CCCAGTGCTG  
6600

CAATGATACC GCGAGACCCA CGCTCACCGG CTCCAGATTT ATCAGCAATA AACCAGCCAG  
6660

CCGGAAGGGC CGAGCGCAGA AGTGGTCCTG CAACTTTATC CGCCTCCATC CAGTCTATTA  
6720

ATTGTTGCCG GGAAGCTAGA GTAAGTAGTT CGCCAGTTAA TAGTTTGCGC AACGTTGTTG  
6780

CCATTGCTAC AGGCATCGTG GTGTCACGCT CGTCGTTTGG TATGGCTTCA TTCAGCTCCG  
6840

GTTCCCAACG ATCAAGGCGA GTTACATGAT CCCCCATGTT GTGCAAAAAA GCGGTTAGCT  
6900

CCTTCGGTCC TCCGATCGTT GTCAGAAGTA AGTTGGCCGC AGTGTTATCA CTCATGGTTA  
6960

TGGCAGCACT GCATAATTCT CTTACTGTCA TGCCATCCGT AAGATGCTTT TCTGTGACTG  
7020

GTGAGTACTC AACCAAGTCA TTCTGAGAAT AGTGTATGCG GCGACCGAGT TGCTCTTGCC  
7080

CGGCGTCAAT ACGGGATAAT ACCGCGCCAC ATAGCAGAAC TTTAAAAGTG CTCATCATTG  
7140

GAAAACGTTC TTCGGGGCGA AACTCTCAA GGATCTTACC GCTGTTGAGA TCCAGTTCGA  
7200

TGTAACCCAC TCGTGCACCC AACTGATCTT CAGCATCTTT TACTTTCACC AGCGTTTCTG  
7260

GGTGAGCAAA AACAGGAAGG CAAAATGCCG CAAAAAAGGG AATAAGGGCG ACACGGAAAT  
7320

GTTGAATACT CATACTCTTC CTTTTTCAAT ATTATTGAAG CATTTATCAG GGTTATTGTC  
7380

TCATGAGCGG ATACATATTT GAATGTATTT AGAAAAATAA ACAAATAGGG GTTCCGCGCA  
7440

CATTTCCCCG AAAAGTGCCA CCTGACGTC  
7469

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 28 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

TGCTTAAGCG GCCGCGAAGG AGAAGTCC  
28

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

CCGAGCTAGC GACTGAAAAT GAG

23

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

CCTCTCGAGA GACAGCAAGA CAC

23

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 11152 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: circular

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

AAGCTTGGGC AGAAATGGTT GAACTCCCGA GAGTGCCTA CACCTAGGGG AGAAGCAGCC  
60

AAGGGGTTGT TTCCCACCAA GGACGACCCG TCTGCGCACA AACGGATGAG CCCATCAGAC  
120

AAAGACATAT TCATTCTCTG CTGCAAACTT GGCATAGCTC TGCTTTGCCT GGGGCTATTG  
180

GGGGAAGTTG CGGTTCGTGC TCGCAGGGCT CTCACCCTTG ACTCTTTTAA TAGCTCTTCT  
240

GTGCAAGATT ACAATCTAAA CAATTCGGAG AACTCGACCT TCCTCCTGAG GCAAGGACCA  
300

CAGCCAACTT CCTCTTACAA GCCGCATCGA TTTTGTCTT CAGAAATAGA AATAAGAATG  
360

CTTGCTAAAA ATTATATTTT TACCAATAAG ACCAATCCAA TAGGTAGATT ATTAGTTACT  
420

ATGTTAAGAA ATGAATCATT ATCTTTTAGT ACTATTTTAA CTCAAATTCA GAAGTTAGAA  
480

ATGGGAATAG AAAATAGAAA GAGACGCTCA ACCTCAATTG AAGAACAGGT GCAAGGACTA  
540

TTGACCACAG GCCTAGAAGT AAAAAAGGGA AAAAAGAGTG TTTTGTCAA AATAGGAGAC  
600

AGGTGGTGGC AACCAGGGAC TTATAGGGGA CCTTACATCT ACAGACCAAC AGATGCCCCC  
660

TTACCATATA CAGGAAGATA TGA CTTAAAT TGGGATAGGT GGGTTACAGT CAATGGCTAT  
720



AAAGTGTTAT ATAGATCCCT CCCTTTTCGT GAAAGACTCG CCAGAGCTAG ACCTCCTTGG  
780

TGTATGTTGT CTCAAGAAGA AAAAGACGAC ATGAAACAAC AGGTACATGA TTATATTTAT  
840

CTAGGAACAG GAATGCACTT TTGGGGAAAG ATTTTCCATA CCAAGGAGGG GACAGTGGCT  
900

GGACTAATAG AACATTATTC TGCAAAAACAT CATGGCATGA GTTATTATGA ATAGCCTTTA  
960

TTGGCCCAAC CTTGCGGTTC CCAGGGCTTA AGTAAGTTTT TGGTTACAAA CTGTTCTTAA  
1020

AACGAGGATG TGAGACAAGT GGTTTCCTGA CTTGGTTTGG TATCAAAGGT TCTGATCTGA  
1080

GCTCTGAGTG TTCTATTTTC CTATGTTCTT TTGGAATTTA TCCAAATCTT ATGTAAATGC  
1140

TTATGTAAAC CAAGATATAA AAGAGTGCTG ATTTTTTGAG TAAACTTGCA ACAGTCCTAA  
1200

CATTCACCTC TTGTGTGTTT GTGTCTGTTT GCCATCCCGT CTCCGCTCGT CACTTATCCT  
1260

TCACTTTCCA GAGGGTCCCC CCGCAGACCC CGGCGACCCT CAGGTCGGCC GACTGCGGCA  
1320

GCTGGCGCCC GAACAGGGAC CCTCGGATAA GTGACCCTTG TCTCTATTTT TACTATTTGG  
1380

TGTTTGTCTT GTATTGTCTC TTTCTTGTCT GGCTATCATC ACAAGAGCGG AACGGACTCA  
1440

CCATAGGGAC CAAGCTAGCG ACTGAAAATG AGACATATTA TCTGCCACGG AGGTGTTATT  
1500

ACCGAAGAAA TGGCCGCCAG TCTTTTGGAC CAGCTGATCG AAGAGGTACT GGCTGATAAT  
1560

CTTCCACCTC CTAGCCATTT TGAACCACCT ACCCTTCACG AACTGTATGA TTTAGACGTG  
1620

ACGGCCCCCG AAGATCCCAA CGAGGAGGCG GTTTCGCAGA TTTTCCCGA CTCTGTAATG  
1680

TTGGCGGTGC AGGAAGGGAT TGACTTACTC ACTTTTCCGC CGGCGCCCCG TTCTCCGGAG  
1740

CCGCCTCACC TTTCCCGGCA GCCCGAGCAG CCGGAGCAGA GAGCCTTGGG TCCGGTTTCT  
1800

ATGCCAAACC TTGTACCGGA GGTGATCGAT CTTACCTGCC ACGAGGCTGG CTTTCCACCC  
1860

AGTGACGACG AGGATGAAGA GGGTGAGGAG TTTGTGTTAG ATTATGTGGA GCACCCCGGG  
1920

CACGGTTGCA GGTCTTGTC A TTATCACCGG AGGAATACGG GGGACCCAGA TATTATGTGT  
1980

TCGCTTTGCT ATATGAGGAC CTGTGGCATG TTTGTCTACA GTAAGTGAAA ATTATGGGCA  
2040

GTGGGTGATA GAGTGGTGGG TTTGGTGTGG TAATTTTTTT TTTAATTTTT ACAGTTTGT  
2100

GGTTTAAAGA ATTTTGTATT GTGATTTTTT TAAAAGGTCC TGTGTCTGAA CCTGAGCCTG  
2160

AGCCCGAGCC AGAACCGGAG CCTGCAAGAC CTACCCGCCG TCCTAAAATG GCGCCTGCTA  
2220

TCCTGAGACG CCCGACATCA CCTGTGTCTA GAGAATGCAA TAGTAGTACG GATAGCTGTG  
2280

ACTCCGGTCC TTCTAACACA CCTCCTGAGA TACACCCGGT GGTCCCGCTG TGCCCCATTA  
2340

AACCAGTTGC CGTGAGAGTT GGTGGGCGTC GCCAGGCTGT GGAATGTATC GAGGACTTGC  
2400

TTAACGAGCC TGGGCAACCT TTGGACTTGA GCTGTAAACG CCCCAGGCCA TAAGGTGTAA  
2460

ACCTGTGATT GCGTGTGTGG TTAACGCCTT TGTTTGCTGA ATGAGTTGAT GTAAGTTTAA  
2520

TAAAGGGTGA GATAATGTTT AACTTGCATG GCGTGTTAAA TGGGGCGGGG CTAAAGGGT  
2580

ATATAATGCG CCGTGGGCTA ATCTTGGTTA CATCTGACCT CATGGAGGCT TGGGAGTGTT  
2640

TGGAAGATTT TTCTGCTGTG CGTAACTTGC TGGAACAGAG CTCTAACAGT ACCTCTTGGT  
2700

TTTGGAGGTT TCTGTGGGGC TCATCCCAGG CAAAGTTAGT CTGCAGAATT AAGGAGGATT  
2760

ACAAGTGGGA ATTTGAAGAG CTTTGTAAAT CCTGTGGTGA GCTGTTTGAT TCTTTGAATC  
2820

TGGGTCACCA GGCGCTTTTC CAAGAGAAGG TCATCAAGAC TTTGGATTTT TCCACACCGG  
2880

GGCGCGCTGC GGCTGCTGTT GCTTTTTTGA GTTTTATAAA GGATAAATGG AGCGAAGAAA  
2940

CCCATCTGAG CGGGGGGTAC CTGCTGGATT TTCTGGCCAT GCATCTGTGG AGAGCGGTTG  
3000

TGAGACACAA GAATCGCCTG CTACTGTTGT CTTCCGTCCG CCCGGCGATA ATACCGACGG  
3060

AGGAGCAGCA GCAGCAGCAG GAGGAAGCCA GCGGCGGCG GCAGGAGCAG AGCCCATGGA  
3120

ACCCGAGAGC CGGCCTGGAC CCTCGGGAAT GAATGTTGTA CAGGTGGCTG AACTGTATCC  
3180

AGAACTGAGA CGCATTTTGA CAATTACAGA GGATGGGCAG GGGCTAAAGG GGGTAAAGAG  
3240

GGAGCGGGGG GCTTGTGAGG CTACAGAGGA GGCTAGGAAT CTAGCTTTTA GCTTAATGAC  
3300

CAGACACCGT CCTGAGTGTA TTACTTTTCA ACAGATCAAG GATAATTGCG CTAATGAGCT  
3360

TGATCTGCTG GCGCAGAAGT ATTCCATAGA GCAGCTGACC ACTTACTGGC TGCAGCCAGG  
3420

GGATGATTTT GAGGAGGCTA TTAGGGTATA TGCAAAGGTG GCACTTAGGC CAGATTGCAA  
3480

GTACAAGATC AGCAAACCTG TAAATATCAG GAATTGTTGC TACATTTCTG GGAACGGGGC  
3540

CGAGGTGGAG ATAGATACGG AGGATAGGGT GGCCTTTAGA TGTAGCATGA TAAATATGTG  
3600

GCCGGGGGTG CTTGGCATGG ACGGGGTGGT TATTATGAAT GTAAGGTTTA CTGGCCCCAA  
3660

TTTTAGCGGT ACGGTTTTCC TGGCCAATAC CAACCTTATC CTACACGGTG TAAGCTTCTA  
3720

TGGGTTTAAC AATACCTGTG TGGAAGCCTG GACCGATGTA AGGGTTCGGG GCTGTGCCTT  
3780

TTACTGCTGC TGGAAGGGGG TGGTGTGTCG CCCCAAAGC AGGGCTTCAA TTAAGAAATG  
3840

CCTCTTTGAA AGGTGTACCT TGGGTATCCT GTCTGAGGGT AACTCCAGGG TGCGCCACAA  
3900

TGTGGCCTCC GACTGTGGTT GCTTCATGCT AGTGAAAAGC GTGGCTGTGA TTAAGCATAA  
3960

CATGGTATGT GGCAACTGCG AGGACAGGGC CTCTCAGATG CTGACCTGCT CGGACGGCAA  
4020

CTGTCACCTG CTGAAGACCA TTCACGTAGC CAGCCACTCT CGCAAGGCCT GGCCAGTGTT  
4080

TGAGCATAAC ATACTGACCC GCTGTTCCTT GCATTGGGT AACAGGAGGG GGGTGTTCCT  
4140

ACCTTACCAA TGCAATTTGA GTCACACTAA GATATTGCTT GAGCCCGAGA GCATGTCCAA  
4200

GGTGAACCTG AACGGGGTGT TTGACATGAC CATGAAGATC TGGAAGGTGC TGAGGTACGA  
4260

TGAGACCCGC ACCAGGTGCA GACCCTGCGA GTGTGGCGGT AAACATATTA GGAACCAGCC  
4320

TGTGATGCTG GATGTGACCG AGGAGCTGAG GCCCGATCAC TTGGTGCTGG CCTGCACCCG  
4380

CGCTGAGTTT GGCTCTAGCG ATGAAGATAC AGATTGAGGT ACTGAAATGT GTGGGCGTGG  
4440

CTTAAGGGTG GGAAAGAATA TATAAGGTGG GGGTCTTATG TAGTTTTGTA TCTGTTTTGC  
4500

AGCAGCCGCC GCCGCCATGA GCACCAACTC GTTTGATGGA AGCATTGTGA GCTCATATTT  
4560

GACAACGCGC ATGCCCCCAT GGGCCGGGGT GCGTCAGAAT GTGATGGGCT CCAGCATTGA  
4620

TGGTCGCCCC GTCCTGCCCC CAAACTCTAC TACCTTGACC TACGAGACCG TGTCTGGAAC  
4680

GCCGTTGGAG ACTGCAGCCT CCGCCGCCGC TTCAGCCGCT GCAGCCACCG CCCGCGGGAT  
4740

TGTGACTGAC TTTGCTTTCC TGAGCCCCGCT TGCAAGCAGT GCAGCTTCCC GTTCATCCGC  
4800

CCGCGATGAC AAGTTGACGG CTCTTTTGGC ACAATTGGAT TCTTTGACCC GGGAACCTAA  
4860

TGTCGTTTCT CAGCAGCTGT TGGATCTGCG CCAGCAGGTT TCTGCCCTGA AGGCTTCCTC  
4920

CCCTCCCAAT GCGGTTTAAA ACATAAATAA AAAACCAGAC TCTGTTTGGA TTTGGATCAA  
4980

GCAAGTGTCT TGCTGTCTCT CGAGGGATCT TTGTGAAGGA ACCTTACTTC TGTGGTGTGA  
5040

CATAATTGGA CAAACTACCT ACAGAGATTT AAAGCTCTAA GGTAAATATA AAATTTTTAA  
5100

GTGTATAATG TGTTAAACTA CTGATTCTAA TTGTTTGTGT ATTTTAGATT CCAACCTATG  
5160

GAAGTATGA ATGGGAGCAG TGGTGGAATG CCTTTAATGA GGAAAACCTG TTTTGCTCAG  
5220

AAGAAATGCC ATCTAGTGAT GATGAGGCTA CTGCTGACTC TCAACATTCT ACTCCTCCAA  
5280

AAAAGAAGAG AAAGGTAGAA GACCCCAAGG ACTTTCCTTC AGAATTGCTA AGTTTTTTGA  
5340

GTCATGCTGT GTTTAGTAAT AGAACTCTTG CTTGCTTTGC TATTTACACC ACAAAGGAAA  
5400

AAGCTGCACT GCTATACAAG AAAATTATGG AAAAATATTC TGTAACCTTT ATAAGTAGGC  
5460

ATAACAGTTA TAATCATAAC ATACTGTTTT TTCTTACTCC ACACAGGCAT AGAGTGTCTG  
5520

CTATTAATAA CTATGCTCAA AAATTGTGTA CCTTTAGCTT TTTAATTTGT AAAGGGGTTA  
5580

ATAAGGAATA TTTGATGTAT AGTGCCTTGA CTAGAGATCA TAATCAGCCA TACCACATTT  
5640

GTAGAGGTTT TACTTGCTTT AAAAAACCTC CCACACCTCC CCCTGAACCT GAAACATAAA  
5700

ATGAATGCAA TTGTTGTTGT TAACTTGTTT ATTGCAGCTT ATAATGGTTA CAAATAAAGC  
5760

AATAGCATCA CAAATTTTAC AAATAAAGCA TTTTTTTCAC TGCATTCTAG TTGTGGTTTG  
5820

TCCAAACTCA TCAATGTATC TTATCATGTC TGGATCCGGC TGTGGAATGT GTGTCAGTTA  
5880

GGGTGTGGAA AGTCCCCAGG CTCCCCAGCA GGCAGAAGTA TGCAAAGCAT GCATCTCAAT  
5940

TAGTCAGCAA CCAGGTGTGG AAAGTCCCCA GGCTCCCCAG CAGGCAGAAG TATGCAAAGC  
6000

ATGCATCTCA ATTAGTCAGC AACCATAGTC CCGCCCCCTAA CTCCGCCCAT CCCGCCCTA  
6060

ACTCCGCCCA GTTCCGCCCA TTCTCCGCCC CATGGCTGAC TAATTTTTTT TATTTATGCA  
6120

GAGGCCGAGG CCGCCTCGGC CTCTGAGCTA TTCCAGAAGT AGTGAGGAGG CTTTTTTTGA  
6180

GGCCTAGGCT TTTGCAAAAA GCTTGGACAC AAGACAGGCT TCGGAGATAT GTTTGAGAAT  
6240

ACCACTTTAT CCCGCGTCAG GGAGAGGCAG TCGGTAAAAA GACGCGGACT CATGTGAAAT  
6300

ACTGGTTTTT AGTGCGCCAG ATCTCTATAA TCTCGCGCAA CCTATTTTCC CCTCGAACAC  
6360

TTTTTAAGCC GTAGATAAAC AGGCTGGGAC ACTTCACATG AGCGAAAAAT ACATCGTCAC  
6420

CTGGGACATG TTGCAGATCC ATGCACGTAA ACTCGCAAGC CGACTGATGC CTTCTGAACA  
6480

ATGGAAAGGC ATTATTGCCG TAAGCCGTGG CGGTCTGGTA CCGGGTGCGT TACTGGCGCG  
6540

TGAACTGGGT ATTCGTCATG TCGATACCGT TTGTATTTC AGCTACGATC ACGACAACCA  
6600

GCGCGAGCTT AAAGTGCTGA AACGCGCAGA AGGCGATGGC GAAGGCTTCA TCGTTATTGA  
6660

TGACCTGGTG GATACCGGTG GTACTGCGGT TCGGATTCGT GAAATGTATC CAAAAGCGCA  
6720

CTTTGTCACC ATCTTCGCAA AACCGGCTGG TCGTCCGCTG GTTGATGACT ATGTTGTTGA  
6780

TATCCCGCAA GATACCTGGA TTGAACAGCC GTGGGATATG GCGTCGTAT TCGTCCCGCC  
6840

AATCTCCGGT CGCTAATCTT TTCAACGCCT GGCAGTCCG GCGTTGTTC TTTTAACTT  
6900

CAGGCGGGTT ACAATAGTTT CCAGTAAGTA TTCTGGAGGC TGCATCCATG ACACAGGCAA  
6960



ACCTGAGCGA AACCCCTGTTT AAACCCCGCT TTAAACATCC TGAAACCTCG ACGCTAGTCC  
7020

GCCGCTTTAA TCACGGCGCA CAACCGCCTG TGCAGTCGGC CCTTGATGGT AAAACCATCC  
7080

CTCACTGGTA TCGCATGATT AACCGTCTGA TGTGGATCTG GCGCGGCATT GACCCACGCG  
7140

AAATCCTCGA CGTCCAGGCA CGTATTGTGA TGAGCGATGC CGAACGTACC GACGATGATT  
7200

TATACGATAC GGTGATTGGC TACCGTGGCG GCAACTGGAT TTATGAGTGG GCCCCGGATC  
7260

TTTGTGAAGG AACCTTACTT CTGTGGTGTG ACATAATTGG ACAAACTACC TACAGAGATT  
7320

TAAAGCTCTA AGGTAAATAT AAAATTTTTA AGTGTATAAT GTGTTAACT ACTGATTCTA  
7380

ATTGTTTGTG TATTTTAGAT TCCAACCTAT GGAAGTATG AATGGGAGCA GTGGTGGAAT  
7440

GCCTTTAATG AGGAAAACCT GTTTTGCTCA GAAGAAATGC CATCTAGTGA TGATGAGGCT  
7500

ACTGCTGACT CTCAACATTC TACTCCTCCA AAAAAGAAGA GAAAGGTAGA AGACCCCAAG  
7560

GACTTTCCTT CAGAATTGCT AAGTTTTTTG AGTCATGCTG TGTTTAGTAA TAGAACTCTT  
7620

GCTTGCTTTG CTATTTACAC CACAAAGGAA AAAGCTGCAC TGCTATACAA GAAAATTATG  
7680

GAAAAATATT CTGTAACCTT TATAAGTAGG CATAACAGTT ATAATCATAA CATACTGTTT  
7740

TTTCTTACTC CACACAGGCA TAGAGTGTCT GCTATTAATA ACTATGCTCA AAAATTGTGT  
7800

ACCTTTAGCT TTTTAATTTG TAAAGGGGTT AATAAGGAAT ATTTGATGTA TAGTGCCTTG  
7860

ACTAGAGATC ATAATCAGCC ATACCACATT TGTAGAGGTT TTAAGTGTCT TAAAAACCT  
7920

CCCACACCTC CCCCTGAACC TGAAACATAA AATGAATGCA ATTGTTGTTG TTAAGTGTCT  
7980

TATTGCAGCT TATAATGGTT ACAAATAAAG CAATAGCATC ACAAATTTCA CAAATAAAGC  
8040

ATTTTTTTCA CTGCATTCTA GTTGTGGTTT GTCCAAACTC ATCAATGTAT CTTATCATGT  
8100

CTGGATCCCC AGGAAGCTCC TCTGTGTCCT CATAAACCTT AACCTCCTCT ACTTGAGAGG  
8160

ACATTCCAAT CATAGGCTGC CCATCCACCC TCTGTGTCCT CCTGTAAATT AGGTCACTTA  
8220

ACAAAAAGGA AATTGGGTAG GGGTTTTTCA CAGACCGCTT TCTAAGGGTA ATTTTAAAT  
8280

ATCTGGGAAG TCCCTTCCAC TGCTGTGTTT CAGAAGTGTG GTTAAACAGC CCACAAATGT  
8340

CAACAGCAGA AACATACAAG CTGTCAGCTT TGCACAAGGG CCCAACACCC TGCTCATCAA  
8400

GAAGCACTGT GGTTGCTGTG TTAGTAATGT GCAAAACAGG AGGCACATTT TCCCCACCTG  
8460

TGTAGGTTCC AAAATATCTA GTGTTTTTCAT TTTTACTTGG ATCAGGAACC CAGCACTCCA  
8520

CTGGATAAGC ATTATCCTTA TCCAAAACAG CTTGTGGTC AGTGTTTCATC TGCTGACTGT  
8580

CAACTGTAGC ATTTTTTGGG GTTACAGTTT GAGCAGGATA TTTGGTCCTG TAGTTTGCTA  
8640

ACACACCCTG CAGCTCCAAA GGTCCCCAC CAACAGCAAA AAAATGAAAA TTTGACCCTT  
8700

GAATGGGTTT TCCAGCACCA TTTTCATGAG TTTTTGTGT CCCTGAATGC AAGTTTAACA  
8760

TAGCAGTTAC CCCAATAACC TCAGTTTTAA CAGTAACAGC TTCCCACATC AAAATATTTT  
8820

CACAGGTAA GTCCTCATTT AAATTAGGCA AAGGAATTCT TGAAGACGAA AGGGCCTCGT  
8880

GATACGCCTA TTTTATAGG TTAATGTCAT GATAATAATG GTTCTTAGA CGTCAGGTGG  
8940

CACTTTTCGG GGAAATGTGC GCGGAACCCC TATTTGTTTA TTTTCTAAA TACATTCAAA  
9000

TATGTATCCG CTCATGAGAC AATAACCCTG ATAAATGCTT CAATAATATT GAAAAAGGAA  
9060

GAGTATGAGT ATTCAACATT TCCGTGTCGC CTTATTCCC TTTTTGCGG CATTTTGCCT  
9120

TCCTGTTTTT GCTCACCCAG AAACGCTGGT GAAAGTAAAA GATGCTGAAG ATCAGTTGGG  
9180

TGCACGAGTG GGTTACATCG AACTGGATCT CAACAGCGGT AAGATCCTTG AGAGTTTTCG  
9240

CCCCGAAGAA CGTTTTCCAA TGATGAGCAC TTTTAAAGTT CTGCTATGTG GCGCGGTATT  
9300

ATCCCGTGTT GACGCCGGGC AAGAGCAACT CGGTCGCCGC ATACACTATT CTCAGAATGA  
9360

CTTGGTTGAG TACTCACCAG TCACAGAAAA GCATCTTACG GATGGCATGA CAGTAAGAGA  
9420

ATTATGCAGT GCTGCCATAA CCATGAGTGA TAACACTGCG GCCAACTTAC TTCTGACAAC  
9480

GATCGGAGGA CCGAAGGAGC TAACCGCTTT TTTGCACAAC ATGGGGGATC ATGTAACTCG  
9540

CCTTGATCGT TGGGAACCGG AGCTGAATGA AGCCATACCA AACGACGAGC GTGACACCAC  
9600

GATGCCTGCA GCAATGGCAA CAACGTTGCG CAACTATTA ACTGGCGAAC TACTTACTCT  
9660

AGCTTCCCGG CAACAATTAA TAGACTGGAT GGAGGCGGAT AAAGTTGCAG GACCACTTCT  
9720

GCGCTCGGCC CTTCCGGCTG GCTGGTTTAT TGCTGATAAA TCTGGAGCCG GTGAGCGTGG  
9780

GTCTCGCGGT ATCATTGCAG CACTGGGGCC AGATGGTAAG CCCTCCCGTA TCGTAGTTAT  
9840

CTACACGACG GGGAGTCAGG CAACTATGGA TGAACGAAAT AGACAGATCG CTGAGATAGG  
9900

TGCCTCACTG ATTAAGCATT GGTAAGTGC AGACCAAGTT TACTCATATA TACTTTAGAT  
9960

TGATTTAAAA CTTCATTTTT AATTTAAAAG GATCTAGGTG AAGATCCTTT TTGATAATCT  
10020

CATGACCAAA ATCCCTTAAC GTGAGTTTTC GTTCCACTGA GCGTCAGACC CCGTAGAAAA  
10080

GATCAAAGGA TCTTCTTGAG ATCCTTTTTT TCTGCGCGTA ATCTGCTGCT TGCAAACAAA  
10140

AAAACCACCG CTACCAGCGG TGGTTTGTTT GCCGGATCAA GAGCTACCAA CTCTTTTTC  
10200

GAAGGTAACCT GGCTTCAGCA GAGCGCAGAT ACCAAATACT GTCCTTCTAG TGTAGCCGTA  
10260

GTTAGGCCAC CACTTCAAGA ACTCTGTAGC ACCGCCTACA TACCTCGCTC TGCTAATCCT  
10320

GTTACCAGTG GCTGCTGCCA GTGGCGATAA GTCGTGTCTT ACCGGGTGG ACTCAAGACG  
10380

ATAGTTACCG GATAAGGCGC AGCGGTCGGG CTGAACGGGG GGTTCGTGCA CACAGCCCAG  
10440

CTTGGAGCGA ACGACCTACA CCGAACTGAG ATACCTACAG CGTGAGCTAT GAGAAAGCGC  
10500

CACGCTTCCC GAAGGGAGAA AGGCGGACAG GTATCCGGTA AGCGGCAGGG TCGGAACAGG  
10560

AGAGCGCACG AGGGAGCTTC CAGGGGGAAA CGCCTGGTAT CTTTATAGTC CTGTCGGGTT  
10620

TCGCCACCTC TGAATTGAGC GTCGATTTTT GTGATGCTCG TCAGGGGGGC GGAGCCTATG  
10680

GAAAAACGCC AGCAACGCGG CCTTTTTACG GTTCCTGGCC TTTTGCTGGC CTTTTGCTCA  
10740

CATGTTCTTT CCTGCGTTAT CCCCTGATTC TGTGGATAAC CGTATTACCG CCTTTGAGTG  
10800

AGCTGATACC GCTCGCCGCA GCCGAACGAC CGAGCGCAGC GAGTCAGTGA GCGAGGAAGC  
10860

- 111 -

GGAAGAGCGC CTGATGCGGT ATTTTCTCCT TACGCATCTG TCGGGTATTT CACACCGCAT  
10920

ATGGTGCACT CTCAGTACAA TCTGCTCTGA TGCCGCATAG TTAAGCCAGT ATACACTCCG  
10980

CTATCGCTAC GTGACTGGGT CATGGCTGCG CCCCACACACC CGCCAACACC CGCTGACGCG  
11040

CCCTGACGGG CTTGTCTGCT CCCGGCATCC GCTTACAGAC AAGCTGTGAC CGTCTCCGGG  
11100

AGCTGCATGT GTCAGAGGTT TTCACCGTCA TCACCGAAAC GCGCGAGGCA GC  
11152

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GACGGATCGG GAGATCTCC  
19

(2) INFORMATION FOR SEQ ID NO:14:

- 112 -

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

CCGCCTCAGA AGCCATAGAG CC

22

## (2) INFORMATION FOR SEQ ID NO:15:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 14455 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: circular

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

AAGCTTGGGC AGAAATGGTT GAACTCCCGA GAGTGTCTTA CACCTAGGGG AGAAGCAGCC

60

AAGGGGTTGT TTCCACCAA GGACGACCCG TCTGCGCACA AACGGATGAG CCCATCAGAC

120

AAAGACATAT TCATTCTCTG CTGCAAACCTT GGCATAGCTC TGCTTTGCCT GGGGCTATTG  
180

GGGGAAGTTG CGGTTCTGTC TCGCAGGGCT CTCACCCTTG ACTCTTTTAA TAGCTCTTCT  
240

GTGCAAGATT ACAATCTAAA CAATTCTGGAG AACTCGACCT TCCTCCTGAG GCAAGGACCA  
300

CAGCCAACCTT CCTCTTACAA GCCGCATCGA TTTTGTCTT CAGAAATAGA AATAAGAATG  
360

CTTGCTAAAA ATTATATTTT TACCAATAAG ACCAATCCAA TAGGTAGATT ATTAGTTACT  
420

ATGTTAAGAA ATGAATCATT ATCTTTTAGT ACTATTTTAA CTCAAATTCA GAAGTTAGAA  
480

ATGGGAATAG AAAATAGAAA GAGACGCTCA ACCTCAATTG AAGAACAGGT GCAAGGACTA  
540

TTGACCACAG GCCTAGAAGT AAAAAAGGGA AAAAAGAGTG TTTTGTCAA AATAGGAGAC  
600

AGGTGGTGGC AACCAGGGAC TTATAGGGGA CCTTACATCT ACAGACCAAC AGATGCCCCC  
660

TTACCATATA CAGGAAGATA TGACTTAAAT TGGGATAGGT GGGTTACAGT CAATGGCTAT  
720

AAAGTGTTAT ATAGATCCCT CCCTTTTCGT GAAAGACTCG CCAGAGCTAG ACCTCCTTGG  
780

TGTATGTTGT CTCAAGAAGA AAAAGACGAC ATGAAACAAC AGGTACATGA TTATATTTAT  
840

CTAGGAACAG GAATGCACTT TTGGGGAAAG ATTTTCCATA CCAAGGAGGG GACAGTGGCT  
900



GGACTAATAG AACATTATTC TGCAAAAACT CATGGCATGA GTTATTATGA ATAGCCTTTA  
960

TTGGCCCAAC CTTGCGGTTT CCAGGGCTTA AGTAAGTTT TGGTTACAAA CTGTTCTTAA  
1020

AACGAGGATG TGAGACAAGT GGTTCCTGA CTTGGTTTGG TATCAAAGGT TCTGATCTGA  
1080

GCTCTGAGTG TTCTATTTTC CTATGTTCTT TTGGAATTTA TCCAAATCTT ATGTAAATGC  
1140

TTATGTAAAC CAAGATATAA AAGAGTGCTG ATTTTTTGAG TAACTTGCA ACAGTCCTAA  
1200

CATTCACCTC TTGTGTGTTT GTGTCTGTC GCCATCCCGT CTCCGCTCGT CACTTATCCT  
1260

TCACTTTCCA GAGGGTCCCC CCGCAGACCC CGGCGACCCT CAGGTCGGCC GACTGCGGCA  
1320

GCTGGCGCCC GAACAGGGAC CCTCGGATAA GTGACCCTTG TCTCTATTTT TACTATTTGG  
1380

TGTTTGTCTT GTATTGTCTC TTTCTTGTCT GGCTATCATC ACAAGAGCGG AACGGACTCA  
1440

CCATAGGGAC CAAGCTAGCG ACTGAAAATG AGACATATTA TCTGCCACGG AGGTGTTATT  
1500

ACCGAAGAAA TGGCCGCCAG TCTTTTGGAC CAGCTGATCG AAGAGGTACT GGCTGATAAT  
1560

CTTCCACCTC CTAGCCATTT TGAACCACCT ACCCTTCACG AACTGTATGA TTTAGACGTG  
1620

ACGGCCCCCG AAGATCCCAA CGAGGAGGCG GTTTCGCAGA TTTTCCCGA CTCTGTAATG  
1680

TTGGCGGTGC AGGAAGGGAT TGACTIONCT ACTTTTCCGC CGGCGCCCGG TTCTCCGGAG  
1740

CCGCCTCACC TTTCCCGGCA GCCCAGAGCAG CCGGAGCAGA GAGCCTTGGG TCCGGTTTCT  
1800

ATGCCAAACC TTGTACCGGA GGTGATCGAT CTTACCTGCC ACGAGGCTGG CTTTCCACCC  
1860

AGTGACGACG AGGATGAAGA GGGTGAGGAG TTTGTGTTAG ATTATGTGA GCACCCCGGG  
1920

CACGGTTGCA GGTCTTGTCA TTATCACCGG AGGAATACGG GGGACCCAGA TATTATGTGT  
1980

TCGCTTTGCT ATATGAGGAC CTGTGGCATG TTTGTCTACA GTAAGTGAAA ATTATGGGCA  
2040

GTGGGTGATA GAGTGGTGGG TTTGGTGTGG TAATTTTTTT TTTAATTTTT ACAGTTTTGT  
2100

GGTTTAAAGA ATTTTGTATT GTGATTTTTT TAAAAGGTCC TGTGTCTGAA CCTGAGCCTG  
2160

AGCCCGAGCC AGAACCGGAG CCTGCAAGAC CTACCCGCCG TCCTAAAATG GCGCCTGCTA  
2220

TCCTGAGACG CCCGACATCA CCTGTGTCTA GAGAATGCAA TAGTAGTACG GATAGCTGTG  
2280

ACTCCGGTCC TTCTAACACA CCTCCTGAGA TACACCCGGT GGTCCTGCTG TGCCCCATTA  
2340

AACCAGTTGC CGTGAGAGTT GGTGGGCGTC GCCAGGCTGT GGAATGTATC GAGGACTTGC  
2400

TTAACGAGCC TGGGCAACCT TTGGACTTGA GCTGTAAACG CCCCAGGCCA TAAGGTGTAA  
2460

ACCTGTGATT GCGTGTGTGG TTAACGCCTT TGTTTGCTGA ATGAGTTGAT GTAAGTTTAA  
2520

TAAAGGGTGA GATAATGTTT AACTTGCATG GCGTGTTAAA TGGGGCGGGG CTTAAAGGGT  
2580

ATATAATGCG CCGTGGGCTA ATCTTGCTTA CATCTGACCT CATGGAGGCT TGGGAGTGTT  
2640

TGGAAGATTT TTCTGCTGTG CGTAACTTGC TGGAACAGAG CTCTAACAGT ACCTCTTGTT  
2700

TTTGGAGGTT TCTGTGGGGC TCATCCCAGG CAAAGTTAGT CTGCAGAATT AAGGAGGATT  
2760

ACAAGTGGGA ATTTGAAGAG CTTTTGAAAT CCTGTGGTGA GCTGTTTGAT TCTTTGAATC  
2820

TGGGTCACCA GCGGCTTTTC CAAGAGAAGG TCATCAAGAC TTTGGATTTT TCCACACCGG  
2880

GGCGCGCTGC GGCTGCTGTT GCTTTTTTGA GTTTTATAAA GGATAAATGG AGCGAAGAAA  
2940

CCCATCTGAG CGGGGGGTAC CTGCTGGATT TTCTGGCCAT GCATCTGTGG AGAGCGGTTG  
3000

TGAGACACAA GAATCGCCTG CTACTGTTGT CTTCCGTCCG CCCGGCGATA ATACCGACGG  
3060

AGGAGCAGCA GCAGCAGCAG GAGGAAGCCA GGCGGCGGCG GCAGGAGCAG AGCCCATGGA  
3120

ACCCGAGAGC CGGCCTGGAC CCTCGGGAAT GAATGTTGTA CAGGTGGCTG AACTGTATCC  
3180

AGAACTGAGA CGCATTTTGA CAATTACAGA GGATGGGCAG GGGCTAAAGG GGGTAAAGAG  
3240

GGAGCGGGGG GCTTGTGAGG CTACAGAGGA GGCTAGGAAT CTAGCTTTTA GCTTAATGAC  
3300

CAGACACCGT CCTGAGTGTA TTACTTTTCA ACAGATCAAG GATAATTGCG CTAATGAGCT  
3360

TGATCTGCTG GCGCAGAAGT ATTCCATAGA GCAGCTGACC ACTTACTGGC TGCAGCCAGG  
3420

GGATGATTTT GAGGAGGCTA TTAGGGTATA TGCAAAGGTG GCACTTAGGC CAGATTGCAA  
3480

GTACAAGATC AGCAAACCTG TAAATATCAG GAATTGTTGC TACATTTCTG GGAACGGGGC  
3540

CGAGGTGGAG ATAGATACGG AGGATAGGGT GGCCTTTAGA TGTCAGCATGA TAAATATGTG  
3600

GCCGGGGGTG CTTGGCATGG ACGGGGTGGT TATTATGAAT GTAAGGTTTA CTGGCCCCAA  
3660

TTTLAGCGGT ACGGTTTTCC TGGCCAATAC CAACCTTATC CTACACGGTG TAAGCTTCTA  
3720

TGGGTTTAAC AATACCTGTG TGGAAGCCTG GACCGATGTA AGGGTTCGGG GCTGTGCCTT  
3780

TTACTGCTGC TGGAAGGGGG TGGTGTGTCG CCCCAAAAGC AGGGCTTCAA TTAAGAAATG  
3840

CCTCTTTGAA AGGTGTACCT TGGGTATCCT GTCTGAGGGT AACTCCAGGG TGCGCCACAA  
3900

TGTGGCCTCC GACTGTGGTT GCTTCATGCT AGTGAAAAGC GTGGCTGTGA TTAAGCATAA  
3960

CATGGTATGT GGCAACTGCG AGGACAGGGC CTCTCAGATG CTGACCTGCT CGGACGGCAA  
4020

CTGTCACCTG CTGAAGACCA TTCACGTAGC CAGCCACTCT CGCAAGGCCT GGCCAGTGTT  
4080

TGAGCATAAC ATACTGACCC GCTGTTCCCTT GCATTTGGGT AACAGGAGGG GGGTGTTCCCT  
4140

ACCTTACCAA TGCAATTGA GTCACACTAA GATATTGCTT GAGCCCGAGA GCATGTCCAA  
4200

GGTGAACCTG AACGGGGTGT TTGACATGAC CATGAAGATC TGAAGGTGC TGAGGTACGA  
4260

TGAGACCCGC ACCAGGTGCA GACCCTGCGA GTGTGGCGGT AACATATTA GGAACCAGCC  
4320

TGTGATGCTG GATGTGACCG AGGAGCTGAG GCCCGATCAC TTGGTGCTGG CCTGCACCCG  
4380

CGCTGAGTTT GGCTCTAGCG ATGAAGATAC AGATTGAGGT ACTGAAATGT GTGGGCGTGG  
4440

CTTAAGGGTG GGAAAGAATA TATAAGGTGG GGGTCTTATG TAGTTTTGTA TCTGTTTTGC  
4500

AGCAGCCGCC GCCGCCATGA GCACCAACTC GTTTGATGGA AGCATTGTGA GCTCATATTT  
4560

GACAACGCGC ATGCCCCCAT GGGCCGGGGT GCGTCAGAAT GTGATGGGCT CCAGCATTGA  
4620

TGGTCGCCCC GTCCTGCCCC CAAACTCTAC TACCTTGACC TACGAGACCG TGTCTGGAAC  
4680

GCCGTTGGAG ACTGCAGCCT CCGCCGCCGC TTCAGCCGCT GCAGCCACCG CCCGCGGGAT  
4740

TGTGACTGAC TTTGCTTTCC TGAGCCCGCT TGCAAGCAGT GCAGCTTCCC GTTCATCCGC  
4800

CCGCGATGAC AAGTTGACGG CTCTTTTGGC ACAATTGGAT TCTTTGACCC GGGA ACTTAA  
4860

TGTCGTTTCT CAGCAGCTGT TGGATCTGCG CCAGCAGGTT TCTGCCCTGA AGGCTTCCTC  
4920

CCCTCCCAAT GCGGTTTAAA ACATAAATAA AAAACCAGAC TCTGTTTGA TTTGGATCAA  
4980

GCAAGTGTCT TGCTGTCTCT CGAGGGATCT TTGTGAAGGA ACCTTACTTC TGTGGTGTGA  
5040

CATAATTGGA CAAACTACCT ACAGAGATTT AAAGCTCTAA GGTAATATA AAATTTTAA  
5100

GTGTATAATG TGTTAACTA CTGATTCTAA TTGTTTGTGT ATTTTAGATT CCAACCTATG  
5160

GAAGTATGA ATGGGAGCAG TGGTGAATG CCTTTAATGA GGAAAACCTG TTTTGCTCAG  
5220

AAGAAATGCC ATCTAGTGAT GATGAGGCTA CTGCTGACTC TCAACATTCT ACTCCTCCAA  
5280

AAAAGAAGAG AAAGGTAGAA GACCCCAAGG ACTTTCCTTC AGAATTGCTA AGTTTTTTGA  
5340

GTCATGCTGT GTTTAGTAAT AGAACTCTTG CTTGCTTTGC TATTACACC ACAAAGGAAA  
5400

AAGCTGCACT GCTATACAAG AAAATTATGG AAAAATATTC TGTAACCTTT ATAAGTAGGC  
5460

ATAACAGTTA TAATCATAAC ATACTGTTTT TTCTTACTCC ACACAGGCAT AGAGTGTCTG  
5520

CTATTAATAA CTATGCTCAA AAATTGTGTA CCTTTAGCTT TTTAATTTGT AAAGGGGTTA  
5580

ATAAGGAATA TTTGATGTAT AGTGCCTTGA CTAGAGATCA TAATCAGCCA TACCACATTT  
5640

GTAGAGGTTT TACTTGCTTT AAAAAACCTC CCACACCTCC CCCTGAACCT GAAACATAAA  
5700

ATGAATGCAA TTGTTGTTGT TAACTTGTTT ATTGCAGCTT ATAATGGTTA CAAATAAAGC  
5760

AATAGCATCA CAAATTTTAC AAATAAAGCA TTTTTTTCAC TGCATTCTAG TTGTGGTTTG  
5820

TCCAAACTCA TCAATGTATC TTATCATGTC TGGATCCGGC TGTGGAATGT GTGTCAGTTA  
5880

GGGTGTGGAA AGTCCCCAGG CTCCCCAGCA GGCAGAAGTA TGCAAAGCAT GCATCTCAAT  
5940

TAGTCAGCAA CCAGGTGTGG AAAGTCCCCA GGCTCCCCAG CAGGCAGAAG TATGCAAAGC  
6000

ATGCATCTCA ATTAGTCAGC AACCATAGTC CCGCCCCCTAA CTCCGCCCAT CCGCCCCCTA  
6060

ACTCCGCCCA GTTCCGCCCA TTCTCCGCCC CATGGCTGAC TAATTTTTTT TATTTATGCA  
6120

GAGGCCGAGG CCGCCTCGGC CTCTGAGCTA TTCCAGAAGT AGTGAGGAGG CTTTTTTGGA  
6180

GGCCTAGGCT TTTGCAAAAA GCTTGGACAC AAGACAGGCT TGCGAGATAT GTTTGAGAAT  
6240

ACCACTTTAT CCCGCGTCAG GGAGAGGCAG TCGGTAAAAA GACGCGGACT CATGTGAAAT  
6300

ACTGGTTTTT AGTGCGCCAG ATCTCTATAA TCTCGCGCAA CCTATTTTCC CCTCGAACAC  
6360

TTTTTAAGCC GTAGATAAAC AGGCTGGGAC ACTTCACATG AGCGAAAAAT ACATCGTCAC  
6420

CTGGGACATG TTGCAGATCC ATGCACGTAA ACTCGCAAGC CGACTGATGC CTTCTGAACA  
6480

ATGGAAAGGC ATTATTGCCG TAAGCCGTGG CGGTCTGGTA CCGGGTGCGT TACTGGCGCG  
6540

TGAACTGGGT ATTCGTCATG TCGATACCGT TTGTATTTCC AGCTACGATC ACGACAACCA  
6600

GCGCGAGCTT AAAGTGCTGA AACGCGCAGA AGGCGATGGC GAAGGCTTCA TCGTTATTGA  
6660

TGACCTGGTG GATACCGGTG GTACTGCGGT TCGGATTCGT GAAATGTATC CAAAAGCGCA  
6720

CTTTGTCACC ATCTTCGCAA AACC GGCTGG TCGTCCGCTG GTTGATGACT ATGTTGTTGA  
6780

TATCCCGCAA GATACCTGGA TTGAACAGCC GTGGGATATG GCGTCGTAT TCGTCCCGCC  
6840

AATCTCCGGT CGCTAATCTT TTCAACGCCT GGCAGTCCG GCGTTGTTC TTTTAACTT  
6900

CAGGCGGGTT ACAATAGTTT CCAGTAAGTA TTCTGGAGGC TGCATCCATG ACACAGGCAA  
6960

ACCTGAGCGA AACCTGTTC AAACCCCGCT TTAAACATCC TGAAACCTCG ACGCTAGTCC  
7020

GCCGCTTTAA TCACGGCGCA CAACCGCCTG TGCAGTCGGC CCTTGATGGT AAAACCATCC  
7080

CTCACTGGTA TCGCATGATT AACCGTCTGA TGTGGATCTG GCGCGGCATT GACCCACGCG  
7140



AAATCCTCGA CGTCCAGGCA CGTATTGTGA TGAGCGATGC CGAACGTACC GACGATGATT  
7200

TATACGATAC GGTGATTGGC TACCGTGGCG GCAACTGGAT TTATGAGTGG GCCCCGGATC  
7260

TTTGTGAAGG AACCTTACTT CTGTGGTGTG ACATAATTGG ACAAACCTACC TACAGAGATT  
7320

TAAAGCTCTA AGGTAAATAT AAAATTTTAA AGTGTATAAT GTGTAAACT ACTGATTCTA  
7380

ATTGTTTGTG TATTTTAGAT TCCAACCTAT GGAACGTATG AATGGGAGCA GTGGTGAAT  
7440

GCCTTTAATG AGGAAAACCT GTTTGCTCA GAAGAAATGC CATCTAGTGA TGATGAGGCT  
7500

ACTGCTGACT, CTCAACATTC TACTCCTCCA AAAAAGAAGA GAAAGGTAGA AGACCCCAAG  
7560

GACTTTCCTT CAGAATTGCT AAGTTTTTTG AGTCATGCTG TGTTTAGTAA TAGAACTCTT  
7620

GCTTGCTTTG CTATTTACAC CACAAAGGAA AAAGCTGCAC TGCTATACAA GAAAATTATG  
7680

GAAAAATATT CTGTAACCTT TATAAGTAGG CATAACAGTT ATAATCATAA CATACTGTTT  
7740

TTTCTTACTC CACACAGGCA TAGAGTGTCT GCTATTAATA ACTATGCTCA AAAATTGTGT  
7800

ACCTTTAGCT TTTTAATTTG TAAAGGGGTT AATAAGGAAT ATTTGATGTA TAGTGCCTTG  
7860

ACTAGAGATC ATAATCAGCC ATACCACATT TGTAGAGGTT TTAATTGCTT TAAAAAACCT  
7920

CCCACACCTC CCCCTGAACC TGAAACATAA AATGAATGCA ATTGTTGTTG TTAACCTGTT  
7980

TATTGCAGCT TATAATGGTT ACAAATAAAG CAATAGCATC ACAAATTTCA CAAATAAAGC  
8040

ATTTTTTTCA CTGCATTCTA GTTGTGGTTT GTCCAAACTC ATCAATGTAT CTTATCATGT  
8100

CTGGATCCCC AGGAAGCTCC TCTGTGTCCT CATAAACCCCT AACCTCCTCT ACTTGAGAGG  
8160

ACATTCCAAT CATAGGCTGC CCATCCACCC TCTGTGTCCT CCTGTTAATT AGGTCACTTA  
8220

ACAAAAAGGA AATTGGGTAG GGGTTTTTCA CAGACCGCTT TCTAAGGGTA ATTTTAAAT  
8280

ATCTGGGAAG TCCCTTCCAC TGCTGTGTTT CAGAAGTGTT GGTAACAGC CCACAAATGT  
8340

CAACAGCAGA AACATACAAG CTGTCAGCTT TGCACAAGGG CCCAACACCC TGCTCATCAA  
8400

GAAGCACTGT GGTGCTGTG TTAGTAATGT GCAAAACAGG AGGCACATTT TCCCCACCTG  
8460

TGTAGGTTCC AAAATATCTA GTGTTTTTCAT TTTTACTTGG ATCAGGAACC CAGCACTCCA  
8520

CTGGATAAGC ATTATCCTTA TCCAAAACAG CCTTGTGGTC AGTGTTTCATC TGCTGACTGT  
8580

CAACTGTAGC ATTTTTTGGG GTTACAGTTT GAGCAGGATA TTTGGTCCTG TAGTTTGCTA  
8640

ACACACCCTG CAGCTCCAAA GGTCCCCAC CAACAGCAAA AAAATGAAAA TTTGACCCTT  
8700

GAATGGGTTT TCCAGCACCA TTTTCATGAG TTTTTGTGT CCCTGAATGC AAGTTTAACA  
8760

TAGCAGTTAC CCCAATAACC TCAGTTTTAA CAGTAACAGC TTCCCACATC AAAATATTTTC  
8820

CACAGGTAA GTCCTCATTT AAATTAGGCA AAGGAATTCT TGAAGACGAA AGGGCCTCGT  
8880

GATACGCTA TTTTATAGG TTAATGTCAT GATAATAATG GTTCTTAGA CGTCAGGTGG  
8940

CACTTTTCGG GGAAATGTGC GCGGAACCCC TATTTGTTTA TTTTCTAAA TACATTCAAA  
9000

TATGTATCCG CTCATGAGAC AATAACCCTG ATAAATGCTT CAATAATATT GAAAAAGGAA  
9060

GAGTATGAGT ATTCAACATT TCCGTGTCGC CCTTATCCC TTTTTGCGG CATTTTGCCT  
9120

TCCTGTTTTT GCTCACCCAG AAACGCTGGT GAAAGTAAAA GATGCTGAAG ATCAGTTGGG  
9180

TGCACGAGTG GGTTACATCG AACTGGATCT CAACAGCGGT AAGATCCTTG AGAGTTTTCG  
9240

CCCCGAAGAA CGTTTTCCAA TGATGAGCAC TTTTAAAGTT CTGCTATGTG GCGCGGTATT  
9300

ATCCCGTGTT GACGCCGGGC AAGAGCAACT CGGTCGCCGC ATACACTATT CTCAGAATGA  
9360

CTTGGTTGAG TACTCACCAG TCACAGAAAA GCATCTTACG GATGGCATGA CAGTAAGAGA  
9420

ATTATGCAGT GCTGCCATAA CCATGAGTGA TAACACTGCG GCCAACTTAC TTCTGACAAC  
9480

GATCGGAGGA CCGAAGGAGC TAACCGCTTT TTTGCACAAC ATGGGGGATC ATGTAAGTCG  
9540

CCTTGATCGT TGGGAACCGG AGCTGAATGA AGCCATACCA AACGACGAGC GTGACACCAC  
9600

GATGCCTGCA GCAATGGCAA CAACGTTGCG CAACTATTA ACTGGCGAAC TACTTACTCT  
9660

AGCTTCCCGG CAACAATTAA TAGACTGGAT GGAGGCGGAT AAAGTTGCAG GACCACTTCT  
9720

GCGCTCGGCC CTTCCGGCTG GCTGGTTTAT TGCTGATAAA TCTGGAGCCG GTGAGCGTGG  
9780

GTCTCGCGGT ATCATTGCAG CACTGGGGCC AGATGGTAAG CCCTCCCGTA TCGTAGTTAT  
9840

CTACACGACG GGGAGTCAGG CAACTATGGA TGAACGAAAT AGACAGATCG CTGAGATAGG  
9900

TGCCTCACTG ATTAAGCATT GGTAAGTGC AGACCAAGTT TACTCATATA TACTTTAGAT  
9960

TGATTTAAAA CTTTATTTTT AATTTAAAAG GATCTAGGTG AAGATCCTTT TTGATAATCT  
10020

CATGACCAAA ATCCCTTAAC GTGAGTTTTT GTTCCACTGA GCGTCAGACC CCGTAGAAAA  
10080

GATCAAAGGA TCTTCTTGAG ATCCTTTTTT TCTGCGCGTA ATCTGCTGCT TGCAAACAAA  
10140

AAAACCACCG CTACCAGCGG TGGTTTGTTT GCCGGATCAA GAGCTACCAA CTCTTTTTC  
10200

GAAGGTAAGT GGCTTCAGCA GAGCGCAGAT ACCAAATACT GTCCTTCTAG TGTAGCCGTA  
10260

GTTAGGCCAC CACTTCAAGA ACTCTGTAGC ACCGCCTACA TACCTCGCTC TGCTAATCCT  
10320

GTTACCAGTG GCTGCTGCCA GTGGCGATAA GTCGTGTCTT ACCGGGTGG ACTCAAGACG  
10380

ATAGTTACCG GATAAGGCGC AGCGGTCGGG CTGAACGGGG GGTTCGTGCA CACAGCCCAG  
10440

CTTGAGCGA ACGACCTACA CCGAACTGAG ATACCTACAG CGTGAGCTAT GAGAAAGCGC  
10500

CACGCTTCCC GAAGGGAGAA AGGCGGACAG GTATCCGGTA AGCGGCAGGG TCGGAACAGG  
10560

AGAGCGCACG AGGGAGCTTC CAGGGGGAAA CGCCTGGTAT CTTTATAGTC CTGTCGGGTT  
10620

TCGCCACCTC TGAATTGAGC GTCGATTTTT GTGATGCTCG TCAGGGGGGC GGAGCCTATG  
10680

GAAAAACGCC AGCAACGCGG CCTTTTTACG GTTCCTGGCC TTTTGCTGGC CTTTTGCTCA  
10740

CATGTTCTTT CCTGCGTTAT CCCCTGATTC TGTGGATAAC CGTATTACCG CCTTTGAGTG  
10800

AGCTGATACC GCTCGCCGCA GCCGAACGAC CGAGCGCAGC GAGTCAGTGA GCGAGGAAGC  
10860

GGAAGAGCGC CTGATGCGGT ATTTTCTCCT TACGCATCTG TGCGGTATTT CACACCGCAT  
10920

ACCGCCTCAG AAGCCATAGA GCCCACC GCA TCCCCAGCAT GCCTGCTATT GTCTTCCCAA  
10980

TCCTCCCCCT TGCTGTCCTG CCCCACCCCA CCCCCAGAA TAGAATGACA CCTACTCAGA  
11040

CAATGCGATG CAATTCCTC ATTTTATTAG GAAAGGACAG TGGGAGTGGC ACCTTCCAGG  
11100

GTCAAGGAAG GCACGGGGGA GGGGCAAACA ACAGATGGCT GGCAACTAGA AGGCACAGTC  
11160

GAGGCTGATC AGCGAGCTCT AGCATTTAGG TGACACTATA GAATAGGGCC CTCTAGATGC  
11220

ATGCTCGAGC GGCCGCTTCT TTATTCTTGG GCAATGTATG AAAAAGTGTA AGAGGATGTG  
11280

GCAAATATTT CATTAATGTA GTTGTGGCCA GACCAGTCCC ATGAAAATGA CATAGAGTAT  
11340

GCACTTGGAG TTGTGTCTCC TGTTTCCTGT GTACCGTTTA GTGTAATGGT TAGTGTTACA  
11400

GGTTTAGTTT TGTCTCCGTT TAAGTAACT TGACTGACAA TGTTACTTTT GGCAGTTTTA  
11460

CCGTGAGATT TTGGATAAGC TGATAGGTTA GGCATAAATC CAACAGCGTT TGTATAGGCT  
11520

GTGCCTTCAG TAAGATCTCC ATTTCTAAAG TTCCAATATT CTGGGTCCAG GAAGGAATTG  
11580

TTTAGTAGCA CTCCATTTTC GTCAAATCTT ATAATAAGAT GAGCACTTTG AACTGTTCCA  
11640

GATATTGGAG CCAAACCTGCC TTAAACAGCC AAAACTGAAA CTGTAGCAAG TATTTGACTG  
11700

CCACATTTTG TTAAGACCAA AGTGAGTTTA GCATCTTTCT CTGCATTTAG TCTACAGTTA  
11760

GGAGATGGAG CTGGTGTGGT CCACAAAGTT AGCTTATCAT TATTTTTGTT TCCTACTGTA  
11820

ATGGCACCTG TGCTGTCAAA ACTAAGGCCA GTTCCTAGTT TAGGAACCAT AGCCTTGTTT  
11880

GAATCAAATT CTAGGCCATG GCCAATTTTT GTTTGAGGG GATTGTGTT TGGTGCATTA  
11940

GGTGAACCAA ATTCAAGCCC ATCTCCTGCA TTAATGGCTA TGGCTGTAGC GTCAAACATC  
12000

AACCCCTTGG CAGTGCTTAG GTTAACCTCA AGCTTTTTGG AATTGTTTGA AGCTGTAAAC  
12060

AAGTAAAGGC CTTTGTGTA GTTAATATCC AAGTTGTGGG CTGAGTTTAT AAAAAGAGGG  
12120

CCCTGTCCTA GTCTTAGATT TAGTTGGTTT TGAGCATCAA ACGGATAACT AACATCAAGT  
12180

ATAAGGCGTC TGTTTTGAGA ATCAATCCTT AGTCCTCCTG CTACATTAAG TTGCATATTG  
12240

CCTTGTGAAT CAAAACCCAA GGCTCCAGTA ACTTTAGTTT GCAAGGAAGT ATTATTAATA  
12300

GTCACACCTG GACCAGTTGC TACGGTCAAA GTGTTTAGGT CGTCTGTTAC ATGCAAAGGA  
12360

GCCCCGTACT TTAGTCCTAG TTTTCCATTT TGTGTATAAA TGGGCTCTTT CAAGTCAATG  
12420

CCCAAGCTAC CAGTGGCAGT AGTTAGAGGG GGTGAGGCAG TGATAGTAAG GGTACTGCTA  
12480

TCGGTGGTGG TGAGGGGGCC TGATGTTTGC AGGGCTAGCT TTCCTTCTGA CACTGTGAGG  
12540

GGTCCTTGGG TGGCAATGCT AAGTTTGGAG TCGTGCACGG TTAGCGGGGC CTGTGATTGC  
12600

ATGGTGAGTG TGTGCCCCG GACCATTAGA GGTGCGGCGG CAGCCACAGT TAGGGCTTCT  
12660

GAGGTAAGTG TGAGGGGTGC AGATATTTCC AGGTTTATGT TTGACTTGGT TTTTGTGAGA  
12720

GGTGGGCTCA CAGTGGTTAC ATTTTGGGAG GTAAGGTTGC CGGCCTCGTC CAGAGAGAGG  
12780

CCGTTGCCCA TTTTGAGCGC AAGCATGCCA TTGGAGGTAA CTAGAGGTTT GGATAGGCGC  
12840

AAAGAGAGTA CCCCAGGGGG ACTCTCTTGA AACCCATTGG GGGATACAAA GGGAGGAGTA  
12900

AGAAAAGGCA CAGTTGGAGG ACCGGTTTCC GTGTCATATG GATACACGGG GTTGAAGGTA  
12960

TCTTCAGACG GTCTTGCGCG CTTTCATCTTG GATCTCAAGC CTGCCACACC TCACCTCGAC  
13020

CATCCGCCGT CTCAAGACCG CCTACTTTAA TTACATCATC AGCAGCACCT CCGCCAGAAA  
13080

CAACCCCGAC CGCCACCCGC TGCCGCCCGC CACGGTGCTC AGCCTACCTT GCGACTGTGA  
13140

CTGGTTAGAC GCCTTTCTCG AGAGGTTTTT CGATCCGGTC GATGCGGACT CGCTCAGGTC  
13200

CCTCGGTGGC GGAGTACCGT TCGGAGGCCG ACGGGTTTCC GATCCAAGAG TACTGGAAAG  
13260

ACCGCGAAGA GTTTGTCCTC AACCGCGAGC CCAACAGCGA GCTCGAATTC AGATCCGAGC  
13320

TCGGTACCAA GCTTGGGTCT CCCTATAGTG AGTCGTATTA ATTTGATAA GCCAGTAAGC  
13380



AGTGGGTTCT CTAGTTAGCC AGAGAGCTCT GCTTATATAG ACCTCCCACC GTACACGCCT  
13440

ACCGCCCAT TGCCTCAATG GGGCGGAGTT GTTACGACAT TTTGGAAAGT CCCGTTGATT  
13500

TTGGTGCCAA AACAACTCC CATTGACGTC AATGGGGTGG AGACTTGGA AATCCCCGTGA  
13560

GTCAAACCGC TATCCACGCC CATTGATGTA CTGCCAAAAC CGCATCACCA TGGTAATAGC  
13620

GATGACTAAT ACGTAGATGT ACTGCCAAGT AGGAAAGTCC CATAAGGTCA TGTACTGGGC  
13680

ATAATGCCAG GCGGGCCATT TACCGTCATT GACGTCAATA GGGGGCGTAC TTGGCATATG  
13740

ATACACTTGA TGTACTGCCA AGTGGGCAGT TTACCGTAAA TAGTCCACCC ATTGACGTCA  
13800

ATGGAAAGTC CCTATTGGCG TTACTATGGG AACATACGTC ATTATTGACG TCAATGGGCG  
13860

GGGGTCGTTG GGCGGTCAGC CAGGCGGGCC ATTTACCGTA AGTTATGTAA CGCGGAACTC  
13920

CATATATGGG CTATGAACTA ATGACCCCGT AATTGATTAC TATTAATAAC TAGTCAATAA  
13980

TCAATGTCAA CGCGTATATC TGGCCCGTAC ATCGCGAAGC AGCGCAAAAC GCCTAACCCCT  
14040

AAGCAGATTC TTCATGCAAT TGTCGGTCAA GCCTTGCCTT GTTGTAGCTT AAATTTTGCT  
14100

CGCGCACTAC TCAGCGACCT CCAACACACA AGCAGGGAGC AGATACTGGC TTAACATATGC  
14160

- 131 -

GGCATCAGAG CAGATTGTAC TGAGAGTCGA CCATAGGGGA TCGGGAGATC TCCCGATCCG  
14220

TCTATGGTGC ACTCTCAGTA CAATCTGCTC TGATGCCGCA TAGTTAAGCC AGTATACACT  
14280

CCGCTATCGC TACGTGACTG GGTCATGGCT GCGCCCCGAC ACCCGCCAAC ACCCGCTGAC  
14340

GCGCCCTGAC GGGCTTGTCT GCTCCCGGCA TCCGCTTACA GACAAGCTGT GACCGTCTCC  
14400

GGGAGCTGCA TGTGTCAGAG GTTTTCACCG TCATACCGA AACGCGCGAG GCAGC  
14455

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10610 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: circular

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

GACGGATCGG GAGATCCGCG CGGTACACAG AATTCAGGAG ACACAACTCC AAGTGCATAC  
60

TCTATGTCAT TTTCATGGGA CTGGTCTGGC CACAACTACA TTAATGAAAT ATTTGCCACA  
120

TCCTCTTACA CTTTTTCATA CATTGCCCAA GAATAAAGAA TCGTTTGTGT TATGTTTCAA  
180

CGTGTTTATT TTTCAATTGC AGAAAATTTC AAGTCATTTT TCATTCAGTA GTATAGCCCC  
240

ACCACCACAT AGCTTATACA GATCACCGTA CCTTAATCAA ACTCACAGAA CCCTAGTATT  
300

CAACCTGCCA CCTCCCTCCC AACACACAGA GTACACAGTC CTTTCTCCCC GGCTGGCCTT  
360

AAAAAGCATC ATATCATGGG TAACAGACAT ATTCTTAGGT GTTATATTCC ACACGGTTTC  
420

CTGTCGAGCC AAACGCTCAT CAGTGATATT AATAAACTCC CCGGGCAGCT CACTTAAGTT  
480

CATGTCGCTG TCCAGCTGCT GAGCCACAGG CTGCTGTCCA ACTTGCGGTT GCTTAACGGG  
540

CGGCGAAGGA GAAGTCCACG CCTACATGGG GGTAGAGTCA TAATCGTGCA TCAGGATAGG  
600

GCGGTGGTGC TGCAGCAGCG CGCGAATAAA CTGCTGCCGC CGCCGCTCCG TCCTGCAGGA  
660

ATACAACATG GCAGTGGTCT CCTCAGCGAT GATTCGCACC GCCCCGAGCA TAAGGCGCCT  
720

TGTCCTCCGG GCACAGCAGC GCACCCTGAT CTCACTTAAA TCAGCACAGT AACTGCAGCA  
780

CAGCACCACA ATATTGTTCA AAATCCCACA GTGCAAGGCG CTGTATCCAA AGCTCATGGC  
840

GGGGACCACA GAACCCACGT GGCCATCATA CCACAAGCGC AGGTAGATTA AGTGCGGACC  
900

CCTCATAAAC ACGCTGGACA TAAACATTAC CTCTTTTGGC ATGTTGTAAT TCACCACCTC  
960

CCGGTACCAT ATAAACCTCT GATTAAACAT GGCGCCATCC ACCACCATCC TAAACCAGCT  
1020

GGCCAAAACC TGCCCGCCGG CTATACACTG CAGGGAACCG GGACTGGAAC AATGACAGTG  
1080

GAGAGCCCAG GACTCGTAAC CATGGATCAT CATGCTCGTC ATGATATCAA TGTGGCACA  
1140

ACACAGGCAC ACGTGCATAC ACTTCCTCAG GATTACAAGC TCCTCCCGCG TTAGAACCAT  
1200

ATCCCAGGGA ACAACCCATT CCTGAATCAG CGTAAATCCC AACTGCAGG GAAGACCTCG  
1260

CACGTAATC ACGTTGTGCA TTGTCAAAGT GTTACATTCG GGCAGCAGCG GATGATCCTC  
1320

CAGTATGGTA GCGCGGGTTT CTGTCTCAA AGGAGGTAGA CGATCCCTAC TGTACGGAGT  
1380

GCGCCGAGAC AACCGAGATC GTGTTGGTCG TAGTGTCATG CCAAATGGAA CGCCGGACGT  
1440

AGTCATATTT CCTGAAGCAA AACCAGGTGC GGGCGTGACA AACAGATCTG CGTCTCCGGT  
1500

CTCGCCGCTT AGATCGCTCT GTGTAGTAGT TGTAGTATAT CCACTCTCTC AAAGCATCCA  
1560

GGCGCCCCCT GGCTTCGGGT TCTATGTAAA CTCCTTCATG CGCCGCTGCC CTGATAACAT  
1620

CCACCACCGC AGAATAAGCC ACACCCAGCC AACCTACACA TTCGTTCTGC GAGTCACACA  
1680

CGGGAGGAGC GGGAAGAGCT GGAAGAACCA TGTTTTTTTT TTTATTCCAA AAGATTATCC  
1740

AAAACCTCAA AATGAAGATC TATTAAGTGA ACGCGCTCCC CTCCGGTGGC GTGGTCAAAC  
1800

TCTACAGCCA AAGAACAGAT AATGGCATTG GTAAGATGTT GCACAATGGC TTCCAAAAGG  
1860

CAAACGGCCC TCACGTCCAA GTGGACGTAA AGGCTAAACC CTCAGGGTG AATCTCCTCT  
1920

ATAAACATTC CAGCACCTTC AACCATGCCC AAATAATTCT CATCTCGCCA CTTTCTCAAT  
1980

ATATCTCTAA GCAAATCCCG AATATTAAGT CCGGCCATTG TAAAAATCTG CTCCAGAGCG  
2040

CCCTCCACCT TCAGCCTCAA GCAGCGAATC ATGATTGCAA AAATTCAGGT TCCTCACAGA  
2100

CCTGTATAAG ATTCAAAGC GGAACATTAA CAAAAATACC GCGATCCCGT AGGTCCCTTC  
2160

GCAGGGCCAG CTGAACATAA TCGTGCAGGT CTGCACGGAC CAGCGCGGCC ACTTCCCCGC  
2220

CAGGAACCTT GACAAAAGAA CCCACACTGA TTATGACACG CATACTCGGA GCTATGCTAA  
2280

CCAGCGTAGC CCCGATGTAA GCTTTGTTGC ATGGGCGGCG ATATAAAATG CAAGGTGCTG  
2340

CTCAAAAAAT CAGGCAAAGC CTCGCGCAAA AAAGAAAGCA CATCGTAGTC ATGCTCATGC  
2400

AGATAAAGGC AGGTAAGCTC CGGAACCACC ACAGAAAAAG ACACCATTTT TCTCTCAAAC  
2460

ATGTCTGCGG GTTCTGTCAT AAACACAAAA TAAAATAACA AAAAAACATT TAAACATTAG  
2520

AAGCCTGTCT TACAACAGGA AAAACAACCC TTATAAGCAT AAGACGGACT ACGGCCATGC  
2580

CGGCGTGACC GTAAAAAAC TGGTCACCGT GATTAAAAAG CACCACCGAC AGCTCCTCGG  
2640

TCATGTCCGG AGTCATAATG TAAGACTCGG TAAACACATC AGGTTGATTC ATCGGTCAGT  
2700

GCTAAAAAGC GACCGAAATA GCCCGGGGGA ATACATACCC GCAGGCGTAG AGACAACATT  
2760

ACAGCCCCCA TAGGAGGTAT AACAAAATTA ATAGGAGAGA AAAACACATA AACACCTGAA  
2820

AAACCCTCCT GCCTAGGCAA AATAGCACCC TCCCGCTCCA GAACAACATA CAGCGCTTCA  
2880

CAGCGGCAGC CTAACAGTCA GCCTTACCAG TAAAAAGAA AACCTATTAA AAAAACACCA  
2940

CTCGACACGG CACCAGCTCA ATCAGTCACA GTGTAAAAAA GGGCCAAGTG CAGAGCGAGT  
3000

ATATATAGGA CTAAAAAATG ACGTAACGGT TAAAGTCCAC AAAAAACACC CAGAAAACCG  
3060

CACGCGAACC TACGCCCAGA AACGAAAGCC AAAAAACCCA CAACTTCCTC AAATCGTCAC  
3120

TTCCGTTTTT CCACGTTACG TAACTTCCCG GATCCTCTCC CGATCCCCTA TGGTCGACTC  
3180

TCAGTACAAT CTGCTCTGAT GCCGCATAGT TAAGCCAGTA TCTGCTCCCT GCTTGTGTGT  
3240

TGGAGGTCGC TGAGTAGTGC GCGAGCAAAA TTTAAGCTAC AACAAAGGCAA GGCTTGACCG  
3300

ACAATTGCAT GAAGAATCTG CTTAGGGTTA GCGGTTTTGC GCTGCTTCGC GATGTACGGG  
3360

CCAGATATAC GCGTTGACAT TGATTATTGA CTAGTTATTA ATAGTAATCA ATTACGGGGT  
3420

CATTAGTTCA TAGCCCATAT ATGGAGTTCC GCGTTACATA ACTTACGGTA AATGGCCCCG  
3480

CTGGCTGACC GCCCAACGAC CCCC GCCCAT TGACGTCAAT AATGACGTAT GTTCCCATAG  
3540

TAACGCCAAT AGGGACTTTC CATTGACGTC AATGGGTGGA CTATTTACGG TAAACTGCCC  
3600

ACTTGGCAGT ACATCAAGTG TATCATATGC CAAGTACGCC CCCTATTGAC GTCAATGACG  
3660

GTAAATGGCC CGCCTGGCAT TATGCCCAGT ACATGACCTT ATGGGACTTT CCTACTTGCC  
3720

AGTACATCTA CGTATTAGTC ATCGCTATTA CCATGGTGAT GCGGTTTTGG CAGTACATCA  
3780

ATGGGCGTGG ATAGCGGTTT GACTCACGGG GATTTCOAAG TCTCCACCCC ATTGACGTCA  
3840

ATGGGAGTTT GTTTTGGCAC CAAAATCAAC GGGACTTCC AAAATGTCGT AACAACTCCG  
3900

CCCCATTGAC GCAAATGGGC GGTAGGCGTG TACGGTGGGA GGTCTATATA AGCAGAGCTC  
3960

TCTGGCTAAC TAGAGAACCC ACTGCTTACT GGCTTATCGA AATTAATACG ACTCACTATA  
4020

GGGAGACCCA AGCTTGGTAC CGAGCTCGGA TCTGAATTCG AGCTCGCTGT TGGGCTCGCG  
4080

GTTGAGGACA AACTCTTCGC GGTCTTTCCA GTACTCTTGG ATCGGAAACC CGTCGGCCTC  
4140

CGAACGGTAC TCCGCCACCG AGGGACCTGA GCGAGTCCGC ATCGACCGGA TCGGAAAACC  
4200

TCTCGAGAAA GCGGTCTAAC CAGTCACAGT CGCAAGGTAG GCTGAGCACC GTGGCGGGCG  
4260

GCAGCGGGTG GCGGTCGGGG TTGTTTCTGG CGGAGGTGCT GCTGATGATG TAATTAAAGT  
4320

AGGCGGTCTT GAGACGGCGG ATGGTCGAGG TGAGGTGTGG CAGGCTTGAG ATCCAAGATG  
4380

AAGCGCGCAA GACCGTCTGA AGATACCTTC AACCCCGTGT ATCCATATGA CACGGAAACC  
4440

GGTCCTCCAA CTGTGCCTTT TCTTACTCCT CCCTTTGTAT CCCCCAATGG GTTTCAAGAG  
4500

AGTCCCCCTG GGGTACTCTC TTTGCGCCTA TCCGAACCTC TAGTTACCTC CAATGGCATG  
4560

CTTGCGCTCA AAATGGGCAA CGGCCTCTCT CTGGACGAGG CCGGCAACCT TACCTCCCAA  
4620

AATGTAACCA CTGTGAGCCC ACCTCTCAAA AAAACCAAGT CAAACATAAA CCTGGAAATA  
4680

TCTGCACCCC TCACAGTTAC CTCAGAAGCC CTAAGTGTGG CTGCCGCCGC ACCTCTAATG  
4740

GTCGCGGGCA ACACACTCAC CATGCAATCA CAGGCCCCGC TAACCGTGCA CGACTCCAAA  
4800



CTTAGCATTG CCACCCAAGG ACCCCTCACA GTGTCAGAAG GAAAGCTAGC CCTGCAAACA  
4860

TCAGGCCCCC TCACCACCAC CGATAGCAGT ACCCTTACTA TCACTGCCTC ACCCCCTCTA  
4920

ACTACTGCCA CTGGTAGCTT GGGCATTGAC TTGAAAGAGC CCATTTATAC ACAAATGGA  
4980

AACTAGGAC TAAAGTACGG GGCTCCTTTG CATGTAACAG ACGACCTAAA CACTTTGACC  
5040

GTAGCAACTG GTCCAGGTGT GACTATTAAT AATACTTCCT TGCAAATAA AGTTACTGGA  
5100

GCCTTGGGTT TTGATTCACA AGGCAATATG CAACTTAATG TAGCAGGAGG ACTAAGGATT  
5160

GATTCTCAA ACAGACGCCT TATACTTGAT GTTAGTTATC CGTTTGATGC TCAAAACCAA  
5220

CTAAATCTAA GACTAGGACA GGGCCCTCTT TTTATAAACT CAGCCCACAA CTTGGATATT  
5280

AACTACAACA AAGGCCTTTA CTTGTTTACA GCTTCAAACA ATTCCAAAAA GCTTGAGGTT  
5340

AACCTAAGCA CTGCCAAGGG GTTGATGTTT GACGCTACAG CCATAGCCAT TAATGCAGGA  
5400

GATGGGCTTG AATTTGGTTC ACCTAATGCA CCAAACACAA ATCCCCTCAA AACAAAAATT  
5460

GGCCATGGCC TAGAATTGA TTCAAACAAG GCTATGGTTC CTAAACTAGG AACTGGCCTT  
5520

AGTTTTGACA GCACAGGTGC CATTACAGTA GGAAACAAAA ATAATGATAA GCTAACTTTG  
5580

TGGACCACAC CAGCTCCATC TCCTAACTGT AGACTAAATG CAGAGAAAGA TGCTAAACTC  
5640

ACTTTGGTCT TAACAAAATG TGGCAGTCAA ATACTTGCTA CAGTTTCAGT TTTGGCTGTT  
5700

AAAGGCAGTT TGGCTCCAAT ATCTGGAACA GTTCAAAGTG CTCATCTTAT TATAAGATTT  
5760

GACGAAAATG GAGTGCTACT AAACAATTCC TTCCTGGACC CAGAATATTG GAACTTTAGA  
5820

AATGGAGATC TTA<sup>~</sup>CTGAAGG CACAGCCTAT ACAAACGCTG TTGGATTTAT GCCTAACCTA  
5880

TCAGCTTATC CAAAATCTCA CGGTAA<sup>~</sup>AACT GCCAAAAGTA ACATTGTCAG TCAAGTTTAC  
5940

TTAAACGGAG ACAA<sup>~</sup>AACTAA ACCTGTAACA CTAACCATTA CACTAAACGG TACACAGGAA  
6000

ACAGGAGACA CAACTCCAAG TGCATACTCT ATGTCATTTT CATGGGACTG GTCTGGCCAC  
6060

AACTACATTA ATGAAATATT TGCCACATCC TCTTACACTT TTTCATACAT TGCCCAAGAA  
6120

TAAAGAAGCG GCCGCTCGAG CATGCATCTA GAGGGCCCTA TTCTATAGTG TCACCTAAAT  
6180

GCTAGAGCTC GCTGATCAGC CTCGACTGTG CCTTCTAGTT GCCAGCCATC TGTTGTTTGC  
6240

CCCTCCCCCG TGCCTTCCTT GACCCTGGAA GGTGCCACTC CCACTGTCCT TTCCTAATAA  
6300

AATGAGGAAA TTGCATCGCA TTGTCTGAGT AGGTGTCATT CTATTCTGGG GGGTGGGGTG  
6360

GGGCAGGACA GCAAGGGGGA GGATTGGGAA GACAATAGCA GGCATGCTGG GGATGCGGTG  
6420

GGCTCTATGG CTTCTGAGGC GGAAAGAACC AGCTGGGGCT CTAGGGGGTA TCCCCACGCG  
6480

CCCTGTAGCG GCGCATTAAAG CGCGGCGGGT GTGGTGGTTA CGCGCAGCGT GACCGCTACA  
6540

CTTGCCAGCG CCCTAGCGCC CGCTCCTTTC GCTTCTTCC CTCCTTTCT CGCCACGTTC  
6600

GCCGGCTTTC CCCGTCAAGC TCTAAATCGG GGCATCCCTT TAGGGTCCG ATTTAGTGCT  
6660

TTACGGCACC TCGACCCCAA AAAACTTGAT TAGGGTGATG GTTCACGTAG TGGGCCATCG  
6720

CCCTGATAGA CGGTTTTTCG CCCTTTGACG TTGGAGTCCA CGTTCTTTAA TAGTGGACTC  
6780

TTGTTCCAAA CTGGAACAAC ACTCAACCCT ATCTCGGTCT ATTCTTTTGA TTTATAAGGG  
6840

ATTTTGGGGA TTTCGGCCTA TTGGTTAAAA AATGAGCTGA TTAAACAAAA ATTTAACGCG  
6900

AATTAATTCT GTGGAATGTG TGTCAGTTAG GGTGTGGAAA GTCCCCAGGC TCCCCAGGCA  
6960

GGCAGAAGTA TGCAAAGCAT GCATCTCAAT TAGTCAGCAA CCAGGTGTGG AAAGTCCCCA  
7020

GGCTCCCCAG CAGGCAGAAG TATGCAAAGC ATGCATCTCA ATTAGTCAGC AACCATAGTC  
7080

CCGCCCCCTAA CTCCGCCCAT CCCGCCCTA ACTCCGCCCA GTTCCGCCCA TTCTCCGCCC  
7140

CATGGCTGAC TAATTTTTTT TATTTATGCA GAGGCCGAGG CCGCCTCTGC CTCTGAGCTA  
7200

TTCCAGAAGT AGTGAGGAGG CTTTTTTGGA GGCCTAGGCT TTTGCAAAAA GCTCCCGGGA  
7260

GCTTGTATAT CCATTTTCGG ATCTGATCAA GAGACAGGAT GAGGATCGTT TCGCATGATT  
7320

GAACAAGATG GATTGCACGC AGGTTCTCCG GCCGCTTGGG TGGAGAGGCT ATTCGGCTAT  
7380

GACTGGGCAC AACAGACAAT CGGCTGCTCT GATGCCGCCG TGTTCGGCT GTCAGCGCAG  
7440

GGGCGCCCGG TTCTTTTGT CAAGACCGAC CTGTCCGGTG CCCTGAATGA ACTGCAGGAC  
7500

GAGGCAGCGC GGCTATCGTG GCTGGCCACG ACGGGCGTTC CTTGCGCAGC TGTGCTCGAC  
7560

GTTGTCACTG AAGCGGGAAG GGACTGGCTG CTATTGGGCG AAGTGCCGGG GCAGGATCTC  
7620

CTGTCATCTC ACCTTGCTCC TGCCGAGAAA GTATCCATCA TGGCTGATGC AATGCGGCGG  
7680

CTGCATACGC TTGATCCGGC TACCTGCCCCA TTCGACCACC AAGCGAAACA TCGCATCGAG  
7740

CGAGCACGTA CTCGGATGGA AGCCGGTCTT GTCGATCAGG ATGATCTGGA CGAAGAGCAT  
7800

CAGGGGCTCG CGCCAGCCGA ACTGTTCGCC AGGCTCAAGG CGCGCATGCC CGACGGCGAG  
7860

GATCTCGTCG TGACCCATGG CGATGCCTGC TTGCCGAATA TCATGGTGGA AAATGGCCGC  
7920

TTTTCTGGAT TCATCGACTG TGGCCGGCTG GGTGTGGCGG ACCGCTATCA GGACATAGCG  
7980

TTGGCTACCC GTGATATTGC TGAAGAGCTT GGC GGCGAAT GGGCTGACCG CTCCTCGTG  
8040

CTTTACGGTA TCGCCGCTCC CGATTGCGAG CGCATCGCCT TCTATCGCCT TCTTGACGAG  
8100

TTCTTCTGAG CGGGACTCTG GGGTTCGAAA TGACCGACCA AGCGACGCCC AACCTGCCAT  
8160

CACGAGATTT CGATTCCACC GCCGCCTTCT ATGAAAGGTT GGGCTTCGGA ATCGTTTTCC  
8220

GGGACGCCGG CTGGATGATC CTCCAGCGCG GGGATCTCAT GCTGGAGTTC TTCGCCCACC  
8280

CCAACTTGTT TATTGCAGCT TATAATGGTT ACAAATAAAG CAATAGCATC ACAAATTTCA  
8340

CAAATAAAGC ATTTTTTTCA CTGCATTCTA GTTGTGGTTT GTCCAAACTC ATCAATGTAT  
8400

CTTATCATGT CTGTATACCG TCGACCTCTA GCTAGAGCTT GGC GTAATCA TGGTCATAGC  
8460

TGTTTCCTGT GTGAAATTGT TATCCGCTCA CAATTCCACA CAACATACGA GCCGGAAGCA  
8520

TAAAGTGTA AGCCTGGGGT GCCTAATGAG TGAGCTAACT CACATTAATT GCGTTGCGCT  
8580

CACTGCCCCG TTTCCAGTCG GGAAACCTGT CGTGCCAGCT GCATTAATGA ATCGGCCAAC  
8640

GCGCGGGGAG AGGCGGTTTG CGTATTGGGC GCTCTTCCGC TTCCTCGCTC ACTGACTCGC  
8700

TGCGCTCGGT CGTTCGGCTG CGGCGAGCGG TATCAGCTCA CTCAAAGGCG GTAATACGGT  
8760

TATCCACAGA ATCAGGGGAT AACGCAGGAA AGAACATGTG AGCAAAAGGC CAGCAAAAGG  
8820

CCAGGAACCG TAAAAAGGCC GCGTTGCTGG CGTTTTTCCA TAGGCTCCGC CCCCCTGACG  
8880

AGCATCACAA AAATCGACGC TCAAGTCAGA GGTGGCGAAA CCCGACAGGA CTATAAAGAT  
8940

ACCAGGCGTT TCCCCCTGGA AGCTCCCTCG TGCGCTCTCC TGTTCCGACC CTGCCGCTTA  
9000

CCGGATACCT GTCCGCCTTT CTCCCTTCGG GAAGCGTGCG GCTTTCTCAA TGCTCACGCT  
9060

GTAGGTATCT CAGTTCGGTG TAGGTCGTTC GCTCCAAGCT GGGCTGTGTG CACGAACCCC  
9120

CCGTTCAGCC CGACCGCTGC GCCTTATCCG GTAACATATCG TCTTGAGTCC AACCCGGTAA  
9180

GACACGACTT ATCGCCACTG GCAGCAGCCA CTGGTAACAG GATTAGCAGA GCGAGGTATG  
9240

TAGGCGGTGC TACAGAGTTC TTGAAGTGGT GGCCTAACTA CGGCTACACT AGAAGGACAG  
9300

TATTTGGTAT CTGCGCTCTG CTGAAGCCAG TTACCTTCGG AAAAAGAGTT GGTAGCTCTT  
9360

GATCCGGCAA ACAAACCACC GCTGGTAGCG GTGGTTTTTT TGTTCGCAAG CAGCAGATTA  
9420

CGCGCAGAAA AAAAGGATCT CAAGAAGATC CTTTGATCTT TTCTACGGGG TCTGACGCTC  
9480

AGTGGAACGA AAACCTCACGT TAAGGGATTT TGGTCATGAG ATTATCAAAA AGGATCTTCA  
9540

CCTAGATCCT TTAAATTAA AAATGAAGTT TTAAATCAAT CTAAAGTATA TATGAGTAAA  
9600

CTTGGTCTGA CAGTTACCAA TGCTTAATCA GTGAGGCACC TATCTCAGCG ATCTGTCTAT  
9660

TTCGTTCATC CATAGTTGCC TGAATCCCCG TCGTGTAGAT AACTACGATA CGGGAGGGCT  
9720

TACCATCTGG CCCAGTGCT GCAATGATAC CGCGAGACCC ACGCTCACCG GCTCCAGATT  
9780

TATCAGCAAT AAACCAGCCA GCCGGAAGGG CCGAGCGCAG AAGTGGTCCT GCAACTTTAT  
9840

CCGCCTCCAT CCAGTCTATT AATTGTTGCC GGAAGCTAG AGTAAGTAGT TCGCCAGTTA  
9900

ATAGTTTGCG CAACGTTGTT GCCATTGCTA CAGGCATCGT GGTGTCACGC TCGTCGTTTG  
9960

GTATGGCTTC ATTCAGCTCC GGTCCCAAC GATCAAGGCG AGTTACATGA TCCCCCATGT  
10020

TGTGCAAAAA AGCGGTTAGC TCCTTCGGTC CTCCGATCGT TGTCAGAAGT AAGTTGGCCG  
10080

CAGTGTTATC ACTCATGGTT ATGGCAGCAC TGCATAATTC TCTTACTGTC ATGCCATCCG  
10140

TAAGATGCTT TTCTGTGACT GGTGAGTACT CAACCAAGTC ATTCTGAGAA TAGTGTATGC  
10200

GGCGACCGAG TTGCTCTTGC CCGGCGTCAA TACGGGATAA TACCGCGCCA CATAGCAGAA  
10260

CTTTAAAAGT GTCATCATT GGAAAACGTT CTTGGGGGCG AAAACTCTCA AGGATCTTAC  
10320

CGCTGTTGAG ATCCAGTTCG ATGTAACCCA CTCGTGCACC CAACTGATCT TCAGCATCTT  
10380

TTACTTTCAC CAGCGTTTCT GGGTGAGCAA AAACAGGAAG GCAAAATGCC GCAAAAAAGG  
10440

GAATAAGGGC GACACGGAAA TGTGAATAC TCATACTCTT CCTTTTCAA TATTATTGAA  
10500

GCATTTATCA GGGTTATTGT CTCATGAGCG GATACATATT TGAATGTATT TAGAAAAATA  
10560

AACAAATAGG GGTTCCGCGC ACATTTCCCC GAAAAGTGCC ACCTGACGTC  
10610

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

TGTACACCGG ATCCGGCGCA CACC24



## (2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 35 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

CACAACGAGC TCAATTAATT AATTGCCACA TCCTC  
35  
G

## (2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 4 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

Thr Leu Trp Thr

1

## (2) INFORMATION FOR SEQ ID NO:20:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 12 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

Pro	Ser	Ala	Ser	Ala	Ser	Ala	Ser	Ala	Pro	Gly	Ser
1				5						10	

We claim:

1. A packaging cell line expressing one or more adenovirus structural proteins, polypeptides, or fragments thereof, wherein said structural protein is selected from the group consisting of:
  - a. penton base;
  - b. hexon;
  - c. fiber;
  - d. polypeptide IIIa;
  - e. polypeptide V;
  - f. polypeptide VI;
  - g. polypeptide VII;
  - h. polypeptide VIII; and
  - i. biologically active fragments thereof.
2. A packaging cell line according to claim 1, which supports the production of a viral vector.
3. A packaging cell line according to claim 2, wherein the viral vector comprises a tripartite leader sequence or a sequence substantially homologous thereto.
4. A packaging cell line according to any one of the preceding claims wherein said structural protein is fiber and wherein said fiber protein has been modified to include a non-native amino acid residue sequence which targets a specific receptor, but which does not disrupt trimer formation or transport of fiber into the nucleus.
5. A packaging cell line according to claim 4 wherein said non-native amino acid residue sequence alters the binding specificity of the fiber for a targeted cell type.
6. A packaging cell line according to, any one of the preceding claims wherein said structural protein is fiber comprising amino acid residue sequences from more than one adenovirus serotype.
7. A packaging cell line according to any one of claims 2 to 6, wherein said viral vector includes a nucleic acid sequence having a deletion or mutation of a DNA sequence encoding an adenovirus structural protein, polypeptide, or fragment thereof.

8. A packaging cell line according to claim 7 wherein said viral vector includes a nucleic acid sequence having a deletion or mutation of the DNA sequences encoding regulatory polypeptides E1A and E1B.
9. A packaging cell line according to claim 7, wherein said viral vector further includes a nucleic acid sequence having a deletion or mutation of a DNA sequence encoding one or more of the following regulatory proteins or polypeptides: E2A, E2B, E3, E4, L4, or fragments thereof.
10. A packaging cell line according to claim 7, wherein a foreign DNA sequence encoding one or more foreign proteins, polypeptides or fragments thereof has been inserted in place of any of said deletions in said therapeutic viral vector.
11. A packaging cell line according to claim 10, wherein said foreign DNA encodes a tumor-suppressor protein or a biologically active fragment thereof.
12. A packaging cell line according to claim 10, wherein said foreign DNA encodes a suicide protein or a biologically active fragment thereof.
13. A packaging cell line according to any one of the preceding claims, wherein said cell line is an epithelial cell line.
14. A packaging cell line according to claim 12, wherein said cell line is selected from the group consisting of 293, A549, W162, HeLa, Vero, 211, and 211A cell lines.
15. A viral vector comprising deletion or mutation of a DNA sequence encoding an adenovirus structural protein, polypeptide, or fragment thereof, wherein said structural protein, polypeptide or fragment thereof is selected from the group consisting of:
  - a. penton base;
  - b. hexon;
  - c. fiber;
  - d. polypeptide IIIa;
  - e. polypeptide V;
  - f. polypeptide VI;
  - g. polypeptide VII;
  - h. polypeptide VIII; and
  - i. biologically active fragments thereof.

16. A viral vector according to claim 15, further comprising deletion or mutation of the DNA sequences encoding regulatory polypeptides E1A and E1B, or fragments thereof.
17. A viral vector according to claim 15, further comprising deletion or mutation of the DNA sequence encoding one or more of the following regulatory proteins or polypeptides: E2A, E2B, E3, E4, L4, or fragments thereof.
18. A viral vector according to claim 15 further comprising a foreign DNA sequence inserted in place of the DNA sequence encoding said structural protein, polypeptide, or fragment thereof.
19. A viral vector according to any one of claims 16, 17 or 18, further comprising foreign DNA sequences inserted in place of the DNA sequences encoding said regulatory polypeptides or fragments thereof.
20. A viral vector lacking all or part of a DNA sequence encoding adenovirus fiber protein, wherein said DNA sequence has been replaced by a foreign DNA sequence encoding a therapeutic molecule.
21. A viral vector having a mutation in a DNA sequence encoding adenovirus fiber protein.
22. A viral vector according to any one of 15 to 21 wherein the vector is a therapeutic vector.
23. A complementing plasmid comprising a promoter nucleotide sequence operatively linked to a nucleotide sequence encoding an adenovirus structural protein, polypeptide, or fragment thereof.
24. A complementing plasmid according to claim 23, further comprising a nucleotide sequence encoding:
  - a. a first adenovirus regulatory protein, polypeptide, or fragment thereof; or
  - b. a second regulatory protein, polypeptide, or fragment thereof; or
  - c. a third regulatory protein, polypeptide, or fragment thereof; or
  - d. any combination of the foregoing.
25. A complementing plasmid according to claim 23, wherein said adenovirus structural protein or polypeptide is selected from the group consisting of:
  - a. penton base;

- b. hexon;
  - c. fiber;
  - d. polypeptide IIIa;
  - e. polypeptide V;
  - f. polypeptide VI;
  - g. polypeptide VII;
  - h. polypeptide VIII; and
  - i. biologically active fragments thereof.
26. A complementing plasmid according to claim 24, wherein said regulatory proteins, polypeptides or fragments thereof are selected from the group consisting of E1A, E1B, E2A, E2B, E3, E4, and L4.
27. A complementing plasmid comprising a promoter nucleotide sequence operatively linked to a nucleotide sequence encoding an adenovirus regulatory protein, polypeptide, or fragment thereof.
28. A composition useful in the preparation of a therapeutic viral vectors, the composition comprising a cell containing a delivery plasmid comprising an adenovirus genome lacking a nucleotide sequence encoding fiber.
29. A composition according to claim 28, wherein said delivery plasmid further comprises a nucleotide sequence encoding a foreign polypeptide.
30. A composition according to claim 29, wherein said polypeptide is a therapeutic molecule.
31. A composition according to claim 28, wherein said delivery plasmid is selected from the group consisting of pDV1, p E1B gal, p E1sp1B, and pFG140-f.
32. A composition according to claim 28, wherein said cell further comprises a complementing plasmid containing a nucleotide sequence encoding fiber, said plasmid being stably integrated into the cellular genome of said cell.
33. A composition according to claim 32, wherein said complementing plasmid has the characteristics of pCLF having ATCC Accession Number 97737.
34. A composition useful in the preparation of therapeutic viral vectors, said composition comprising a cell containing:

- a. a first delivery plasmid comprising an adenovirus genome lacking a nucleotide sequence encoding fiber and incapable of directing the packaging of new viral particles in the absence of a second delivery plasmid; and
  - b. a second delivery plasmid comprising an adenoviral genome capable of directing the packaging of new viral particles in the presence of said first delivery plasmid.
- 35. A composition according to claim 34, wherein said first and second delivery plasmids interact within said cell to produce a therapeutic viral vector.
  - 36. A composition according to claim 34, wherein said cell further comprises a complementing plasmid containing a nucleotide sequence encoding fiber, said plasmid being stably integrated into the cellular genome of said cell.
  - 37. A composition according to claim 34, wherein said first or second delivery plasmid further comprises a nucleotide sequence encoding a foreign polypeptide.
  - 38. A composition according to claim 37, wherein said polypeptide is a therapeutic molecule.
  - 39. A composition according to claim 34, wherein said first delivery plasmid lacks adenovirus packaging signal sequences.
  - 40. A composition according to claim 34, wherein said second delivery plasmid contains a LacZ reporter construct.
  - 41. A composition according to claim 34, wherein said second delivery plasmid further lacks a nucleotide sequence encoding an adenovirus regulatory protein.
  - 42. A composition according to claim 41, wherein said regulatory protein is E1.
  - 43. A composition according to claim 36, wherein said complementing plasmid has the characteristics of pCLF having ATCC Accession Number 97737.
  - 44. A composition according to claim 34, wherein said first delivery plasmid lacks a nucleotide sequence encoding adenovirus E4 protein and said second delivery plasmid lacks a nucleotide sequence encoding adenovirus E1 protein.
  - 45. A composition according to claim 44, wherein said cell contains at least one complementing plasmid encoding an adenoviral regulatory protein and a structural protein.
  - 46. A composition according to claim 45, wherein said regulatory protein is E4 and said structural protein is fiber.

47. A composition according to claim 45, wherein said regulatory protein is E1 and said structural protein is fiber.
48. A composition according to claim 45, wherein said regulatory protein is both E1 and E4 and said structural protein is fiber.
49. A composition according to claim 45, wherein said adenoviral regulatory protein and said structural protein are encoded by separate complementing plasmids.
50. A composition according to claim 45, wherein said cell is selected from the group consisting of 293, A549, W162, HeLa, Vero, 211, and 211A.






Figure 1




Figure 2




FIGURE 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8






Figure 9



FIGURE 10




Figure 11




FIGURE 12




Figure 13




Figure 14

A

Ad2/5	11	K	R	A	R	P	S	E	D	T	F	N	P	V	P	Y	D	T	E	T	G	P	P	T	V	P	F	L	T	P	P	F	V	S	P	N	G	F	O	E	S	P	P	
Ad3	11	A	K	R	A	R	L	S	-	T	S	F	N	P	V	P	Y	E	D	S	S	-	Q	H	P	F	I	M	P	Q	F	I	S	P	D	G	F	T	O	S	P	N		
Ad18/37	11	S	K	R	L	R	V	E	D	-	F	N	P	V	P	V	G	Y	A	R	N	-	N	I	P	F	L	T	P	P	F	V	S	S	D	G	F	K	N	F	P	P		
Ad40 (1)	11	-	K	R	A	R	F	-	E	D	-	F	N	P	V	P	Y	E	H	Y	N	-	P	L	D	I	P	F	I	T	P	P	F	A	S	S	N	G	L	O	E	K	P	P
Ad40 (2)	11	-	K	R	T	R	-	I	E	D	-	F	N	P	V	P	Y	D	T	S	S	T	P	S	-	I	P	Y	V	A	D	P	F	V	S	S	D	G	L	O	E	N	P	P

B




Figure 15




Figure 16






## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> : C12N 15/34, 15/86, 15/85, 5/10, C07K 14/075, A61K 48/00</p>	<p>A2</p>	<p>(11) International Publication Number: <b>WO 98/13499</b></p> <p>(43) International Publication Date: 2 April 1998 (02.04.98)</p>
<p>(21) International Application Number: PCT/EP97/05251</p> <p>(22) International Filing Date: 24 September 1997 (24.09.97)</p> <p>(30) Priority Data: 08/719.806 25 September 1996 (25.09.96) US</p> <p>(71) Applicants (for all designated States except US): NOVARTIS AG (CH/CH); Schwarzwaldallee 215, CH-4058 Basel (CH). THE SCRIPPS RESEARCH INSTITUTE (US/US); 10666 North Torrey Pines Road, La Jolla, CA 92037 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): NEMEROW, Glen, R. [US/US]; 462 Cerro Street, Encinitas, CA 92024 (US). VON SEGGERN, Daniel, J. [US/US]; 7528 Charmant Drive #428, San Diego, CA 92122 (US).</p> <p>(74) Agent: ROTH, Bernhard, M.; Novartis AG, Patent- und Markenabteilung, Lichtstrasse 35, CH-4002 Basel (CH).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> Without international search report and to be republished upon receipt of that report.</p>

(54) Title: PACKAGING CELL LINES FOR USE IN FACILITATING THE DEVELOPMENT OF HIGH-CAPACITY ADENOVIRAL VECTORS



## (57) Abstract

The present invention relates to gene therapy, especially to adenovirus-based gene therapy, and related cell lines and compositions. In particular, novel packaging cell lines are disclosed, for use in facilitating the development of high-capacity vectors. The invention also discloses a variety of high-capacity adenovirus vectors and related compositions and kits including the disclosed cell lines and vectors. Finally, the invention discloses methods of preparing and using the disclosed vectors, cell lines and kits.

**FOR THE PURPOSES OF INFORMATION ONLY**


Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12N 15/34, 15/86, 15/85, 5/10, C07K 14/075, A61K 48/00</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 98/13499</b> <b>(43) International Publication Date:</b> 2 April 1998 (02.04.98)
<b>(21) International Application Number:</b> PCT/EP97/05251 <b>(22) International Filing Date:</b> 24 September 1997 (24.09.97) <b>(30) Priority Data:</b> 08/719,806 25 September 1996 (25.09.96) US <b>(71) Applicants (for all designated States except US):</b> NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH). THE SCRIPPS RESEARCH INSTITUTE [US/US]; 10666 North Torrey Pines Road, La Jolla, CA 92037 (US). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> NEMEROW, Glen, R. [US/US]; 462 Cerro Street, Encinitas, CA 92024 (US). VON SEGGERN, Daniel, J. [US/US]; 7528 Charmant Drive #428, San Diego, CA 92122 (US). <b>(74) Agent:</b> ROTH, Bernhard, M.; Novartis AG, Patent- und Markenabteilung, Lichtstrasse 35, CH-4002 Basel (CH).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). <b>Published</b> <i>With international search report.</i> <b>(88) Date of publication of the international search report:</b> 27 August 1998 (27.08.98)

**(54) Title:** PACKAGING CELL LINES FOR USE IN FACILITATING THE DEVELOPMENT OF HIGH-CAPACITY ADENOVIRAL VECTORS

**(57) Abstract**

The present invention relates to gene therapy, especially to adenovirus-based gene therapy, and related cell lines and compositions. In particular, novel packaging cell lines are disclosed, for use in facilitating the development of high-capacity vectors. The invention also discloses a variety of high-capacity adenovirus vectors and related compositions and kits including the disclosed cell lines and vectors. Finally, the invention discloses methods of preparing and using the disclosed vectors, cell lines and kits.

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## INTERNATIONAL SEARCH REPORT

Intern: Application No

PCT/EP 97/05251

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/34 C12N15/86 C12N15/85 C12N5/10 C07K14/075  
A61K48/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	KROUGLIAK V ET AL: "DEVELOPMENT OF CELL LINES CAPABLE OF COMPLEMENTING E1, E4, AND PROTEIN IX DEFECTIVE ADENOVIRUS TYPE 5 MUTANTS" HUMAN GENE THERAPY, vol. 6, no. 12, 1 December 1995, pages 1575-1586, XP000575816 see the whole document	23,24, 26,27
X	--- CARAVOKYRI C ET AL: "CONSTITUTIVE EPISOMAL EXPRESSION OF POLYPEPTIDE IX (PIX) IN A 293-BASED CELL LINE COMPLEMENTS THE DEFICIENCY OF PIX MUTANT ADENOVIRUS TYPE 5" JOURNAL OF VIROLOGY, vol. 69, no. 11, 1 November 1995, pages 6627-6633, XP000561195 see the whole document --- -/-	23,24



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*Z\* document member of the same patent family

Date of the actual completion of the international search

2 March 1998

Date of mailing of the international search report

27.05.98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

HORNIG H.

## INTERNATIONAL SEARCH REPORT

Intern 1st Application No

PCT/EP 97/05251

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 95 02697 A (RHONE POULENC RORER SA ;PERRICAUDET MICHEL (FR); VIGNE EMMANUELLE) 26 January 1995 see the whole document ---	23,24,27
X	WO 95 11984 A (CANJI INC) 4 May 1995 cited in the application see the whole document ---	23,24,27
X	WO 95 27071 A (UNIV TEXAS) 12 October 1995  see page 33, line 4 - page 35, line 2 ---	1,2, 15-19
X	WO 95 34671 A (GENVEC INC ;KOVESDI IMRE (US); BROUGH DOUGLAS E (US); MCVEY DUNCAN) 21 December 1995 see claims 4,13 ---	1,2,15, 27
X	WO 96 14061 A (CELL GENESYS INC) 17 May 1996 see the whole document ---	27
X	WO 96 22378 A (RHONE POULENC RORER SA ;DEDIEU JEAN FRANCOIS (FR); LATTA MARTINE ( ) 25 July 1996 see the whole document ---	27
A	WO 96 07734 A (GENVEC INC) 14 March 1996  see the whole document ---	1-3, 7-19, 22-27
P,X	WO 97 21826 A (GENVEC INC) 19 June 1997 see page 48, line 25 - page 49, line 25; figure 8 -----	27

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP 97/05251

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
  
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see continuation-sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
  
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
  
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
  
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-3,7-19,22,25 partially and 23,24,26,27 complete (subject 1. on continuation-sheet)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: (1-3,7-19,22,25)-partially, (23,24,26,27)-complete

A packaging cell line expressing one or more adenovirus structural proteins, polypeptides, or fragments thereof, wherein at least one of said structural protein is p e n t o n b a s e ; and biologically active fragments thereof; said cell line, which supports the production of a viral vector; a complementing plasmid comprising a promoter nucleotide sequence operatively linked to a nucleotide sequence encoding an adenoviral structural protein, polypeptide, or fragment thereof; said plasmid wherein said structural protein, or polypeptide is p e n t o n b a s e ; a complementing plasmid comprises a promoter nucleotide operatively linked to a nucleotide sequence encoding an adenoviral regulatory protein, polypeptide, or fragment thereof;

2. Claims: (1-3,7-19,22,25)-partially

A packaging cell line expressing one or more adenovirus structural proteins, polypeptides, or fragments thereof, wherein at least one of said structural protein is h e x o n , and biological active fragments thereof; said cell line, which supports the production of a viral vector; a complementing plasmid comprising a promoter nucleotide sequence operatively linked to a nucleotide sequence encoding an adenovirus structural protein, polypeptide, or fragment thereof, wherein said structural protein, or polypeptide is h e x o n ;

3. Claims: (1-3,7-19,22,25)-partially, (4-6,20,21, 28-50)-complete

Idem as subject 2, but limited to the structural polypeptide f i b e r ; a composition useful in the preparation of a therapeutic viral vectors, the composition comprising a cell containing a delivery plasmid comprising an adenoviral genome lacking a nucleotide sequence encoding f i b e r ; a composition useful in the preparation of therapeutic viral vectors, said composition comprising a cell containing: a. a first delivery plasmid comprising an adenovirus genome lacking nucleotide sequences encoding fiber and incapable of directing the packaging of new viral particles in the absence of a second delivery plasmid; and b. a second delivery plasmid comprising an adenoviral genome capable of directing the packaging of new viral particles in the presence of the first delivery plasmid;

4. Claims: (1-3,7-19,22,25)-partially

Idem as subject 2, but limited to the structural polypeptide IIIa;



FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

5. Claims: (1-3,7-19,22,25)-partially

Idem as subject 2, but limited to the structural polypeptide V;

6. Claims: (1-3,7-19,22,25)-partially

Idem as subject 2, but limited to the structural polypeptide VI;

7. Claims: (1-3,7-19,22,25)-partially

Idem as subject 2, but limited to the structural polypeptide VII;

8. Claims: (1-3,7-19,22,25)-partially

Idem as subject 2, but limited to the structural polypeptide VIII;

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern: al Application No

PCT/EP 97/05251

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9502697 A	26-01-95	FR 2707664 A	20-01-95
		FR 2718749 A	20-10-95
		AU 7264694 A	13-02-95
		CA 2144040 A	26-01-95
		CN 1113390 A	13-12-95
		CZ 9500639 A	15-11-95
		EP 0667912 A	23-08-95
		FI 951138 A	13-04-95
		HU 72558 A	28-05-96
		JP 8501703 T	27-02-96
		NO 950939 A	10-03-95
		NZ 269156 A	26-03-96
		PL 308122 A	24-07-95
		SK 31295 A	08-05-96
		ZA 9405012 A	20-02-95
WO 9511984 A	04-05-95	US 5591209 A	07-01-97
		AU 687117 B	19-02-98
		AU 8125094 A	22-05-95
		BR 9407956 A	26-11-96
		CA 2173975 A	04-05-95
		CZ 9601143 A	13-11-96
		EP 0797676 A	01-10-97
		FI 961755 A	04-06-96
		JP 9507051 T	15-07-97
		NO 961639 A	24-06-96
		NZ 275956 A	22-09-97
		PL 314239 A	02-09-96
		SK 51096 A	01-10-96
		AU 2637295 A	18-12-95
		WO 9532020 A	30-11-95
		US 5534015 A	09-07-96
WO 9527071 A	12-10-95	AU 2238695 A	23-10-95
		EP 0754237 A	22-01-97
WO 9534671 A	21-12-95	AU 2770495 A	05-01-96
		CA 2192442 A	21-12-95
		EP 0784690 A	23-07-97

## INTERNATIONAL SEARCH REPORT

information on patent family members

Internat'l Application No

PCT/EP 97/05251

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9614061 A	17-05-96	AU 4109296 A CA 2204357 A EP 0797436 A	31-05-96 17-05-96 01-10-97
WO 9622378 A	25-07-96	FR 2729674 A FR 2741891 A FR 2738575 A AU 4544396 A CA 2210168 A CZ 9702306 A EP 0805868 A FI 973055 A NO 973332 A	26-07-96 06-06-97 14-03-97 07-08-96 25-07-96 15-10-97 12-11-97 18-07-97 18-07-97
WO 9607734 A	14-03-96	US 5559099 A AU 684220 B AU 3212995 A CA 2198861 A EP 0778889 A US 5712136 A US 5731190 A	24-09-96 04-12-97 27-03-96 14-03-96 18-06-97 27-01-98 24-03-98
WO 9721826 A	19-06-97	AU 1417797 A	03-07-97